LICK OBSERVATORY

UNIVERSITY OF CALIFORNIA

BULLETIN No. 6

OBSERVATIONS OF THE SPECTROSCOPIC BINARY CAPELLA

The first magnitude star Capella was discovered to be a spectroscopic double star early in August, 1899, from an examination of the plates of its spectrum secured with the MILLS Spectrograph in 1896. Announcement of the fact was made to the Astronomical and Astrophysical Society of America at the meeting of September 7, 1899, and in the Astrophysical Journal for October, 1899.

Independent discovery of its binary character was made by Mr. H. F. NEWALL of Cambridge, England, in November, 1899, and announced in the Monthly Notices of the Royal Astronomical Society for November.

The spectra of the two components are distinguishable on most of the plates—the exceptions being those taken when the radial velocities of the two were nearly equal, producing a superposition of the two sets of lines. The spectrum of the principal star is of the solar type, whereas that of the secondary is intermediate between the solar and Sirian types.

The velocities of the principal component, as observed with the MILLS Spectrograph, are given in the following table:

No.	DATE. Greenwich M.T.	VELOCITY. Kilometers.
1	1896 September 1.036, . . .	+ 36.4 C
2	17.005, . . .	53.8 C
3	October 4.003, . . .	50.3 C
4	6.029, . . .	46.9 C
5	November 12.865, . . .	4.2 C
6	1899 August 12.999, . . .	48.3 C
7	27.052, . . .	26.1 C
8	September 12.950, . . .	5.7 W
9	20.006, . . .	5.1 W
10	20.919, . . .	3.5 W
11	20.933, . . .	5.4 W
12	25.909, . . .	6.6 C
13	October 3.988, . . .	14.8 C
14	16.912, . . .	32.9 C
15	16.929, . . .	32.0 C
16	31.892, . . .	52.0 C

No.	DATE. Greenwich M.T.	VELOCITY. Kilometers.
17	1899 November 6.026, . . .	+ 54.8 C
18	27.952, . . .	43.2 W
19	27.966, . . .	44.0 W
20	December 3.730, . . .	35.2 C
21	18.648, . . .	12.6 W
22	24.882, . . .	7.7 W
23	1900 January 10.649, . . .	7.7 C
24	21.740, . . .	21.7 W
25	February 11.724, . . .	50.0 W
26	26.726, . . .	55.2 W
27	26.740, . . .	54.8 C
28	August 2.012, . . .	3.6 C
29	September 19.944, . . .	55.5 C
30	24.950, . . .	53.9 C
31	27.005, . . .	53.7 C

[Measures of the plates by CAMPBELL and WRIGHT are indicated by C and W respectively.]

The presence of the second component's spectrum interferes considerably with the measures of that of the first component, and the probable error of a single observation, ± 0.50 kilometer, deduced by Dr. REESE, is as small as could be expected. Measures of the speed of the second component are somewhat uncertain, but an estimated range of from -3 to $+63$ kilometers will not be far from the truth. The velocity of the principal component in the line of sight ranges from $+4.2$ to $+55.7$ kilometers. The masses of the two components are therefore as 1.26 to 1.

The solar-type component is estimated to be half a magnitude brighter, photographically, than the bluer component. In the visual portion of the spectrum the solar component is probably at least a whole magnitude the brighter of the two.

Inasmuch as the spectroscope takes account of the component of speed in the line of sight, and is powerless to measure the component at right angles to the line of sight, the spectroscopic orbit is determinate in