LICK OBSERVATORY BULLETIN

NUMBER 80

THE ORBIT OF ξ SCORPII = Σ 1998

This interesting binary system was placed on my regular observing list in 1897. Since then the following measures have been secured:

1898.17	222°6	0'84	3 ⁿ	12-in.	A. N. 150 , 140, 1899.
1899.35	228.4	0.70	4-3	12-in.	A. N. 152 , 230, 1900.
1901.47	242.2	0.50	3	36-in.	L. O. B. 1 , 73, 1901.
1903.448	269.1	0.34	1000 power	36-in.	Unpublished.
.495	269.1	0.35	1600	36	
.599	273.2	0.28	1500	36	
1903.51	270°5	0'32			
1904.269	296.8	0.22	1500	36-in.	Unpublished.
.532	309.6	0.20	1500	36	
1904.40	303°2	0'21			
1905.288	358.2	0.22	1500	36-in.	Unpublished.
.310	11.7	0.22	1500	36	
1905.30	5°0	0'22			

These observations when compared with See's* elements, which closely resemble those obtained by Schorr† in 1889, give the following residuals [O-C]:

1898.17	+	1°2	+0'03
1899.35	+	3.3	-0.05
1901.47	+	9.9	-0.19
1903.51	+	27.9	-0.29
1904.40	+	55.3	-0.37
1905.30	+	100.9	-0.34

The first three residuals indicated that the commonly accepted orbit needed revision, but the last three show that it is radically in error. This will be even more evident if the observations are plotted on the diagram of the apparent orbit given by See. It will at once be seen that the true orbit of this binary, instead of being nearly circular, is highly eccentric; and that the revolution period, instead of being over one hundred years, cannot exceed half that value.

So far as the periodic time is concerned, this

result has been anticipated by Thiele, who in 1859 computed an orbit for this pair with a period of forty-nine years. His other elements, however, were in error, and as his orbit failed to represent later observations it has been generally disregarded.

I have computed a new orbit by the graphical method using the observations to 1895 inclusive, as given by Dr. See,* and, besides my own given above, the following by Doolittle:

1899.42	227°2	0'81	6 ⁿ
1900.54	232.3	0.74	5 ⁿ

To do this it was necessary to change by 180° the position angles for the years 1782 to 1858 inclusive, or those from 1862 to date. I have adopted the former alternative. The two components are so nearly equal in brightness that no objection can be made to this procedure.

My elements are:

P =	44.5 years
T =	1905.4
e =	0.767
a =	0'701
Ω =	20°4
i =	± 29.1
ω =	352.6
Position angles increasing.	

APPARENT ORBIT.

Length of major axis	= 1'464
Length of minor axis	= 0'802
Angle of major axis	= 11°0
Angle of periastron	= 13°5
Distance of star from centre	= 0'561

EPHEMERIS.

1905.4	12°3	0'16
1906.4	83.1	0.21
1907.4	117.7	0.31
1908.4	135.1	0.41
1909.4	145.6	0.51
1910.4	153.0	0.60

* Evolution of the Stellar Systems 1, 176, 1896.

† Untersuchungen über ξ Scorpii. München, 1889.

The diagram reproduces the apparent orbit and the more important observations, combined into annual means, from 1825 to 1905. The distance

measures at apastron range from $0''.70$ to $1''.54$ and many are necessarily poorly represented. The measures of Dembowski, Schiaparelli, Hall and other good observers from 1875 to 1886 agree with the earlier ones of Struve, Dawes, and Kaiser in proving that the maximum separation of the two components is about $1''.25$. This distance, in connection with the angle measures, also satisfies the law of areas. The table which follows gives all the observations used and the residuals (O-C) resulting from their comparison with the above elements.

The present orbit is considered as preliminary and is published at this time more particularly to call the attention of double star observers to ξ Scorpil in the hope that numerous and careful observations may be secured in the next few years while the companion is moving rapidly from periastron to the position it occupied in 1867. Observations in this part of the orbit are almost entirely wanting and will be of great value in correcting the elements now given, especially the inclination. It is hardly profitable at

this time, therefore, to try to reduce further the residuals of the table, or to take account of the possible disturbing influence of the third star of the system or of the systematic errors of the observers. The residuals, as they stand, show that the orbit given above is approximately correct; in three or four years, when the necessary data are available, I hope to undertake a more complete discussion.

OBSERVATIONS AND RESIDUALS.

t	θ_0	ρ_0	n	Observer	$\Delta\theta$	(O-C) $\Delta\rho$
1782.36	188°0	...	1	Herschel	+14.5	...
1825.47	175.3	1.15	3	Struve	+ 5.5	+0.27
1828.48	181.0	...	1	Herschel	+ 4.0	...
1830.25	181.4	1.46	4-3	Herschel	+ 0.8	+0.36
1831.38	189.4	1.32	2-1	Herschel	+ 6.8	+0.19
1831.48	183.5	1.21	1	Struve	+ 2.5	+0.07
1832.52	184.8	1.24	1	Struve	+ 0.3	+0.09
1833.37	185.0	1.19	1	Struve	- 0.9	+0.01
1833.39	186.2	1.15	1	Dawes	- 2.0	-0.03
1834.45	188.3	1.24	2-1	Herschel	+ 0.6	+0.04
1834.45	186.7	1.24	1	Struve	- 1.0	+0.04
1834.50	187.1	1.17	4	Dawes	- 0.8	-0.03
1834.51	194.6	...	3	Mädler	- 6.8	...
1835.39	190.6	1.58	5-1	Herschel	+ 1.5	+0.37
1835.48	191.0	...	4	Mädler	+ 1.7	...
1836.49	189.5	1.02	1	Dawes	- 1.3	-0.20
1836.50	191.0	...	3	Mädler	+ 0.2	...
1837.33	191.4	...	1	Herschel	- 0.6	...
1839.61	196.7	1.28	2	Dawes	+ 1.3	+0.04
1840.56	198.6	1.19	3	Dawes	+ 1.9	-0.04
1840.57	197.2	0.96	1	O. Struve	+ 0.5	-0.27
1841.48	196.7	1.28	4-3	Mädler	- 1.3	+0.05
1841.57	200.8	0.84	1	O. Struve	+ 2.7	-0.38
1841.58	199.0	1.20	3-2	Dawes	+ 0.9	-0.03
1841.61	197.7	1.30	2-1	Kaiser	- 0.5	+0.07
1842.42	200.4	1.05	4-2	Mädler	+ 0.9	-0.16
1842.46	201.6	...	2	Dawes	+ 2.1	...
1842.53	201.0	...	1	Kaiser	+ 1.5	...
1843.40	203.5	1.09	2	Dawes	+ 2.5	-0.11
1843.40	203.8	1.16	6-4	Mädler	+ 2.8	-0.04
1843.62	200.8	1.20	11-1	Kaiser	- 0.5	± 0.00
1844.40	203.7	1.82	3	Mädler	+ 1.1	+0.64
1846.17	203.2	1.00	3-1	Jacob	- 2.3	-0.14
1846.47	204.1	0.97	9-8	Mitchell	- 1.9	-0.15
1847.58	206.0	1.71	1	Mitchell	- 2.0	+0.61
1848.54	210.6	1.19	3	Dawes	+ 0.8	+0.13
1848.54	207.2	0.84	1	Mitchell	- 2.6	-0.26
1853.53	226.3	...	1	Dawes	+ 3.5	...
1855.36	228.2	...	4	Dembowski	- 2.1	...
1855.53	233.1	0.46	3	Secchi	+ 2.0	-0.19

<i>t</i>	θ_0	ρ_0	<i>n</i>	Observer	(O-C)		<i>t</i>	θ_0	ρ_0	<i>n</i>	Observer	(O-C)	
					$\Delta\theta$	$\Delta\rho$						$\Delta\theta$	$\Delta\rho$
1856.20	245.5	0.63	3	Jacob	+10.5	+0.04	1879.41	189.2	1.22	5	Howe	+0.8	+0.02
1856.41	238.1		4	Dembowski	+1.6		1879.42	186.7	1.44	3	Stone	-1.7	+0.24
1856.49	250.3	0.36	10-8	Secchi	+13.3	-0.20	1879.47	187.6	1.45	3	Egbert	-0.9	+0.25
1856.55	239.6		2	Winnecke	+2.2		1879.54	189.8	1.07	3	Hall	+1.1	-0.13
1856.58	249.8	0.47	1	O. Struve	+12.3	-0.10	1879.56	186.8	1.29	7	Schiaparelli	-1.9	+0.09
							1879.58	185.6	1.47	2	C. W. Pritchett	-3.1	+0.27
1857.68	261.4	0.50	1	Jacob	+14.1	+0.05	1879.60	188.8	1.16	3	Burnham	$\pm$ 0.0	-0.04
							1879.67	194.5	0.70	3-1	Sea. & Smith	+4.6	-0.50
1858.13	259.4	0.40	1	Jacob	+6.4	$\pm$ 0.00	1880.36	188.8	1.12	2	Egbert	-1.1	-0.10
1858.22	296.8	0.30	1	Jacob	+42.6	-0.09	1880.40	189.7	1.17	4-2	Doberck	-0.2	-0.05
							1880.52	185.7	1.13	1	Frisby	-4.4	-0.09
1862.56	137.9		3	Dembowski	+29.1		1880.54	189.0	1.24	6	Schiaparelli	-1.2	+0.02
							1880.87	189.6	1.10	3	H. S. Pritchett	-1.0	-0.12
1863.44	142.1		9	Dembowski	+13.7								
1864.45	147.8	0.21	4	Secchi	+6.3	-0.25	1881.24	191.3	1.03	1	Doberck	+0.2	-0.20
1864.51	150.9		10	Dembowski	+8.9		1881.40	190.8	1.21	2-1	Bigourdan	-0.6	-0.02
1865.44	151.4		10	Dembowski	+1.4		1882.27	193.6	1.19	1	Doberck	+0.9	-0.04
1865.51	155.5	0.35	7	Secchi	+5.1	-0.21	1882.43	196.7	1.44	1	Englemann	+3.8	+0.21
1865.55	166.9	0.49	7	Englemann	+16.2	-0.07	1882.46	192.7	1.12	3	Hall	-0.2	-0.11
							1882.54	191.8	1.31	5	Schiaparelli	-1.3	+0.08
1866.46	156.6	0.53	8-3	Dembowski	+0.2	-0.11	1882.59	192.1	1.35	3	Frisby	-1.0	+0.12
1866.52	161.0	0.40	2-1	Secchi	+5.5	-0.25							
							1883.45	191.0	1.33	4	Frisby	-3.4	+0.09
1867.45	160.7	0.83	7-4	Dembowski	-0.2	+0.11	1883.49	195.3		1	Küstner	+0.8	
							1883.49	195.5	1.20	3	Englemann	+1.0	-0.04
1868.40	165.0	0.90	7-4	Dembowski	+0.3	+0.12	1883.51	193.9	1.38	2	Hall	-0.6	+0.14
1868.48	166.5	0.99	1	Knott	+5.4	+0.20	1883.52	191.5	1.16	3	Perrotin	-3.0	-0.08
							1883.55	193.5	1.24	12	Schiaparelli	-1.1	$\pm$ 0.00
1869.51	172.5	0.83	6	Dunér	+4.1	-0.02							
1869.52	168.2	0.88	5	Dembowski	-0.2	+0.03	1884.38	195.8	1.34	4-3	H. C. Wilson	$\pm$ 0.0	+0.11
							1884.44	195.6	1.46	3	Englemann	-0.3	+0.23
1870.21	168.2		1	Gledhill	-2.3		1884.50	194.6	1.28	5	Hall	-1.4	+0.05
1870.39	169.8	0.89	7-5	Dembowski	-1.2	-0.01	1884.53	195.1	1.27	3	Perrotin	-0.9	+0.04
1870.54	173.3	0.88	2	Dunér	+1.9	-0.03	1884.54	195.6	1.41	1	H. S. Pritchett	-0.5	+0.18
							1884.54	195.0	1.26	9	Schiaparelli	-1.1	+0.03
1871.41	173.1	1.06	7-5	Dembowski	-0.5	+0.11	1885.53	196.2	1.34	8	Schiaparelli	-1.1	+0.11
1871.49	174.0	1.00	1	Gledhill	+0.2	+0.04	1885.57	198.1	1.38	5	Englemann	+0.7	+0.15
1871.60	174.8	0.88	5	Dunér	+0.8	-0.08							
							1886.35	197.4	1.19	1	H. C. Wilson	-1.2	-0.03
1872.45	177.3	0.95	1	W. & S.	+1.4	-0.06	1886.46	197.5	1.24	2	Perrotin	-1.2	+0.02
1872.46	176.9	1.12	1	Knott	+1.0	+0.11	1886.49	198.6	1.54	2-1	Smith	-0.2	+0.32
1872.46	173.8	1.12	8-5	Dembowski	-2.2	+0.11	1886.51	198.1	1.29	3	Tarrant	-0.7	+0.07
1872.50	175.8	1.10	2	Ferrari	-0.2	+0.09	1886.55	197.2	1.19	3	Schiaparelli	-1.7	-0.03
1872.53	177.4	0.96	3	Dunér	+1.3	-0.05	1886.56	198.0	1.07	3	Hall	-0.9	-0.15
							1886.63	198.9	1.07	7	Englemann	-0.1	-0.15
1873.36	180.4	1.04	1	W. & S.	+0.5	$\pm$ 0.00							
1873.36	176.8	1.19	5-3	Dembowski	-2.1	+0.15	1887.54	199.6	1.16	9	Schiaparelli	-0.9	-0.05
1873.68	176.5	1.10	1	Gledhill	-2.0	+0.05							
							1888.50	200.4	1.24	2	Leavenworth	-1.5	+0.05
1874.44	183.1	1.19	1	W. & S.	+2.7	+0.10	1888.56	200.6	0.96	2	Hall	-1.4	-0.23
1874.49	178.7	1.05	5	Dembowski	-1.4	-0.04	1888.57	201.9	1.14	7	Schiaparelli	-0.1	-0.05
1875.44	180.5	1.10	5	Dembowski	-1.4	-0.02	1889.43	197.5	1.20	2	Hodges	-5.9	+0.03
1875.51	182.0	1.18	5	Schiaparelli	+0.1	+0.06							
1875.51	180.0	1.33	1	W. & S.	-1.9	+0.21	1890.39	205.2		2	Glaserapp	+0.2	
1875.56	180.9	0.96	4	Dunér	-1.1	-0.16							
							1891.46	200.6	1.27	2	Collins	-6.3	+0.16
1876.44	185.6	1.04	1	Howe	+2.0	-0.11	1891.48	208.7	2.87	1	See	+2.7	+1.76
1876.45	181.8	1.21	6	Dembowski	-1.8	+0.06							
1876.52	183.9	1.14	3	Hall	+0.2	-0.01	1892.53	208.0	1.23	3	Maw	-0.9	+0.15
1876.52	183.6	1.18	4	Schiaparelli	-0.1	+0.03	1892.58	206.5	0.82	4	Comstock	-2.5	-0.26
1876.54	182.5	1.0 $\pm$ 1	1	Plummer	-1.3	-0.15							
1876.61	186.5		3	Doberck	+2.6		1893.46	211.1	1.01	2	Burnham	+0.5	-0.03
							1893.49	209.5	1.10	1	Schiaparelli	-1.2	+0.06
1877.43	179.5	0.97	2-1	Doberck	-5.6	-0.20	1893.51	210.9	0.89	2	Leavenworth	+0.2	-0.15
1877.43	183.3	1.20	5	Dembowski	-1.8	+0.03	1893.60	209.7	1.07	5	Bigourdan	-1.2	+0.03
1877.43	184.1	1.61	1	Upton	-1.0	+0.44							
1877.46	184.9	1.27	1	W. & S.	-0.2	+0.07	1894.59	207.5	1.0 $\pm$ 2-1	2-1	Glaserapp	-5.7	$\pm$ 0.00
1877.47	187.0	1.12	4-1	Howe	+1.9	-0.05							
1877.55	184.0	1.25	9	Schiaparelli	-1.3	+0.08	1895.31	210.3	1.04	3	See	-4.6	+0.08
1877.55	182.5	1.27	3	Jedrzejewicz	-2.8	+0.10	1895.41	213.9	0.91	2	Schiaparelli	-1.4	-0.05
							1895.53	213.4	0.81	3	Comstock	-2.0	-0.14
1878.46	186.2	1.22	5-4	Dembowski	-0.7	+0.03							
1878.54	186.1	1.31	6	Schiaparelli	-1.0	+0.12	1898.17	222.6	0.84	3	Aitken	-0.5	+0.05

t	θ_0	ρ_0	n	Observer	(O—C)	
					$\Delta\theta$	$\Delta\rho$
1899.35	228°4	0".70	4—3	Aitken	+ 0".6	±0".00
1899.42	227.2	0.81	6	Doolittle	— 0.8	+0.11
1900.54	232.3	0.74	5	Doolittle	— 1.7	+0.13
1901.47	242.2	0.70	3	Aitken	+ 1.6	—0.02
1903.51	270.5	0.32	3	Aitken	+ 0.5	+0.01

t	θ_0	ρ_0	n	Observer	(O—C)	
					$\Delta\theta$	$\Delta\rho$
1904.40	303°2	0".21	2	Aitken	+ 1".6	—0".01
1905.30	5.0	0.22	2	Aitken	+ 0.1	+0.06

May 2, 1905.

R. G. AITKEN.