UNIVERSITY OF CALIFORNIA PUBLICATIONS

ASTRONOMY

LICK OBSERVATORY BULLETIN

NUMBER 183

NOTES ON AN ATTEMPT TO DETECT POLARIZATION IN THE BRIGHT
H γ LINE IN THE SPECTRUM OF α CETI.

The maximum of α Ceti (*Mira*) for the year 1909 occurred in August and a number of observations were made at that time for the purpose of determining, if possible, whether the light of the bright H γ line is polarized in any way. When it was found in 1898 that this line is triple during a certain part of the star's period¹ it was at once suspected that the observed phenomenon might be the result of magnetic activity in the star, but the necessary apparatus for making polariscopic observations was not available. Later, both in a letter to Dr. Campbell and in published articles,² Miss Agnes M. Clerke made suggestions along this line. Polariscopic observations were therefore planned for the maximum of 1899, but on this occasion the star was disappointingly faint, and the observations could not be undertaken with any hope of success. A number of direct photographs of the line were secured, however, showing it to be double, instead of triple, the more refrangible component being perhaps a little brighter than the other. The detection by Professor Hale of the Zeeman effect in sun-spots added interest to this subject, and indicated the importance of investigating it as fully as possible.

The observations described below do not shed much light on the question as to the existence of magnetic forces in the star. The complete solution of the problem is hedged about with difficulties, chief of which is the fact that if the direction of the magnetic field varies considerably in the source the characteristic polarizations of the various components will be destroyed. If the magnetic field maintains a constant direction throughout the source, then we

should expect to find plane polarization of the central component and elliptic polarization of the other two. All that is claimed for these observations is that they show pretty conclusively that the multiple character of the lines is not due to the action of such a uniform field.

About thirty photographic exposures were made, of which eleven are selected as of interest. In all cases the H γ line was observed, being photographed with the Mills spectrograph, mounted as described by Dr. Campbell³ and attached to the thirty-six inch telescope. The short camera (focal length 405.5 mm.) was used.

In one set of experiments a Nicol was placed in front of the slit, and so oriented that the plane of polarization of its transmitted light was parallel to the slit of the spectrograph. On account of the polarizing action of a spectroscope this is the best position for the Nicol. The entire apparatus,—spectrograph and Nicol,—could be rotated, through a limited angle, around the axis of the telescope. The position angles given below are those of the slit, and are measured from the north point of the eye end of the telescope around through the east. With this arrangement the following observations were secured:

Position angle 1909 August 13	Position angle August 20
90°	110°
45°	88°
0°	45°
	0°
	335°

The central component is in all cases the brightest. Next in intensity is the more refrangible one, while

¹ Notes on the spectrum of α Ceti. W. W. Campbell, *Astrophysical Journal*, 9, 30, 1899.

² *The Observatory*, 22, 152, 1899.

³ *The Mills Spectrograph of the Lick Observatory*, *Astrophysical Journal*, 8, 123, 1898.

the component having the greatest wave-length is comparatively faint. It is difficult to decide whether the relative intensities of the three are exactly the same in all the exposures. There is, however, no such variation as would exist if any two of them were polarized in planes at right angles to each other. The measured position of the triplet, taken as a single line, is the same for all position angles.

A number of observations were made with a Fresnel rhomb in front of the Nicol. Only the photograph described below is sufficiently exposed to be of value. The position angles of the rhomb refer to the plane perpendicular to its reflecting surfaces.

1909 August 14.

(1) Exposure without rhomb. Position angle of Nicol and slit 90° .

(2) With rhomb, position angle 135° . Position angle of Nicol and slit 90° .

(3) With rhomb, position angle 45° . Position angle of Nicol and slit 90° .

The effective exposures are not all the same in (1), (2) and (3), but there is no evidence of variation in the relative intensities of the components. The position of the triplet, measured as a single line, remains unchanged in the three exposures, showing that the relative intensities of the outer components remain substantially the same.

I may suggest, for the benefit of any one caring to repeat these experiments or to carry them further, that an instrument of greater resolving power, or at least greater separating power linearly than the Mills spectrograph with the short camera, be used. Under good conditions this instrument readily shows the bright stellar line to be multiple; but when it comes to a matter of determining the relative densities of the components the grain of the plate and a number of other factors seriously interfere with the observation.

W. H. WRIGHT.

June, 1910.

NOTE ON SOME LINES IN THE SPECTRA OF THE NEBULAE.

In a former paper I published the wave-lengths of some of the brighter nebular lines.¹ It has been the intention for some time to extend the investigation so as to include, as far as possible, all the lines that can be observed with the means now at our command. With this end in view, a number of photographic observations have been made for the purpose of detecting any lines which might hitherto have escaped observation. A few have been found which have not been observed before, so far as the writer is aware, and provisional values of these are herewith given.

ORION NEBULA.

In addition to lines reported by other observers the following, unquestionably belonging to hydrogen, are present on the photograph of 1908 December 17:

¹ *Lick Observatory Bulletin*, 1, 154, 1902.

² This line was not observed by Ames in the vacuum tube.

The value given is the computed one, $\lambda = 3647.20 \cdot \frac{m^2}{m^2 - 4}$.

In nebula		In hydrogen tube (Ames)
$\lambda 3734$	faint	$\lambda 3734$
3722	v. faint	3722
3712	v. v. faint	3712
3704	v. faint	(3704) ²

An additional line is suspected near $\lambda 4137$.

N. G. C. 7027.

The following lines (in addition to $H\alpha$ and others reported by previous observers) have been observed on more than one plate. The wave-lengths are merely approximate, being based on an interpolation curve passed through the nebular lines $H\alpha$, D_3 and N_1 .

$\lambda 6301$	faint
6548	brighter than above line
6583	fairly bright

The second and third make with $H\alpha$ a conspicuous triplet. $H\alpha$ is very bright in all the nebulae that have been observed.

W. H. WRIGHT.

July, 1910.