ing to about 400 km/sec, sharply bounded and strong.

3. The violet component of a doubled hydrogen line is usually two to three times as strong as the red one. For $H\gamma$ the tendency is less conspicuous, and for $Ca II$ it may be reversed.

4. During both minima $H\beta$ shows emission on the red side.

5. The spectrum undergoes drastic changes from cycle to cycle.

6. The difference in absolute magnitudes, deduced from the radii and temperatures, is in sharp contradiction to that derived from the well determined light-curve.

Further spectroscopic and photometric studies are needed. The system may be tentatively considered as consisting of two components of about the same mass, but very different sizes. The bluer component has apparently a rotation (see No. 2) about 15 times as fast as would be required if it rotated with the revolution period. At the same time, the blue component seems to eject a continuous stream of material from its far side, which spirals around the whole system, forming a tenuous envelope.

*Harvard College Observatory,
Cambridge, Mass.*

Gaposchkin, Sergei. Computing elements in an eclipsing system.

A modification of the method of Sharbe¹ is developed in detail. It seems to be fast and accurate, and to involve a minimum of auxiliary tables for computing the elements for eclipsing variables.

1. *Bull. Obs. Central Poulkovo* **10**, Part I, 31, 1923.

*Harvard College Observatory,
Cambridge, Mass.*

Gaposchkin, Sergei. The variable star HD 26750, BD —10°863, HV 11087.

A star in the field of the variable star YY Eridani was found to show a drop in brightness during the year 1944. The Harvard patrol plates, extending from March 7, 1888, to December 23, 1945, show only one minimum, lasting at most 321 days, with constant brightness at the bottom. If we ascribe the variable to the eclipsing class, the period must be over 53, perhaps over 57, years; the range is 0.8 magnitudes, 8.5 to 9.3 photographic. The spectrum is that of an M6 giant; no secondary spectrum was visible in December, 1944.

*Harvard College Observatory,
Cambridge, Mass.*

Henroteau, F. A rapid calculator.

This is a device to attach to what is now called a fully automatic mechanical calculator such as the Monroe calculator or the Friden Calculator.

The device takes the numbers from the machine without having to read them and writes them on the paper. It also takes the numbers from the paper and places them on the keyboard of the machine without having to read them.

Each figure 1, 2, 3, 4, 5, 6, 7, 8, 9 and 0 is written on the machine or on the paper by means of different colors. A rotating drum formed of ten successive sectors, each of which is displaced by one unit rank with respect to the other, and each of which contains ten color filters, revolves in front of the numbers to be picked up.

When the proper filter passes in front of the proper number it activates the single photo-cell of the system for a very short time and trips the thyatron which during this short time is just connected to the cell by means of a rotating commutator. As the thyatron remains energized it pushes down its corresponding key of the machine's keyboard by means of an electromagnet, if the number has to be placed on the machine, or pushes down its corresponding colored pencil, if the number has to be written on the paper.

*Farnsworth Television and Radio Corporation,
Fort Wayne, Ind.*

Henroteau, F. A wide-field reflecting telescope.

This telescope which would replace the Schmidt telescope with advantage is composed of two concentric spherical mirrors which neutralize each other's aberrations. The first mirror which first receives the light of the stars is convex while the second which has a greater radius than the first and which also has a central aperture to allow the light of the stars to reach the first mirror, is concave.

According to computations it should give images as good as those of the Schmidt telescope. It is, however, much simpler to build.

*Farnsworth Television and Radio Corporation,
Fort Wayne, Ind.*

Hertz, Hans G. A theory of the Trojan group, with application to (659) Nestor.

A general theory for the asteroids of the Trojan group has been outlined by Brown and Shook in their *Planetary Theory*. The details have been carried out and have led to a development of an approximate numerical theory that may be con-