The Brackett series of hydrogen is shown in Sirius as well as Paschen β , γ and δ (α is blended with atmospheric Ω). Betelgeuse, Mira Ceti and R Leonis (minimum) are stars of increasing redness; apart from two new bands, near 1.1 and 1.2 μ , the spectra are unexpectedly smooth. The infrared spectrum of ϵ Aurigae is recorded but appears due entirely to the F component. Evidently the infrared component is even cooler than estimated on the assumption that the star falls on the mass-luminosity relation.

An extension of the program is outlined.

*McDonald Observatory,
Fort Davis, Tex.*

Ledoux, P. On stellar models with convection and discontinuity of the mean molecular weight.

The recent development of nuclear physics has led to the consideration of the stars (main sequence) as being composed of an external part in radiative equilibrium where the total flux of energy $L(r)$ can be considered as constant and a convective core where the generation of energy takes place. As this implies a progressive exhaustion of hydrogen, the mean molecular weight $\bar{\mu}$ will increase uniformly (mixing) in the convective core.

Thus in the course of the evolution, a state will be reached where μ changes abruptly from one value $\bar{\mu}_e$ in the envelope to a value $\bar{\mu}_i$ in the core. This is particularly true if the carbon cycle (Bethe's theory) is responsible for the generation of energy.

The change in $\bar{\mu}$ can be assimilated to a discontinuity as diffusion in a star is extremely slow. But, as pointed out by Hoyle and Lyttleton when the ratio $\bar{\mu}_i/\bar{\mu}_e$ exceeds a certain limit, say α_c , some difficulties arise in connection with the continuity of $L(r)$ across the surface r_c where the discontinuity of $\bar{\mu}$ takes place.

This paper presents a general discussion of the problem. It is shown that when $\bar{\mu}_i/\bar{\mu}_e$ exceeds α_c , turbulent mixing takes place at the interface, leading to a region of transition in stable radiative equilibrium where $\bar{\mu}$ varies continuously from its value $\alpha_c\bar{\mu}_e$ to $\bar{\mu}_i$ according to a law $\mu \div P^{1/5}m(r)$. The solutions for that region are described in the plane of Milne's variables (U , V) generally used for the fitting together of parts of the star built on different models. In particular, if the region of transition must be followed immediately by the convective core (as it should be in the process of evolution indicated above),

then only one solution is possible for that region.

The numerical values of some characteristics of stars built on that model are given. Their interpretation as a sequence of evolution, the central temperature remaining the same, gives an appreciable increase of the radius, luminosity and central condensation as $\bar{\mu}_i$ increases. The mass of the convective core starts by increasing, fresh material becoming available for the nuclear reaction. But as soon as the chemical composition has been altered appreciably, its mass decreases rapidly while the total mass m_t of the core plus the region of transition varies more slowly and goes on increasing for some time. However it has a flat minimum for $\bar{\mu}_i/\bar{\mu}_e \simeq 1.5$, and it is difficult to follow the evolution any further as, in reality, $(M - m_t)$ cannot increase.

To illustrate some aspects of the problem, the case where the discontinuity of $\bar{\mu}$ occurs in the external layers, where a complete analytical treatment is possible, is also considered. The same law as in the interior, for the variation of $\bar{\mu}$ in the transition layer, is recovered by considering in detail the process of mixing. It is also shown that the time taken for the establishment of that region of transition is very short compared to the time required for the building up of the discontinuity of $\bar{\mu}$. Contrary to the opinion expressed by Hoyle and Lyttleton who considered especially this case in connection with their theory of accretion of interstellar hydrogen, such a discontinuity will not lead to a dynamic state with large scale motions.

*Yerkes Observatory,
Williams Bay, Wis.*

McCuskey, S. W. Variations in the stellar luminosity function.

The important assumption of constancy in the relative proportions of stars of different luminosities at all points in the galactic plane makes possible an analysis of galactic structure based upon general star counts. It is admitted, however, that there are serious fluctuations in this luminosity function from point to point and even over comparatively small distance ranges. In an attempt to evaluate systematically these variations, observations based on spectral types, colors and star counts are being made in eleven regions of the galactic plane from $l = 12^\circ$ to $l = 200^\circ$. The spectral classifications are complete to $m_{po} = 12.25$.

Analyses have been completed for two regions LF1 at 19^h27^m , $+6^\circ9$ in Aquila and LF2 at