Eggen, Olin J. A photoelectric study of the eclipsing variable TX Ursae Majoris.

The eclipsing variable star TX Ursae Majoris has been observed on 40 nights from 1935 to 1942 with the photoelectric photometer on the 15-inch refractor of the Washburn Observatory. A measure of precision of the observations is indicated by the probable error of a single normal magnitude of ± 0.0046 mag.

The light elements of Rügemer were used in the computation of the phases with the exception that the amplitude of the sine term for the variation of the period was reduced to $0^d.015$.

After rectification for small reflection and ellipticity effects, the Algol-type light-curve revealed minima of depths 2.00 mag. and 0.032 mag. for the primary and secondary respectively. Anticipating the darkening coefficients as 0.4 and 0.6 for the B8 and gF2 components, a solution for the elements has been made. Because of the uncertainty of the value of the mass ratio at present available, no attempt has been made to proceed beyond "intermediate elements."

*Washburn Observatory,
Madison, Wis.*

Fath, E. A. The triple variability of 12 Lacertae.

On the basis of 63 maxima from 1917 to 1940 the mean period of variability is $0^d.19308902$ with a mean range of 0.082 mag. Besides this mean period there is a second period of $0^d.164850$ with a range of 0.031 mag. and a third period of about $0^d.1316+$ with a range of 0.017 mag.

*Goodsell Observatory,
Northfield, Minn.*

Gaposchkin, Cecilia Payne. The remarkable variable Y Ceti.

The star named Y Ceti has been recorded on only two plates, taken at the Cape Observatory on 1893 October 7 and 1894 September 26 respectively. Harvard plates taken within ten days of each date do not show the star, and it has never since been seen by any observer.

Through the kindness of the Kapteyn Astronomical Laboratory at Groningen the author has received contact prints of the two discovery plates. Each bore two exposures; one on Cetus, one on a region in Sagittarius; the earlier was a photovisual plate. On comparison with a Harvard photograph of the Sagittarius region it is clear that the two images identified as Y Ceti are in

fact those of two Sagittarius stars. The variable Y Ceti must therefore be regarded as non-existent.

*Harvard College Observatory,
Cambridge, Mass.*

Gaposchkin, Cecilia Payne, Sergei Gaposchkin, and Margaret Mayall. The variations of ρ Cassiopeiae.

The variations of ρ Cassiopeiae have been studied on all available Harvard photographs. Observations are recorded as far back as JD 2409850 (November, 1885) and are fairly continuous for the past 45 years.

The star has varied continuously in brightness, for the most part with small range. Maximum is about 5.5 mag. A well-marked minimum, 6.7 mag., occurred at JD 2416800, and many less prominent ones are recorded. The star is too bright for accurate study with this material, but most of the visual maxima and minima recorded by Hassenstein are shown by the Harvard curves, which strengthens the evidence for their objective existence. During the past year the star has fallen to about the eighth photographic magnitude. No periodicity is evident in the variations, which are executed in an interval of the order of a hundred days.

A number of spectra have been recorded with various instruments. A few were classified by Miss Cannon; these and all others have been independently classified by Mrs. Mayall and Mrs. Gaposchkin. The spectrum of earliest recorded type is supergiant F8p (1885); the latest, M5p (1946). During the intermediate time the spectrum has varied between F8 and K5. In general, the spectrum has been of latest type when the star was faintest; but at JD 2419600 it was K5 when the star was near maximum.

At all times the spectrum is most peculiar; for example in 1946, while displaying titanium oxide bands appropriate to class M5, it shows no corresponding strengthening of the Ca I line 4227. The impression is that the late-type characteristics overlie the G supergiant during times of minimum, producing the overall effect of a later spectrum. Guthnick, in his discussion of the Potsdam spectra, considered that they gave the impression of being complex.

*Harvard College Observatory,
Cambridge, Mass.*

Green, Louis C. Oscillator strengths for the continua of CaII.

A calculation of $df/d\epsilon$ at three or more points in each of five continua of CaII, 4s-p, 4p-s, 4p-d,