results are obtained from meteors with sub-points less than 150 km distant from the centroid of the three observing stations. A total of 41 meteors was found which satisfied the above condition, had radar echoes from all stations lasting over one second, and were visually observed from at least one point. The values of mean radar height for each meteor varied from 79 to 112 km with an average height of 94 km. The corresponding values for the same 41 meteors determined by combining radar ranges with visual plots gave heights from 68 to 142 km with an average height of 100 km. The addition of 21 meteors of long range or with very short duration echoes did not materially alter these figures. Heights computed purely from visual plots showed a much greater spread and were not accurate enough to be of value individually. The mean height of 26 meteors computed from visual observation alone was 98 km, agreeing well with the other results.

It would appear that, for the August period, the heights computed from radar alone average somewhat lower than those found from a combination of visual and radar data. This indicates that there is a tendency for the radar echo to occur near the end of the visual trail—for the Perseids at least. Confirmation of this fact, and a check on the reliability of the correlation of the radar echoes with visual meteors, was found by computing from the radar ranges alone the position of each echoing source in the sky and then comparing this with the visual meteor plot. The mean angular distance on the celestial sphere of all radar points from the corresponding visual meteor plots was only 7 degrees, a good agreement since this figure includes the summed effect of at least five separate sources of error. Over twice as many radar points were plotted near the end of the visual trails as near the beginning.

Additional radar heights were computed for meteors appearing on the same night but not visually observed. A compilation of all radar heights determined, corresponding to a total of 94 different meteors, gave a mean height of 95 km. The moving type of echo (*b* or *h* characteristic in our classification) was observed for 16 of these meteors. Mean height of appearance of this echo was 110 km and mean height of disappearance 95 km.

It seems clear from the above results that meteors produce ionization clouds which may be satisfactorily detected on the 30 megacycle band from widely separated points on the earth's surface. Triangulation of these clouds by range

measures alone is practical and yields results which compare favourably with those obtained by any other methods. This technique, when applied to individual meteor echoes of certain types, makes possible the accurate determination of daytime meteor paths and orbits. It may also be noted that these results confirm our earlier conclusions concerning echo heights and the relation between the echo form and distance from the radiant.

*Dominion Observatory,
Ottawa, Ont.,
and National Research Council of Canada.*

Morgan, H. R. A possible lag in the nutation.

In a recent paper¹ on the lunar nutation Dr. Jeffreys asks for an examination of observations for a possible lag in the nutation. The Washington observations of circumpolar stars, 1903-1925, recently discussed for the nutation constant,²

CORRECTIONS TO THE NUTATION CONSTANT, $N = 9''.210$

From the Right Ascensions

Star	ΔN	<i>W</i>	$\Delta\theta$	<i>W</i>
43 H Cep	−.076	7	+°.49	10
α UMi	+ .060	22	+ .26	16
Grmb 750	− .027	11	+ .25	11
Grmb 944	+ .004	8	+ .62	10
51 H Cep	− .086	17	+ .20	15
4 B UMi	− .079	8	+ .36	7
1 H Dra	+ .092	16	− .23	13
30 H Cam	+ .035	9	− .77	8
6 B UMi	− .032	11	+ .58	11
57 B UMi	− .137	10	+ .44	10
ϵ UMi	− .133	6	+ .13	7
δ UMi	− .005	9	+ .12	9
λ UMi	+ .041	7	+ .23	6
76 Dra	− .072	6	+ .25	6
39 H Cep	− .011	10	+ .32	12
Means	−.016 ± .013		+°.22 ± .06	

From the Declinations

Star	ΔN	<i>W</i>	$\Delta\theta$	<i>W</i>
43 H Cep	−.149	6	+°.50	5
α UMi	+ .054	12	− .07	14
Grmb 750	+ .007	6	− .07	6
Grmb 944	− .117	11	− .73	9
51 H Cep	+ .047	9	− .08	11
4 B UMi	+ .020	8	− .32	10
1 H Dra	+ .039	10	− .38	12
30 H Cam	+ .110	5	− .29	6
6 B UMi	+ .077	5	+ .61	5
57 B UMi	− .047	8	− .24	8
ϵ UMi	+ .033	13	+ .62	11
δ UMi	+ .019	15	+ .08	15
λ UMi	− .008	6	− .64	6
76 Dra	− .030	5	+ .86	7
39 H Cep	− .116	5	+ .93	5
Means	.000 ± .012		.00 ± .08	

have been re-solved using equations of the following form:

$$\Delta\alpha = (-.557 a \sin \Omega - 15 b \cos \Omega)\Delta N \\ + (15 b \sin \Omega - .557 a \cos \Omega)N\Delta\theta,$$

$$\Delta\delta = (-.037 a' \sin \Omega - b' \cos \Omega)\Delta N \\ + (b' \sin \Omega - .037 a' \cos \Omega)N\Delta\theta.$$

where $\Delta\theta$ represents the phase angle of any lag; a , a' , b and b' are the Besselian star constants; and N is 9^m.210. The results are given in the table. The residuals indicate considerable uncertainty in the results, but it seems important to present them. The large residuals are due, partly to poor distribution over the period of the moon's node, and partly to large accidental range in the observations. A later Washington series is to be examined.

The quantity $\Delta\theta = +0^{\circ}.11$, if real, means a lag of two days in the principal term of the nutation.

These observations are satisfied by the nutation:

$$\Delta\alpha = +9''.202 A + ''018 B, \\ \Delta\delta = +9''.202 A' + ''018 B',$$

where A , B , A' , and B' are the coefficients of ΔN and of $N\Delta\theta$ given above.

1. *M. N.* 108, 209, 1948.
2. *A. J.* 50, 125, 1943.

*Yale University Observatory,
New Haven, Conn.,
and U. S. Naval Observatory,
Washington, D. C.*

Nassau, J. J. and D. A. MacRae. A study of the galactic structure in a clear region in Cygnus.

The present investigation is a part of a general program, at the Observatory, dealing in a broad sense with the study of the fluctuations of the stellar luminosity function. The field under investigation has an area of 11 square degrees with a center at R.A. = 20^h0^m and Dec. = 38^o0. It is relatively free from space absorption with the exception of about one square degree near its southern boundary.

The photographic and photo-red magnitudes and spectra of 1230 stars were determined. Beginning with Fo and later, all stars with few exceptions were classified either giants or dwarfs up to the limiting photographic magnitude of 12.15. For stars brighter than 11.0 four luminosity classes were used.

The mean value of interstellar absorption in the clear region was derived from 683 stars and

found equal to 0.53 mag. per kiloparsec. The color excess of the stars in the obscured region seems to indicate that an interstellar cloud sets in at about 500 parsecs with a maximum value of about 0.6 mag. over and above the general absorption of the region. This result however is provisional as our present data are limited to 75 stars only.

The division of the stars with giants and dwarfs in each spectral group indicates a rapid decrease in the ratio of dwarfs to giants from Fo-F5 group to G8-K3 group. In the former group we have about 90 per cent dwarfs while in the latter about 5 per cent.

Space densities, corrected for absorption, for all the spectral groups were computed for the stars in the relatively clear region. The densities have been used to compute a luminosity function at distances of 100, 200, 400 and 600 parsecs and show no significant differences between the four regions. Likewise no significant difference exists where our results are compared with the standard van Rhijn luminosity function, with one exception at absolute magnitude between -1.5 and +1.0. Here the maximum excess in our case in the log $\varphi(M)$ is 0.3.

*Warner and Swasey Observatory,
Case Institute of Technology,
Cleveland, Ohio.*

O'Connell, D. J. K., S. J. RY Scuti, an eclipsing binary of the V444 Cygni type.

A photographic light curve of RY Scuti was obtained from estimates on 774 plates taken at Lembang and Riverview observatories. The writer was at the time unaware of the work done on this variable by S. Gaposchkin¹ and Popper.² The period agrees well with Gaposchkin's. The ranges at primary and secondary minimum are 0.52 and 0.47 mag. respectively. The ellipticity constant, z , and the reflection constant, b , were determined from the light curve as 0.31 and 0.01 respectively.

Primary minimum is much wider than secondary, the ratio of the widths being 1.76. Secondary minimum is 0^d.11 earlier than the midpoint between two primary minima. It would follow that the eccentricity of the orbit is 0.27 and the longitude of periastron 268^o. On the other hand, Popper's radial velocity measures are altogether inconsistent with such a large orbital eccentricity. These apparently conflicting photometric and spectroscopic observations can be reconciled, if one assumes that the broadening of primary minimum is caused by an "atmospheric eclipse"