servations are still too few to regard this period as definitive. So far there is a curious scarcity of observations of No. 10 near maximum. For No. 6, the indications are that the rise from minimum to maximum takes at least 40 days. The recent observations of No. 2 so far show such a small range that the variation, if real, would not have been noticed. No. 3 behaves like a slow variable, but it may prove to have a short period. Prof. Joy is investigating these brighter variables spectroscopically.

There appear to be seven variables which are probably RR Lyrae type and cluster members. The median magnitude of these variables is 15.62. Measured with the same sequence, the magnitude of the 25 brightest stars in the cluster is 14.73, with the 6th star at 14.3 and the 30th, 14.9. The value last determined by Shapley in 1918 for the 25 brightest stars was 14.89. The two values are in remarkably close agreement, in view of independent sequences and selection of stars. Four variables regarded as field stars are very faint, and at considerable distance from the cluster center. Because of its low southern declination, the cluster is accessible at the Dunlap Observatory for only a limited time each night, but it will be followed over a period of years until definitive periods can be derived.

The modulus of the cluster, uncorrected for absorption, may be taken as the mean magnitude

of the RR Lyrae stars, 15.62, and is half a magnitude brighter than the modulus of 16.14 published by Shapley and Sawyer in 1929, without regard to variables. Since the color excess as measured by Stebbins and Whitford is very high, $+0.29$, the cluster is probably relatively near to us, possibly as close as 3500 parsecs.

Table I gives a summary of the information on the variables in this cluster. The first nine variables were found by Bailey, the last seven by Sawyer. The coordinates of the new variables were measured to correspond as nearly as possible to the system used by Bailey.

1 *Circ. Harv. Coll. Obs.* 33.

2. *Ann. Harv. Coll. Obs.* 38.

*David Dunlap Observatory,
Richmond Hill, Ontario.*

Seeger, Charles L. The variation of the Cygnus "radio star" at 205 Mc.

Seventy-five two-hour observations were made of the 205 Mc radio frequency radiation originating in the direction of Cygnus. The observations were grouped half in November and December 1948, and half in April and May 1949. The early observations show only occasional variability, while those of the later group were seldom non-variable. For all times when substantial activity was observed (approximately 1700 minutes) the average peak to peak period was 0.8 minutes. During any given two-hour period the rate might be 0.1 to 10.0 times this average value. The peak to peak period seldom varied by more than two to one during a two-hour period. This variability has been observed at substantially all altitudes of the source. The amplitude of the variations on occasion has reached values one hundred times the minimum detectable value. At these times the energy of the variable component has been equivalent to the total steady radiation received from approximately two hundred square degrees of the sky, centered on 42 degrees and 20 hours.

*Cornell University,
Ithaca, N. Y.*

Shapley, Harlow and Virginia McKibben Nail. Further evidence of supergiant long-period variables.

In 1923 the three brightest variable stars in the southern globular cluster 47 Tucanae were shown to be long-period variables of extraordinarily high luminosity at maximum. Their crowding toward the center of the cluster argued against the possibility of their being superimposed galactic stars. Some long-period variables were later

TABLE I. VARIABLE STARS IN NGC 6626

No.	x	y	Max.	Min.	Mean	Notes
Known Variables						
1	+174.0	+188.5	15.1	16.4	15.75	1
2	-47.3	+63.1	14.3	14.8	14.55	2
3	-32.9	+111.0	14.6	15.4	15.00	2
4	-34.5	+33.6	13.6	14.8	14.20	3
5	-44.8	+16.4	14.9	15.8	15.35	1
6	+34.1	+50.4	14.3	15.2	14.75	4
7	+172.2	+102.7	15.9	17.0	16.45	5
8	+227.3	-222.3	14.9	16.3	15.60	1
9	-158.6	-252.4	14.9	15.9	15.40	1
New Variables						
10	+96	-79	13.5	14.6	14.05	6
11	-14	+35	15.0	16.3	15.65	1
12	+148	-49	14.9	16.5	15.70	1
13	-92	-24	15.2	16.7	15.95	1, 7
14	-131	-100	15.7	16.2	15.95	5, 8
15	-472	-186	15.8	17.0	16.40	5
16	+432	-372	15.9	17.0	16.45	5

NOTES

1. Probably RR Lyrae type and cluster member.
2. Possibly slow variable.
3. Period probably 14 days, W Vir. type.
4. Period probably many weeks.
5. Probably field star.
6. Slow variable, type not yet known.
7. Double, south preceding varies.
8. Double, probably south component varies.