Journal to the American Astronomical Society, with the continued support of the Benjamin Apthorp Gould Fund. This transfer provided at last a more secure foundation for existence and continuity.

A tremendous broadening of the range of astronomical research has taken place since Gould's day, rather than a passing of active interest from one field to another. The content of recent volumes proves the vitality of those branches of astronomy served by the *Astronomical Journal*, and encourages the hope that it will continue to realize the aim of its founder during its second century.

Editor

APSIDAL MOTION IN THE ECLIPSING BINARY AG PERSEI

By JOSEPH ASHBROOK

1. The principal points of interest in the eclipsing variable AG Persei = HD 25833 are the rapid apsidal motion, and the fact that both stars have been observed spectrographically. The variable is described in N. L. Pierce's Finding List¹ as follows:

$P = 2^d.03$; Spectrum B3 + B3; Max. = 6.5 mag.; $A_1 = 0.3$ mag.;

$A_2 = 0.2$ mag.; $D = 6^h$; eclipses partial.

In this paper, a fresh determination of the rate of apsidal motion is made, in which new photoelectric observations will be used.

2. *Previous Investigations.* Attention was first directed to this star when it was discovered to be a double-lined spectroscopic binary. The orbital elements were determined by Plaskett² from 24 Victoria spectrograms. The light variations were discovered independently by Huffer³ and at Leiden.⁴ Huffer's photoelectric light curve showed a marked displacement of the secondary minimum. A uniform solution was not possible, but a satisfactory fit of the observations was had on the assumption of total limb darkening; the larger star was found to be the brighter.

In 1937-38 AG Persei was reobserved photographically by Martin⁵ at Leiden. The displacement of the secondary minimum was found to have changed in sign since Huffer's observations, indicating apsidal motion. Each displacement of the secondary provided an observed value of $e \cos \omega$; by taking $e = 0.0512$ following Luyten's⁶ rediscussion of Plaskett's radial velocities, Martin concluded that the revolution of the apse required 53 years.

This procedure does not determine the quadrant of the apse unambiguously, and indeed Oosterhoff⁷ pointed out in 1943 that the alternative apsidal period of 20 years seemed supported by the very uncertain values of $e \sin \omega$ furnished by the relative widths of the minima. But as we shall see below, more recent observations rule out the possibility of any period near 20 years.

3. *The Visual Companion.* Any discussion of the light variations of AG Persei is embarrassed by the fact that the variable is the brighter component of the visual double star ADS 2990 = OΣ71. The latest micrometer measurement is by Baize:⁸

1934.93 214°.5 0".96 4 nights.

There has been practically no change in relative position since the first micrometer measures in 1846, so that it is evident that all photometric observers have measured the combined light. Accordingly the apparent magnitude of the visual companion is an important datum, but no photometric determination exists. We have only visual estimates by the double star observers, from which we obtain the following values of Δm :

Observer	Δm	Nights	Aperture
O. Struve ⁹	2.0	2	15-inch
Dembowski ¹⁰	2.2	6	7
Engelmann ¹¹	2.0	6	7½
Burnham ¹²	3.0	1	18

The unweighted mean $\Delta m = 2.3$ is probably systematically too large, in view of the tendency among double star observers to underestimate the fainter star of an unequal pair. In view of our uncertainty of the magnitude and ignorance of the color of the visual companion, no reliable correction can as yet be made to the observed light curves. Huffer's photometric solution, in which no allowance was made for the light of the visual companion, must therefore await revision.

4. *New Observations.* During the interval JD 2432923-946 I secured 96 photoelectric observations of AG Persei at the Amherst College Observatory with the 18-inch Clark refractor. The photometer used a 1P21 photomultiplier tube behind a Corning 5543 blue glass filter 5.10 mm in thickness. This combination has an effective wave length of approximately 4240 Å for a B3 star. The output of the photomultiplier was fed through a DC amplifier designed and constructed by John Jewett of Amherst, and was registered by a Brown recording potentiometer. Use of the

TABLE I. AMHERST OBSERVATIONS OF AG PERSEI

HD 25539-AG								
JD	Mag.	ϕ	JD	Mag.	ϕ	JD	Mag.	ϕ
2432900 +			2432900 +			2432900 +		
23.5018	-.066	.2525	31.5287	+.177	.2091	36.6339	-.061	.7256
.5088	-.092	.2560	.5337	+.171	.2116	.6377	-.049	.7275
.5131	-.111	.2581	.5711	+.094	.2300	.6428	-.047	.7300
.5178	-.101	.2604	.5770	+.074	.2329	.6473	-.031	.7322
.5226	-.101	.2628	.5839	+.024	.2363	37.4850	+.177	.1451
.5276	-.109	.2652	.5898	+.006	.2392	.4892	+.188	.1472
.5375	-.098	.2701	.5953	-.028	.2420	.4946	+.194	.1498
.5423	-.080	.2724	.6003	-.054	.2444	.5012	+.187	.1531
.5471	-.074	.2748	.6125	-.093	.2504	.5068	+.200	.1559
.5518	-.063	.2772	.6177	-.114	.2530	.5746	+.185	.1893
.5576	-.038	.2800	.6233	-.091	.2558	.5796	+.181	.1917
25.6598	+.160	.3158	.6292	-.100	.2587	.5858	+.195	.1948
.6657	+.177	.3191	.6339	-.082	.2610	.5904	+.180	.1971
.6722	+.172	.3223	.6405	-.117	.2642	39.6627	+.164	.2185
.6782	+.164	.3253	.6455	-.108	.2667	.6673	+.145	.2208
.6849	+.182	.3286	.6514	-.102	.2696	.6756	+.117	.2249
.6896	+.169	.3309	.6567	-.083	.2722	.6772	+.112	.2257
.6949	+.172	.3335	.6625	-.069	.2750	.6817	+.106	.2279
.7010	+.172	.3365	.6669	-.065	.2772	.6902	+.079	.2321
.7072	+.174	.3396	.6729	-.044	.2802	.6944	+.052	.2342
.7120	+.180	.3420	.6779	-.017	.2827	.6984	+.038	.2361
31.4723	+.196	.1813	.6877	.000	.2875	.7020	+.016	.2379
.4765	+.192	.1834	.6930	+.019	.2901	.7066	+.014	.2402
.4809	+.188	.1856	.6990	+.052	.2930	.7102	+.008	.2420
.4855	+.188	.1878	.7076	+.079	.2973	.7141	-.015	.2439
.4898	+.187	.1900	36.5882	-.017	.7031	46.5403	+.179	.6086
.4952	+.176	.1926	.5935	-.030	.7057	.5456	+.190	.6113
.5001	+.183	.1950	.5995	-.029	.7086	.5504	+.191	.6136
.5048	+.188	.1973	.6038	-.047	.7107	.5549	+.176	.6158
.5096	+.186	.1997	.6060	-.055	.7118	.5570	+.179	.6189
.5156	+.181	.2026	.6102	-.056	.7139	.5632	+.191	.6199
.5219	+.173	.2058	.6140	-.064	.7158	.5699	+.186	.6232
			.6175	-.074	.7175	.5743	+.178	.6254
			.6281	-.075	.7227			

Brown recorder made one-man operation of the photometer feasible.

The comparison star used was HD 25539, whose constancy has been vouched for by Huffer.³ The Draper Catalogue gives 6.53 for the photographic magnitude, and spectrum B3. Differential extinction was computed under the assumption that the zenithal extinction was 0.50 mag. The correction for differential extinction never exceeded 0.015 mag.

The observing procedure consisted in taking deflections of 1 to 2 minute duration on the variable and comparison star in turn. While this practice increased the time lost in moving the telescope, it had the important advantage of minimizing the effect of the unstable transparency of the New England sky. Each observation rests therefore on a single deflection on the variable and on the two adjacent deflections on the comparison star. The mean error of a single observation is ± 0.0066 mag. from the measures at maximum light, which for this purpose was assumed constant.

Since the period of AG Persei differs so slightly from two days and since the interval during which the star could be observed was short,

relatively few measures could be made at maximum, and the quarter points of the light curve are not covered.

The Amherst observations of AG Persei are given in Table I. Here, and throughout this paper, the phase ϕ is defined by the formula

$$\phi = (\text{JD}_{\odot} - 2,420,000.0000) \text{d}^{-1} 492920.$$

5. *Displacement of Secondary Minimum.* The observed phases of primary and secondary minimum have been determined from the mean light curve by the analytical process described by Oosterhoff,¹³ and used by Martin in his discussion of the Madison and Leiden observations of AG Persei. The method is to reflect the light curve about an arbitrarily chosen trial phase of minimum, and to form the sum of the squares of the differences in ordinates of the two branches, taken at equidistant phases. This procedure is repeated for several trial phases, the phase of minimum being that for which the sum of the squares is least. This method presupposes that the light curve is symmetrical, but it may easily be shown¹⁴ that the asymmetry in the light curve of AG Persei arising from orbital eccentricity is negligible.

The phases of primary and secondary minimum thus found are

$$\begin{aligned} p &= 0^{\text{P}}2630 \\ s &= 0^{\text{P}}7202. \end{aligned}$$

The displacement of the secondary minimum is therefore

$$\delta = s - p - 0^{\text{P}}5 = -0^{\text{P}}0428.$$

6. *Apsidal Motion.* The component $e \cos \omega$ of the orbital eccentricity is given, when powers of e beyond the first are neglected, by

$$e \cos \omega = \pi \delta / (1 + \operatorname{cosec}^2 i).$$

If, following Huffer, we put $i = 78^{\circ}3$,

$$e \cos \omega = 1.5375\delta.$$

Martin's values of δ , which he obtained from the Madison and Leiden light curves by the process described above, are included in Table II.

TABLE II. PHOTOMETRIC VALUES OF $e \cos \omega$

J.D.	δ	$e \cos \omega$	Observer
2424946	+0 ^P 0220	+0.0388	Huffer
28946	-0.0181	-0.0278	Martin
32930	-0.0428	-0.0658	Ashbrook

From these three values of $e \cos \omega$, we derive $e = 0.0670$, and ω at these three epochs as $59^{\circ}7$, $114^{\circ}5$ and $169^{\circ}2$. The longitude of periastron is therefore

$$\omega = 59^{\circ}7 + 5^{\circ}00 (t - 1927.18)$$

where t is the date in years. This corresponds to an apsidal period of 72 years. Luyten's rediscussion of Plaskett's radial velocity observations gives $\omega = 42^{\circ} \pm 16^{\circ}$ for 1924.95, which is only $6^{\circ}6$ smaller than the formula predicts.

The new orbital eccentricity 0.0670 agrees satisfactorily with the spectroscopic value 0.0512 ± 0.015 (m.e.).

The adopted orbital inclination, $78^{\circ}3$, may be a few degrees too small because of Huffer's disregard of the unresolved visual companion. However the error so introduced into the apsidal period is quite minor; even putting $i = 90^{\circ}$ will increase the period by only two per cent. Oosterhoff's suggestion that the apsidal period is about 20 years cannot be reconciled with the new observations, as it would require that ω be about 70° for their epoch; this would make δ positive instead of negative as observed. Moreover, the 20-year period gives a less close fit for the spectroscopic value of ω .

If the parameters K_1 and K_2 characterizing the internal density distributions of the two com-

ponents are supposed equal, their common value K is given by¹⁵

$$K = \frac{P}{P'} / \left[r_1^5 \left(1 + 16 \frac{m_2}{m_1} \right) + r_2^5 \left(1 + 16 \frac{m_1}{m_2} \right) \right].$$

Here P and P' are the orbital and apsidal periods, r_1 and r_2 the radii of the stars in terms of the radius of the orbit, and m_1 and m_2 the masses. Taking the four latter quantities from Huffer's paper, we find $K = 0.0033$. For polytropic stars, this corresponds to a polytropic index $n = 3.6$. The most important source of uncertainty in these values of K and n comes from the radii of the stars. While the ratio of the radii is $k = 0.70$ from Huffer's solution, R. M. Petrie's¹⁶ spectrophotometric determination of the magnitude difference between the components gives $k = 0.89$.

7. *Elements.* For the prediction of times of minimum of AG Persei the following formulae have been derived, for primary and secondary minimum, respectively.

$$\begin{aligned} \text{JD } \odot \text{ Min}_1 &= 2424946.5156 + 2^{\text{d}}02872933\text{E} \\ &\quad - 0^{\text{d}}0442 \cos \omega \\ \text{JD } \odot \text{ Min}_2 &= 2424947.5300 + 2^{\text{d}}02872933\text{E} \\ &\quad + 0^{\text{d}}0442 \cos \omega \end{aligned}$$

where $\omega = 59^{\circ}7 + 0^{\circ}0278\text{E}$.

These formulae were based on the Washburn, Leiden and Amherst observations only. Some visually determined minima by Skoberla¹⁷ were not used, because of their large and erratic residuals.

8. *Summary.* New photoelectric observations of AG Persei made at Amherst are reported, and epochs of primary and secondary minimum obtained. Discussion of all available photometric observations of sufficient precision gives 72 years for the period of the apsidal motion, and 0.067 for the orbital eccentricity. The photometric solution must remain uncertain until the magnitude and color of the close visual companion are determined. The present data give $n = 3.6$ for the polytropic index. New ephemeris formulae are given in the last section.

9. *Acknowledgments.* The new observations of AG Persei were made possible by the Office of Naval Research, under contract N8onr-574. It is a pleasant duty to thank Prof. Warren K. Green for the use of the facilities of the Amherst College Observatory, and Mr. John Jewett for his very valuable assistance in electronic matters.

REFERENCES

1. *Contr. Princeton Univ. Obs.* No. 22, 1947.
2. *Pub. Dominion Astroph. Obs.* 3, 188, 1925.