3. *Washburn Obs. Pub.* **15**, 192, 1931.
4. *B. A. N.* **3**, 242, 1927.
5. *B. A. N.* **8**, 286, 1938.
6. *Pub. Astr. Obs. Univ. Minnesota* **2**, No. 2, 42, 1935.
7. *B. A. N.* **9**, 416, 1943.
8. *J. Observateurs* **18**, 92, 1935.
9. S. W. Burnham, *General Catalogue of Double Stars* **1**, 40, 1906.
10. *Misure Micrometriche* **1**, 141, Rome, 1883.
11. *A. N.* **115**, 81, 1886.

12. *Mem. R. A. S.* **47**, 234, 1883.
13. *B. A. N.* **5**, 37, 1929.
14. H. N. Russell, *Ap. J.* **36**, 59, 1912.
15. *Ibid.* **90**, 649, equation (1), 1939.
16. *Harv. Obs. Monograph* No. 7, 242, 1948.
17. *Zs. Astroph.* **11**, 15, 1935.

Yale University Observatory,
New Haven, Conn.,
1949 May.

THE CEPHEID X VULPECULAE

By M. S. DAVIS

The purpose of this investigation is to revise the light elements of the Cepheid X Vulpeculae. The new observational material consists of 90 plates taken with the 3-inch Ross camera of the Yale University Observatory from 1947 to 1948. The comparison star sequence used was that determined by Nassau and Ashbrook.¹

Each plate was estimated at least twice by Pickering's method, suitable precautions having been taken against preconception. Estimates on about three-fourths of these plates had previously been made by Dr. Ashbrook. Since no systematic differences in magnitude between the two observers were apparent, his estimates were combined directly with my own. The probable error of a single plate is ± 0.07 magnitude.

Table I gives the photographic mean light curve. The period $6^d.319490$ was used in the formula

$$\Phi = 0.15824062 (\text{JD} - 2427000)$$

to compute the phases. The period is that used by Nassau and Ashbrook,¹ but the reciprocal is there erroneously given as 0.15842062 instead of 0.15824062. An amplitude of 0.92 mag. was

found between maximum light 9.71 and minimum 10.63.

TABLE I. PHOTOGRAPHIC MEAN LIGHT CURVE

Φ	Mag.	No. Plates	Φ	Mag.	No. Plates
0 ^h 0208	9.80	10	0 ^h 5411	10.57	7
.0903	9.91	4	.5960	10.61	5
.1892	10.09	9	.6325	10.55	3
.2731	10.33	7	.7171	10.64	8
.3461	10.29	6	.7683	10.49	6
.4155	10.44	6	.8592	10.16	7
.4865	10.50	6	.9437	9.68	6

Beginning with the Nassau-Ashbrook formula

$$\text{JD}_{\odot} \text{ Max.} = 2420636.231 + 6^d.319381E,$$

the $O - C_{N,A}$ were computed for ten sets of observations from 1905 to 1948 as shown in Table II. As too long a period was indicated, a least-squares solution was made, giving $+0^d.016$ for the correction to the initial epoch and $-0^d.000080$ for the correction to the period. The residuals $O - C_D$ in Table II refer to the revised elements.

It is well known that the maximum of a Cepheid occurs slightly later visually than photographically. A second least-squares solution was made with this lag as an additional unknown, but its

TABLE II. COMPARISON OF OBSERVED MAXIMA OF X VULPECULAE WITH OLD AND NEW ELEMENTS

Observed Max $\odot$	E	$O - C_{N,A}$	$O - C_D$	Observations	Observer
2417299.853	- 528	+ ^d 26	- ^d 11	273 vis	Seares-Haynes
17438.755	- 506	+ .13	+ .02	49 vis	Luizet
19808.519	- 131	+ .13	- .01	74 vis	Luizet
20636.248	0	+ .02	+ .08	~200 ptg	Robinson
25843.478	+ 824	+ .08	- .04	144 ptg	Nassau-Towson
26481.59	+ 925	- .07	+ .11	82 vis	Kukarkin
27499.037	+ 1086	- .04	+ .06	217 ptg	Nassau-Ashbrook
28004.616	+ 1166	- .01	+ .02	166 ptg	Nassau-Ashbrook
28857.785	+ 1301	+ .04	- .04	187 ptg	Nassau-Ashbrook
32421.920	+ 1865	+ .04	- .08	90 ptg	Davis

value, vis. max. - ptg. max. = $-0^d.005 \pm 0^d.048$ (p.e.), did not differ significantly from zero. Therefore the first solution was used to give the new light elements

$$\text{JD}_{\text{max}} = 2420636.337 + 6^d.319301E \\ \pm .020 \pm .000020 \text{ (p.e.)}$$

Thanks are due to Dr. Ashbrook for friendly advice.

REFERENCE

1. *A. J.* **50**, 99, 1943.

Yale University Observatory,
New Haven, Conn.,
1949 June.