tial, no clouds or wind. An examination of the objective with the eye at the focal plane will indicate the approach of atmospheric stagnation and an anemometer will be a useful meteorological instrument.

It appears that 1954 and 1956 are the only suitable oppositions for canal photography in the next 22 years. We expect to assign the 60-inch telescope to explore the problem of canal photography in 1952, and use it and the 100-inch in 1954 and 56. It seems that an electrically controlled motion-picture camera at the Cassegrain focus will produce best results. It is expected that arrangements can be made to keep the planet under continuous visual observation, and when the canals begin to come out run the motion-picture cameras until the interval of opportunity passes. If the whole pattern becomes visible, it is believed that the optical quality of the atmosphere is so perfect that exposures as short as a hundredth of a second can be given without the momentary distortions that appear in ordinary good seeing.

*Mount Wilson Observatory,
Pasadena, Calif.*

Pettit, Edison. Nebular photometry.

Some 18 years ago Hubble and Humason showed that a nearly linear relation exists between the distances and velocities of galactic systems. The distances of eight nearby systems were determined from recognizable stars and the rest were determined from the apparent magnitudes. They showed that the range of the absolute photographic magnitude was from -11.3 to -16.3 and the velocity diagrams could be expressed by

$$\log V_{\text{km}} = .2m_{\text{ph}} + .507.$$

From the material then available it was also thought the color index was about 1.1, and showed little relation to distance.

It was then determined to secure more abundant material and in the intervening years Humason has secured 555 velocities of nebulae for which magnitudes (m_{ph}) and color indices are being measured with the photomultiplier cell.

In Humason's list 351 nebulae have now been completed, of which 43 overlap Stebbins and Whitford. Of these the average difference Δm_{ph} , Pettit minus Stebbins and Whitford is zero; average without regard to sign is .1 mag. The average difference Pettit minus Shapley and Ames is $-.1$ mag. and the average without re-

gard to sign is .4 mag. The average difference in color index Pettit minus Stebbins and Whitford for the 24 identities is $+.05$ mag. A study of the reductions of the North Polar Sequence stars used for calibration and of the Selected Area central stars used for extinction seem to confirm the idea that a blanket .07 mag. should be subtracted from the color index. Early- and late-type stars are being run to test this principle.

While the list of nebulae is not well suited to statistical analysis, the following points of interest may be mentioned.

(1) Of the 108 nebulae for which the inner zone and the approximately complete object were measured no sensible difference of color index was found between the inner zone and the whole system in 30 objects. In 45 cases the inner zone was redder and in 33 bluer. However, 12 nebulae have been measured for the nucleae. These are all redder than the entire object except two, which are slightly bluer.

(2) The range of color index is considerable, varying from .2 mag. for six such objects as the Lundmark-Mallotte object, several spirals and irregulars to 1.71 for 16-17 mag., E-types. Emission patches in certain nebulae, like NGC 604, have color indices as low as 0. The bluest E-types are 0.48 (1770 Eo), 0.65 (7343 E3), 0.68 mag. (4478 E2). Spirals cover the range .2 to 1.3 mag. with one object IC 342 at 1.54 mag. The E-types run from .5 to 1.71 mag. and irregulars .28 to .90 mag.

(3) The average color index for the magnitude classes shows a practically constant value 0.8 for the brighter nebulae down to the 14th mag. At this point the color index rises to 1.02, reaching 1.12 at 16 mag. and 1.71 at 17 mag. There are only two nebulae in the 17 mag. class and the result for this magnitude must be regarded as not representative. The same investigation was made using only E-type nebulae, reducing the spectral class to Go from Humason's spectral types. No substantial change was seen in the results. The effect of absolute magnitude on color index is part of the investigation.

*Mount Wilson Observatory,
Pasadena, Calif.*

Popper, Daniel M. Notes on the spectra of eclipsing binaries.

Fifteen years ago Wyse compiled observational data on the spectra of eclipsing binaries. A considerable number of those systems showed a