to be derived from (2). The use of the existing elimination equations is in fact made possible by the differential correction of c_1 and c_3 , since they yield new values of the three geocentric distances when new values of c_1 and c_3 are known.

In the differential correction of c_1 and c_3 one must realize that he is dealing with three sets of these quantities:

- (P) Preliminary, from the first approximation;
- (C) Computed, from Gibbs's expressions or equations (1), or other;
- (O) Objective, the aim of the differential correction, equivalent to "observed" (O) of $\Delta\alpha, \Delta\delta$ (O-C).

Thus we have also the three sets of differences:

$$(P-C) \quad \Delta'c_i = c_i(P) - c_i(C), \quad (4)$$

residuals from the "representation" of c_i ,

$$(O-C) \quad \Delta c_i = -B_i''\Delta\rho_2, \quad (5)$$

to be sought, where

$$B_i'' = 3(c_i - A_i)(\rho_2 - R_2 \cos\psi_2)/r_2^2, \quad (6)$$

the A_i being the ratios of the sectors or time-intervals;

$$(O-P) \quad \Delta''c_i = \Delta c_i - \Delta'c_i. \quad (7)$$

These differences we use in conjunction with the well-known equation

$$E\rho_2 = F_1c_1 - F_2 + F_3c_3, \quad (8)$$

where the E and the F 's are numerical constants resulting from the elimination of ρ_1 and ρ_3 from the "fundamental equations"—or of $\rho_1, \rho_3, \dot{x}, \dot{y}, \dot{z}$ from equations (3). Equation (8) holds for (P) and (O) values of c_1 and c_3 , and thus

$$(O-P) \quad E\Delta''\rho_2 = F_1\Delta''c_1 + F_3\Delta''c_3, \quad (9)$$

or, since $\Delta''\rho_2 = \Delta\rho_2$, the (P) and (C) values of ρ_2 being identical,

$$E'\Delta\rho_2 = -(F_1\Delta'c_1 + F_3\Delta'c_3), \quad (10)$$

where

$$E' = E + F_1B_1'' + F_3B_3''. \quad (11)$$

The differential correction of c_1 and c_3 involves equations (4), (6), (11), (10), and (5), in that order. It is thus complete for the method of Gauss-Gibbs. For the Lagrangian method additional expressions similar to (5) and (6) are available for the correction of $\dot{x}, \dot{y}, \dot{z}$.

*Department of Astronomy,
University of California,
Los Angeles, Calif.*

Huang, Su-Shu. Line intensities for an atmosphere with varying absorption.

The departure of stellar atmospheres from the Milne-Eddington model has been investigated theoretically by many authors.¹ All these authors tried to solve the equation of radiative transfer for the spectral lines by one kind of approximation or another. Here we shall give a different approach to the problem. It is semi-empirical in nature; hence, physical ideas rather than mathematical rigor has been emphasized in the following considerations.

In the equation of transfer for a spectral line

$$\mu \frac{dI}{dt} = I - J, \quad (1)$$

the source function J ,

$$J(t) = \frac{1}{2} \frac{\sigma_\nu}{\kappa + l_\nu + \sigma_\nu} \int_{-1}^1 Id\mu + \frac{\kappa + l_\nu}{\kappa + l_\nu + \sigma_\nu} B(\tau[t]), \quad (2)$$

consists of two terms. The first term represents scattering, and the second term, thermal emission. Here t represents the optical depth in the line, and τ , the optical depth in the continuum. They are related in the following way:

$$t = \int_0^\tau \frac{\kappa + l_\nu + \sigma_\nu}{\kappa} d\tau. \quad (3)$$

The other notations are apparent.

In the source function the dependence of the factors $B(\tau)$, σ_ν/κ and l_ν/κ on t is through the relation $\tau(t)$ as implied in eq. (3), because by their physical nature these quantities are directly functions of τ . If the scattering is not dominant, i.e., weak lines and wings of strong lines, we may therefore write

$$J(t) \equiv J(\tau[t], t), \quad (4)$$

and the main variation of $J(t)$ with t will be a result of the variation of τ . The direct dependence of $J(t)$ on t should be regarded as a small correction for the scattering. Or in other words if we approximate the source function $J(t)$ by a double series in t and τ , i.e.,

$$J(\tau[t], t) = \sum_{n=0}^m \sum_{i+j=n} c_{ij} t^i \tau^j, \quad (5)$$

the coefficients of the τ terms will be large compared with the other terms. In the case of local thermodynamical equilibrium,

$$J(t) = B(\tau[t]), \quad (6)$$