Although the separation of the components never exceeds $1''.5$ or $.08$ mm, this separation is large enough to throw some doubt on the interpretation and evaluation of the value β . While for well-blended images the relation $\beta = (1 + 10^{.4\Delta m})^{-1}$ holds very closely, this is no longer the case for incomplete blending.^{9, 10} The result for B may, therefore, have been systematically influenced, and should be accepted with reservation.

The residuals indicate a systematic behaviour over the decade 1939–1949; the material however is not sufficient to permit any conclusions about a possible perturbation. It should be pointed out that an unseen companion of the fainter component might be an explanation for the seemingly high mass of the latter.

I am indebted to Dr. van de Kamp for putting the plate material at my disposal.

REFERENCES

1. *Ann. Obs. Strasbourg*, 5, Fasc. 2, 7, 1949.
2. *Sproul Obs. Pub. No.* 4, 32.
3. *General Catalogue of Stellar Parallaxes*, Introduction, 1935.
4. *A. J.* 33, 92, 1920.
5. *Pub. McCormick Obs.* 4, 6, 1927.
6. *A. J.* 45, 175, 1936.
7. *Astronomie* 57, 44, 1943.
8. *A. J.* 46, 38, Table III, 1937.
9. *A. J.* 54, 188, 1949.
10. *A. J.* 55, 215, 1951.

*Sproul Observatory,
Swarthmore College,
Swarthmore, Pa.,
1950, November.*

NOTES AND OBSERVATIONS

THE DUPLICITY OF φ SAGITTARII

The star φ Sagittarii is a spectroscopic binary which has been observed as a slow occultation on three occasions and as a visual double by R. H. Wilson, Jr.¹ The data available are as follows:

1926.58 Single	In B	26½-inch
1929.47 Appeared to bury itself in moon before disappearing.	W	
1931.46 Single	In Rst	27-inch
1933.29 Glided (occultation)	Wl	
1933.66 Slow (occultation)	Wl	
1937.42 27°.4, 0".13	4n R. Wil. J	18-inch
1948.71 Single (<0".101)	2n φ	26½-inch
1949.54 17°.0, 0".21	3n R. Wil. J	18-inch
1950.79 Single (<0".100)	In φ	26½-inch

The last four are interferometer measurements.

The minimum zenith distance at which this star could have been observed by Wilson is 67° and the apparent duplicity observed by him must undoubtedly be ascribed to the effect of atmospheric dispersion.

REFERENCE

1. *A. J.* 55, 155, 1950.

W. S. FINSEN,
*Union Observatory,
Johannesburg,
1950 October.*

NOTE ON THE DUPLICITY OF ROSS 79
AND OF CAPELLA H

The duplicity of Ross 79 ($6^h5^m4, +10^\circ21'$, 1900) recently reported by Van de Kamp¹ has been confirmed on five plates taken with the

20-inch refractor of the Van Vleck Observatory in 1949 and 1950, but not on any plates of the regular parallax series taken in the years 1937 to 1940. However, the residuals resulting from the parallax solution gave some indication of a variation in the proper motion, and the star was continued on the observing list, but the plates taken since 1940 have not yet been measured. The presence of the companion, together with possible variations in the effective exposure, may introduce considerable uncertainty into attempts to explain the variable proper motion.

A faint companion to Capella H ($5^h10^m, +45^\circ44'$, 1900) was detected on Van Vleck Observatory plates in 1936² and its presence was confirmed visually by Kuiper.³ The two components are still not separated on the plates but the following approximate measurements suggest a change in the position angle and perhaps in the separation.

Date	d	p
1935.6	1".8	117°
1940.7	1.6	132
1948.7	2.2	133
1950.8	2.0	137

REFERENCES

1. *A. J.* 55, 160, 1950.
2. *A. J.* 45, 120, 1936.
3. *Ap. J.* 84, 359, 1936.

CARL L. STEARNS,
*Van Vleck Observatory,
Wesleyan University,
Middletown, Conn.,
1950 November.*