critical characteristic for checking any theory of stellar evolution. The observational data on rotation can now be significantly enlarged since with present high dispersion spectrographs rather slow rotations can be measured. Whether the magnetic moment of a star is entirely a relic of the star's past or whether at least in part the magnetic field is continuously regenerated in the present stellar status does not yet seem clear. In the first alternative the magnetic fields presently observed would give direct information regarding the magnetic fields at the birth of the stars.

Third class: chemical abundances in stellar atmospheres. Many features of the chemical composition of stars are presumably determined by the origin or evolution of the universe as a whole, which is outside the present discussion. However, differences in chemical composition between different types of stars seem to be of great importance for theoretical work on stellar evolution. Among the many suspected differences the following three may be the most active examples at present.

a) Hydrogen-poor stars. The number of hydrogen-poor stars thus far discovered is still small, but their importance is evident, as hydrogen-poorness is presumably caused by the burning up of hydrogen in the stellar interior and hence indicates directly an advanced stage in stellar development.

b) Carbon-rich stars. An unusual high abundance of carbon has long been recognized in the R and N types stars and possibly also in some of the Wolf-Rayet stars. Recently it has been suggested that this carbon-richness may possibly be explained by the contraction of a star, after the exhaustion of its interior hydrogen, until the

internal temperature is sufficiently high to permit the formation of carbon from helium. If this suspicion is substantiated carbon-richness will be a direct indication of old age of a star.

c) Abundance peculiarities of high-velocity stars. It seems at present possible to interpret the various spectroscopic peculiarities of high-velocity stars in terms of a relatively low abundance of all the heavier elements in these stars. Such an interpretation tends to support the suspicion that the low-velocity stars may have originated in places unusually rich in interstellar dust, which in itself is rich in the heavier elements.

Fourth class: kinetic characteristics. The kinetic characteristics of groups of stars, such as their speed of galactic rotation, their concentration toward the galactic plane, and their average random velocities, are in themselves presumably determined by the history of the galaxy as a whole rather than by the history of the individual stars. However, since these kinetic characteristics can change during the lifetime of a star only little if at all, these characteristics act as permanent labels for all groups of stars. Thus they indicate which types of stars may represent consecutive stages of one sequence of evolution and which types of stars must belong to separate sequences of evolution.

The above listing of classes of inherited and acquired stellar characteristics is certainly incomplete and of a roughly-summarizing nature. The listing, however, may suffice to indicate what a large variety of observational and theoretical work can directly contribute toward our knowledge of stellar evolution.

*Princeton University Observatory,
Princeton, N. J.*

ABSTRACTS*

Bappu, M. K. Vainu. The variation of emission-line intensities in CQ Cephei.

The binary nature of the Wolf-Rayet star HD 214419 was first noted by McLaughlin and Hiltner,¹ at Michigan in 1941. Shortly thereafter Gaposchkin² showed the star to be an eclipsing system with two well-defined minima. Hiltner³ recently studied the system photoelectrically and obtained light-curves in the ultraviolet and yellow regions of the spectrum as well as a light

curve of the emission line 4686A of HeII. The variation of 4686 differed from that of the integrated case in that it had maxima at conjunctions and minima at elongations, or in other words the geometry of the system is such that 4686 is most intense when either star is eclipsed.

To determine whether other lines of HeII and

* Of papers presented at the Eighty-sixth Meeting of the American Astronomical Society, East Cleveland, Ohio, December 26-29, 1951.