metric substitutions and eliminates apparent indeterminacies usually encountered when small numbers of significant figures are carried.

The procedure entails the usage of differences of given quantities, rather than the quantities themselves. A seven-place table of natural sines is used at the outset, to find sines and cosines of all three right ascensions and three declinations. This table is not used again. Each quantity, such as $\sin \alpha$, is differenced. This same procedure is followed for the observed times and the solar coordinates. Orbital elements, found from one tabulated value and two differences in this way, are accurate enough to indicate the physical orbit of the body in space. They are not, however, accurate enough for use in an ephemeris computation. Intermediary quantities, rather than the elements, are used for this purpose. The ephemeris so computed compares favorably in accuracy with that computed by the usual procedure.

The method was tried out for the case of three observations of Comet 1951a. As a standard of comparison, an eight-place machine computation was made, by the same method. Other computations, using the same method and same observational data, are carried out entirely independently of this or of each other.

The results include, in addition to the machine computation, three separate logarithmic computations—a 3-place, a 4-place, and a 5-place computation—and a slide rule computation. For the logarithmic computer, it is suggested that the 4-place computation is amply accurate, while for a bright comet even the 3-place computation might suffice. The slide rule computation gives a good search ephemeris for more than 30 days ahead of the two-day interval on which the orbit was based, and gives elements about one decimal lower in accuracy than that required in an astronomical telegram. Residuals for the middle place are, in all but the machine computation, found by finite difference formulae, and in all cases indicate with sufficient accuracy how well the parabolic assumption was justified.

*Howard University,
Washington, D. C.*

Menzel, Donald H. Origin of the aurora borealis.

Geophysicists have generally supposed that the aurora borealis results from the penetration of the earth's upper atmosphere by a neutral

stream of ions and electrons from the sun. The ion stream may arise from spicule activity, the energetic surgelike prominences that occur near the sun's poles. Future work is needed to establish the origin of the corpuscular stream.

Measures of the abundances of cosmic material lead us to suspect that this stream from the sun consists largely of ionized hydrogen: protons and electrons in nearly equal numbers. Focused by solar and terrestrial magnetic fields, these ion clouds reach the earth. In exceptional cases, as Gartlein and Meinel have shown, the aurora borealis displays broadened and shifted lines of the Balmer series.

The physics of the emission of hydrogen gas, as developed for studies of the planetary nebulae, applies also to the ion clouds in the vicinity of the earth. The emission probably occurs well outside of the earth's atmosphere, but relatively close to the earth. From the observed brightness of the hydrogen emission we can get rough estimates of the density of the ion cloud. Some uncertainty in this density arises because we have no direct way of measuring the temperature. However, for the most likely values of the temperature, which lie between 10^4 and 10^5 degrees K, the corresponding densities are in the range of 10^7 and 10^8 ions per cm^3 . This figure is somewhat higher than the value of $10^3/\text{cm}^3$ calculated by Whipple and Gossner for electron gas uniformly distributed within the orbit of the earth.

The total energy carried by such an ion stream is about 10^5 times greater than the visible energy flux of the brighter aurora and only 20 times smaller than the flux of solar radiation. However, the limited size of the clouds saves the energy situation. Even so, the total energy of the bombardment probably far exceeds the value usually quoted for the total energy of a magnetic storm derived from the variability of the magnetic field.

One important problem relates to the relative stability of certain auroral forms: streamers, curtains, and arcs, which may maintain an almost unchanging pattern for times up to one hour. These luminous regions are fixed relative to axes attached to and rotating with the earth. The earth's motion over the interval traverses regions of space thousands of times more extended than the original auroral streamer. To explain this persistence of auroral forms, one may assume that currents in the auroral levels significantly modify the earth's magnetic field,