1. J. P. Wild and L. L. McCready, *Australian J. Sci. Res.*, Ser. A, **3**, 387, 1950.
2. *Ark. Fys.* **1**, 569, 1950.
3. *Phys. Rev.* **82**, 232, 1951.
4. *U.S.S.R. Acad. Sci. J. Phys.* **10**, 25, 1946.

*Central Radio Propagation Laboratory,
National Bureau of Standards,
Washington, D. C.*

Sharpless, Stewart. A study of the Orion aggregate of B stars.

A preliminary Hertzsprung-Russell diagram for the Orion aggregate of B stars has been constructed on the basis of a distance modulus of 8.0 mag. Spectral types and luminosity classes were obtained from spectrograms taken on W. W. Morgan's general B star survey and on a special survey of the fainter stars in Orion earlier than Ao.

The reddening in Orion has been investigated by means of these spectroscopic data and the observations of color excess by Stebbins, Huffer, and Whitford¹ and by the writer. The results indicate an abnormally high ratio of total to selective absorption in the Orion region of about 16. This is in agreement with previous results obtained for θ^1 Orionis.^{2,3} Application of the higher ratio in correcting the points on the H-R diagram for interstellar absorption tends to decrease the scatter in the main sequence and move the earliest type stars to higher luminosities.

1. *Ap. J.* **91**, 20, 1940.
2. W. Baade and R. Minkowski, *Ap. J.* **86**, 123, 1937.
3. Joel Stebbins and A. E. Whitford, *Ap. J.* **102**, 318, 1945.

*Yerkes Observatory,
Williams Bay, Wis.*

Thomas, Richard N. Chromospheric kinetic temperature from radio measures.

A major anomaly in the interpretation of the solar chromosphere at the present time is the apparent disagreement between temperatures inferred from optical observations and temperatures inferred from radio measures. In each case the relevant temperature is only a kinetic temperature. In the interpretation of several of the optical observations the departure from thermodynamic equilibrium of the solar chromosphere has appeared to influence appreciably the observations. The radio measures are, currently, interpreted as coming from free-free transitions in the field of the hydrogen ion. Heretofore, the influence of a departure from Maxwellian distribution of the electrons about the kinetic temperature of the chromosphere has been

ignored. It appears however that an average relative departure between the high energy and the low energy ends of the velocity spectrum amounting to one part in one hundred thousand will suffice to cause an error of a factor 2 in the inferred value of the temperature.

*University of Utah,
Salt Lake City, Utah.*

Urey, Harold C. The surface of the moon.

The planetesimal 100 km in radius which collided with the moon, forming Mare Imbrium, contained iron-nickel alloy objects some kilometers in dimensions which produced the grooves in the moon's surface and grey silicates which formed radiating ridges as far from the collision area as the grooves. The Mare Nectaris planetesimal contained larger objects. It is suggested that these objects were raised to a high temperature on their surfaces in the presence of cosmic gases which reduced iron oxides and caused the formation of an iron shell some distance below the surface. Black ridges of the Imbrium system west of Copernicus may be unreduced primordial planetary material. The order of fall of planetesimals forming the maria was approximately Foecunditatis, Nectaris, Crisium, Humorum, Serenitatis and Imbrium. All but the last contain no large craters and Imbrium none in the area of collision. The depths of lava produced were 10 to 15 km. The time of formation was about 10^5 years, judging from the time required for lavas to cool through a layer of pumice. The Mare Serenitatis planetesimal fell from the vertical, produced the black lavas of Tranquillitatis and was covered by the grey lavas of Imbrium. The entire process occurred at the terminus of the formation of the moon and its surface has been modified only slightly since.

*University of Chicago,
Chicago, Illinois.*

van Wijk, Uco. A photoelectric magnitude sequence in the Large Magellanic Cloud.

In the period from January to March, 1951, the magnitudes and colors of 16 stars between apparent magnitudes 10 and 14 in the Large Magellanic Cloud were obtained with the Linnell-King photometer attached to the 60-inch Rockefeller reflector of the Harvard Boyden Station in South Africa. A total of 63 comparisons were made between these stars, and at least three comparisons were obtained for each star. The zero point of the sequence was determined by