the difference in axial rotation between the main-sequence stars and the giants and sub-giants. The distribution of observed $v \sin i$'s for these two groups appears to be quite different, the more luminous stars having generally smaller rotational velocities. If this difference is real, the relation between axial rotation and spectral type among the early-type stars should be considered by luminosity class.

*Perkins Observatory,
Delaware, Ohio.*

Warwick, James W. Remarks on the structure of sunspots.

Three lines of evidence concur in indicating that sunspots possess a fine structure small compared with optical depth unity.

Brief summaries of magnetic field strength and visual observations are given. The main body of the paper deals with a curve-of-growth analysis of the spot spectrum. This analysis shows that the separation of the photospheric and spot curves of growth is to be understood as a turbulence effect, rather than as a magnetic intensification effect.

An intensification parameter is introduced. The limitations of the parameter are discussed in some detail. The paper concludes with a resumé of the ten Bruggencate-von Klüber interpretation of their data, which have formed the basis for the present paper.

*Sacramento Peak Station,
Harvard College Observatory,
Cloudcroft, N. Mex.*

Weston, Edwin B. The variable stars in the region of NGC 7023.

The stars in the region of the reflection nebula NGC 7023 are being examined for their distribution in space, presence or absence of variability in light, proper motion, and the presence or absence of emission lines in their spectra. The plates used in this study were taken with the Curtis Schmidt telescope of the University of Michigan during the years 1950 to 1952 and with the Crossley reflector of the Lick Observatory between 1902 and 1950. Some of the Curtis Schmidt plates were taken with an objective prism.

Within the region of obscuration caused by the nebulosity there are two apparent clusterings of stars. One of these is in the center of the nebula, and the other is located about $5'$ west of the center. The numbers of stars brighter than the

nineteenth photographic magnitude (approximately), the incidence of variability among these stars, and the incidence of $H\alpha$ emission among these that are bright enough to be examined on objective prism spectra have been compared for the stars in both apparent clusterings, in regions of very heavy obscuration, and in relatively clearer areas outside of the nebula. Stars were not classified as variable unless the observed light variation was one-half magnitude or more, and caution was exercised in the recognition of emission lines. Hence, all variables of small amplitude and emission objects with weak lines were excluded. Light curves were determined for many of the stars.

It was found that nearly all of the stars in the central clustering are irregular variables that resemble the RW Aurigae and T Tauri types. Some even show appreciable light variation during intervals of only a few hours. Nearly all of these show $H\alpha$ emission, and in addition other emission lines of hydrogen, Ca II, etc. were recognized in the spectra of a few of the brightest. However, few variables and emission-line objects were discovered in the apparent clustering to the west; one of the two variables discovered in this region appears to be an eclipsing variable. Evidence for only a small number of variables and emission-line objects was found for the regions of heaviest obscuration. Approximately normal incidences of variables and emission-line stars were indicated for the regions of relatively lower obscuration outside the nebula. The peculiar proper motions of all stars in the region of the nebula were found to be small, and there is no evidence of common proper motion groups among the stars in the apparent clusterings.

The central clustering of stars thus appears to be a physical cluster, and its members to constitute a T-association of the form defined by Ambartsumian.¹ On the other hand, the apparent clustering to the west appears to be neither a physical cluster nor a T-association. It is an apparent optical cluster produced by a window or lane of relatively high transparency between the individual clouds and filaments that form this nebula and through which we see the distant stars beyond the obscuration. As to the central cluster, the absence of common proper motion indicates that it is not a moving cluster that encountered the clouds and filaments composing the nebula, but rather that it has a genetic connection with the nebula in the sense that it may represent the birthplace of stars along the lines