THE PRESENT STATUS OF DOUBLE STAR ASTRONOMY

By K. AA. STRAND

Abstract. A general survey has been made of the present status of double star astronomy in regard to discoveries of new pairs, the measurements of relative and absolute motions of known pairs, as well as their magnitudes and radial velocities.

It is proposed that more emphasis should be placed on obtaining photographic observations of the wider pairs. The visual observers should be allowed the facilities of large refractors and reflectors for the difficult pairs most in need of observations. Further progress in the field also depends upon accurate data on parallaxes, mass ratios, magnitudes, spectral types and radial velocities.

The results of past surveys up to 1927 for double stars north of -30° are listed by Aitken and show a total of 17,180 pairs.¹ The inclusion of a pair in this catalogue was based upon the magnitude-separation formula, $\log d'' = 2.8 - 0.2m$, where d'' is the separation and m is the combined magnitude of the components of the double star. In order to bring the Aitken catalogue up to date, the large number of pairs discovered since in the various surveys and falling within the limits of the catalogue should be added. The Bloemfontein and Johannesburg surveys would contribute the largest number of brighter pairs with additional pairs from Kuiper's survey of stars brighter than 6.51 mag. H.S.² The additional large number of faint pairs will be those discovered visually by Jonckheere, Espin, Milburn and others, and from Barton's search of the astrographic catalogues.

Rossiter has recently published the final list of new double stars discovered at Bloemfontein.³ Based upon the magnitude-separation formula, $\log d'' = 2.5 - 0.2m$, this survey has contributed 7,350 new double stars over a period of 24 years; approximately 30,000 measures of these pairs are being prepared for publication.

The completed Johannesburg survey, including pairs to 9.0 mag. CPD and closer than $5''$, has added another 4,000 double stars, and according to a letter from Dr. van den Bos, who has completed the manuscript for the first six hours of right ascension, The *General Catalogue of Double Stars South of -19°* now contains approximately 16,000 pairs as compared with the 6,902 entries in the Innes loose leaf edition of 1926-27.

With the systematic double star surveys of Lick, Bloemfontein, and Johannesburg completed, it is safe to assume that only a relatively small number of new double stars brighter than the ninth magnitude will be discovered with separations within the limits set by these surveys, although the survey by Kuiper indicates that perhaps as many as 7 per cent additional pairs,

which were either overlooked or were too close at the epoch of the Lick survey, may be discovered.²

On the basis of van den Bos's estimate for the catalogue of southern double stars, we find that at the present time approximately 40,000 double stars are known, falling within Aitken's inclusion formula, even though the survey is incomplete for magnitudes fainter than 9.0. A continuation of visual work by Jonckheere, Espin, Milburn and others on the fainter stars as well as the search by Barton and others of the Astrographic Catalogues will result in the addition of large numbers of faint pairs.

An examination of the Aitken Catalogue and the lists of observations published since show as a whole the lack of adequate, systematic measures. The extensions of previous surveys to fainter magnitudes will only tend to make this problem more acute. Moreover, most of these pairs will consist of optical doubles and pairs with extremely slow orbital motions, while a very small number will have components which are either M dwarfs or white dwarf stars. These pairs can be discovered much more effectively from a search among stars with large parallaxes or proper motions, as has been done by Kuiper and Luyten, and in the various astrometric programs with long-focus refractors.

Since any statistical problem dealing with double stars requires the knowledge of accurate magnitudes of the components, spectra and colors, such a program must necessarily be confined to a lower limiting magnitude than ninth, supplemented by pairs selected on the basis of large parallax to keep the program within manageable proportions.

The number of astronomers engaged in routine visual observations of known pairs has steadily declined over a period of many years. From time to time a new name appears with a series of a few hundred measures but the bulk of the observations are made by astronomers who have been engaged in this field for more than twenty-five

years. The steady decline in the number of observers is alarming, but natural on account of the competition with the many new fields in astronomy where the chance of reward for hard work in the form of significant discoveries is much greater. Another problem facing the worker in the double star field is that the general catalogues are complete only to 1927, making it difficult to rely upon them in regard to pairs in need of measures at the present time. Only the observer who maintains a catalogue of measures published since 1927 is able to cope with this problem, until new general catalogues with representative up-to-date measures are published. These catalogues are now in the planning stage.

Visual observations of double stars with the micrometer continue to be the main source of information in regard to orbital motions. The conventional wire micrometer still seems to give the best results for the wider pairs, especially in position angle, while the modern form of double-image micrometer gives very accurate determinations of both position angles and distances for the close pairs.

For very close pairs with nearly equal components, the interferometer can give much higher accuracy than the micrometer measures. This method of observing relative positions of double stars was first applied by J. A. Anderson to observations of Capella with the 100-inch reflector,⁴ and interferometer measures have since been made primarily by Maggini, van den Bos, Finsen, R. H. Wilson and Jeffers. It has been claimed in the past that the error of measurement in this type of observations is of the order of 1 per cent. According to Finsen,⁵ this evaluation of the method is rather optimistic, and in the majority of cases the observational errors are more nearly of the order of 10 per cent, approaching but not reaching 1 per cent in the most favorable cases. Some interferometer observations have been found to be spurious, usually due to faulty collimation, effects of atmospheric dispersion especially at large zenith distances, or exceeding the practical limiting difference of magnitude, which is about 1.0 magnitude and for the closest pairs 0.5 or even less.⁶ In a recent note van den Bos lists a fairly large number of observations by Maggini and R. H. Wilson which he considers spurious.⁷

For double stars with definitely separated images on the photographic plate, Hertzsprung was the first to show that one good exposure with

a telescope of focal length of the order of 10 meters gives about the same accuracy as an ordinary visual measure with the micrometer.⁸ While the mean of repeated visual measures by the same observer during the same epoch is not materially improved by more than three nights of observations on account of the systematic personal errors, these errors are much smaller for the photographic measures. As a result it is possible to reach a much higher accuracy by using the photographic method when thirty to fifty measures are made on the same plate.

Investigations by van den Bos⁹ and Strand¹⁰ have shown that if the components on a photographic plate have a separation larger than .15 mm and are properly exposed under good seeing conditions, there are no systematic errors due to such photographic effects as the Kostinsky, Eberhardt, and turbidity effects. This means that if a telescope of 10 meters focal length is employed, pairs with separations larger than 3".0 are not affected by systematic errors due to the photographic effects, provided the magnitude difference between the components is less than one magnitude. For larger magnitude differences the limiting separation has to be increased to approximately 4".5 for magnitude differences equal to four magnitudes.

An estimate of the relative accuracy of the visual and photographic observations for the wider pairs can be obtained from a comparison of the systematic personal errors in the two methods of observation. For the visual observations this error varies considerably from observer to observer, from one pair to another, and also with the change with time in the separation of a measured pair. Estimates from binaries with accurately observed motions show that the error is of the order of ".05 and primarily affects the measured distance. The systematic personal measuring errors in the photographic observations are of the order of ".006. This means that we are able to reach nearly a decimal place further in accuracy by the photographic method than with visual observations. It could be expected that the personal error of measurement for the visual observations would cancel out as a random error if many observers contributed to the formation of a mean place. Only in the case of a few classical binaries do we find sufficient observations to check on this matter and it is found that in most cases the distances are systematically in error over a long period of time, or the errors change with the separation of the

components. In the case of η Cassiopeiae the distances attained from the means are systematically too small by $^{\circ}.04 \pm ^{\circ}.01$ (m.e.) over the entire period of observations,¹¹ and for the double star 70 Ophiuchi the observed distances are measured systematically too small when the components have separations larger than $2''.5$ and are measured systematically too large when the separation is less than $2''.5$.¹² In this case we find that one observer who has been very active over the past 25 years has measured all his distances systematically too large by $0''.17 \pm ^{\circ}.04$ (m.e.) over the entire period.

Fortunately, the visually observed position angles, when averaged over many observers, are essentially free from systematic errors, and the mean result of 100 nights' observations by various observers has a mean error of $^{\circ}.018$ (or $^{\circ}.21$ for a separation of $5''$).¹³

In comparison, we find that the mean position of a photographic normal place from four multiple-exposure plates is usually obtained with a mean error of $^{\circ}.010$, clearly indicating the much higher precision of the photographic method.

Up until 1938 Hertzprung was nearly the sole contributor to this field with his observations made with the Potsdam, Johannesburg, and Lick refractors. In 1938 his work at Lick was continued by Jeffers and a systematic program was started by Strand with the Sproul refractor and later on with the Yerkes and Dearborn refractors. Within the past three years a similar program for the southern double stars was initiated with the Lembang refractor by van Albada. The method initiated by Hertzprung has been retained unchanged as to its essential features; it includes the use of multiple exposure plates with 40 to 60 exposures, nearly monochromatic light to eliminate atmospheric dispersion, and gratings for compensation of magnitude differences. Improvements have been introduced due to the increased speed in photographic emulsions and the use of cameras with automatic timing and plate transport.

A continuation, and preferably an extension of the present photographic double-star programs, should relieve the burden on the visual observer who should be encouraged to concentrate on those pairs which cannot be measured accurately by photography. It should be emphasized that the photographic process offers a great deal of economy in that there are very few cases in which continuous series would lead to important orbit improvements, whereas the

majority of the pairs need accurate relative positions at intervals of ten to thirty years. The taking of the individual plates takes about half an hour, but present methods of measurement and reduction require approximately six hours. Thus far it has been possible to have the measurements keep pace with the output of plates. All plates of the past programs have been measured and on the current Sproul, Yerkes, Dearborn program 90 per cent of the plates have been measured with 600,000 settings. A letter from Professor Hertzprung informs me that he and his associates have completed 450,000 settings on the Lembang program. Even a large backlog in such a program is not a serious matter since the photographic plate, if properly stored, constitutes a permanent record which perhaps in the future can be measured and reduced more quickly by electronic means.

With the completion of their original parallax programs, several observatories have undertaken extensive series of observations of selected double stars for the purpose of determining orbital motions as well as mass ratios. Notable contributors to this field are the McCormick, Sproul and Yerkes Observatories. The initial plates in these series usually consist of a set of plates used in a regular parallax determination, but the plates are now being taken regardless of parallax factors in increased numbers at each opposition, and the individual mass-ratio determination is usually based upon a hundred or more plates taken over an interval of approximately 35 years. From the individual plate of unit weight the position of the star is determined with a probable error of $\pm ^{\circ}.030$ in each coordinate and usually no systematic errors exist for the plates within a series. This accuracy is considerably greater than the accuracy obtained in meridian observations where in general a position of unit weight has a probable error $\pm ^{\circ}.30$,¹⁴ and where systematic errors are common between observations obtained at different observatories. For this reason the photographic mass-ratio determinations are gradually replacing the mass ratios obtained from meridian observations for those binaries which have shown sufficient curvature in their orbital motion over the past 35 years.¹⁵

At the McCormick Observatory 28 binaries are at the present time under observation with approximately 10 preliminary determinations published or ready for publication. On this same program are 10 unresolved and 15 suspected astrometric binaries.

The Sproul program has, since 1937, been devoted to a systematic search for orbital motions of all "single" stars with parallaxes larger than $0''.1$, which are within reach of the 24-inch telescope. So far, orbital motions have been determined or found for the single stars Ross 614,¹⁶ Lalande 21185,¹⁷ and BD 20°2465. A letter by Dr. van de Kamp permits me to include the third star, the observational data of which have not yet been published. Strong indications of variation have been found for Ci 18, 2354, while Barnard's star, from intensive observations of about 200 plates each year over the past 15 years, shows definite deviations which so far have not been satisfactorily explained.

While 70 per cent of the current output of plates at the Sproul Observatory is on the program outlined above, the remaining 30 per cent is spent on a mass-ratio program essentially based upon the list of double stars given by Kuiper in his discussion of the mass-luminosity relation.¹⁸ A number of mass ratios have been finished and published, but observations of all the pairs are continued, so that the mass ratios as well as the parallaxes may be improved.

The Yerkes mass-ratio program lists at the present time 30 double stars. Most of the objects are classical binaries with large orbital motion. For some series the earliest plates were taken in 1903 (Krüger 60) and 1904 (η Cas), and the majority of the stars have been observed fairly regularly since 1914, although extensive series at each opposition were not started until 1946. For 10 binaries mass ratios and parallaxes have been published, and accurate orbits have been established from these series for Procyon¹⁹ and Krüger 60.²⁰

For several binaries in the mass-ratio programs mentioned above the components are not resolved photographically, and accordingly the light center of the two components relative to their center of mass is observed. If we assume that the center of light is situated on the line connecting the components, at a distance $sL_2/(L_1 + L_2)$ from the brighter star, where L_1 and L_2 are the luminosities of the components, measured in the wave-length considered, and where s is the separation, then the observed value of the mass ratio is not $B = m_2/(m_1 + m_2)$, but $B - \beta$, where $\beta = L_2/(L_1 + L_2)$.²¹ An investigation by Hall has shown that the empirical curve for β deviates from the theoretical curve for Δm larger than 2 mag.²² The uncertainty in the value of β gives a corresponding uncertainty in the derived

mass ratio; however, comparisons of mass ratios for the same stars obtained with the Sproul and Yerkes refractors, which have focal length in the ratio of almost 1:2 and therefore corresponding ratios in actual separations on the photographic plates, give nearly identical results if the empirically determined values for β are used. We may therefore assume that the mass-ratio determinations of unresolved binaries can be accurately determined if the magnitude difference between the components has been determined accurately by other means. For the unresolved binaries which remain unresolved visually as well as spectroscopically, the masses of the components cannot be rigorously determined. However, with a reasonable assumption about the value for $M_A + M_B$, a value for B , and hence β , can be found, and the corresponding value of M_A or M_B can be derived.

In regard to the magnitudes of the binaries, we still have to rely mainly upon the Henry Draper Catalogue, which includes nearly all stars as bright as 8.0, omits many of magnitude 8.0 to 8.5, and is very incomplete for fainter stars, especially in the northern hemisphere. Moreover, a large percentage of the magnitudes listed are not based upon direct photometric measures but upon *Durchmusterung* magnitudes, reduced to the Harvard scale by application of corrections derived statistically.

As far as magnitude differences between the components are concerned, we have for the brighter pairs, until recently, only the photometric measures of the Harvard Observatory.²³ Attempts have been made by Öpik²⁴ and Baize²⁵ to reduce the estimates of Δm to a photometric system. For the more difficult pairs, those with separations less than $2''.5$, as well as the pairs with large Δm , these values are considerably in error.

Accurate Δm 's have been obtained by Kuiper for all double stars north of -18° , brighter than 6.5 mag. H.S. and separations between $0''.5$ and $5''.0$.²⁶ However, the data have as yet not been published, although Dr. Kuiper has from time to time very generously communicated his results for individual stars when need arose for an accurate value of Δm . Kuiper's method of determining the magnitude differences between the components by means of objective gratings may perhaps be considered superior to any other method for pairs with separations less than $4''.0$, with the possible exception of the method developed by Muller with his double image micrometer.²⁷

Within recent years several series of magnitude differences have been published. Detre has published a list of magnitude differences for about 200 systems using a wedge photometer attached to a $10\frac{1}{2}$ -inch refractor.²⁸ Observations with a similar photometer attached to a $14\frac{1}{2}$ -inch refractor have been published by Wallenquist,²⁹ who also made a comparison between his measures and those published by Harvard and Detre, as well as measures published by Kooreman³⁰ and unpublished measures by Strand. The data by Kooreman and Strand were obtained from multiple exposure plates, but while Kooreman measured the plates in a photometer reaching an internal accuracy of .03 mag. (m.e.), those by Strand are estimates to nearest .1 mag. The magnitude scales were obtained from the grating images on the plates.

Several series of magnitude differences have been published by Muller, and his final results are published in a catalogue of Δm 's for 315 stars.³¹ These measures have been made with his double-image micrometer attached to the 16.2 cm and 48.6 cm Strasbourg refractors, and to the 23 cm and 60 cm telescopes at Pic-du-Midi. Series of measures have also been published by Hoffmann for 115 pairs³² and Miczaika for 827 pairs.³³

It appears from this summary that a great deal of effort has been spent in recent years on improving our knowledge of Δm 's for binaries. It is therefore of considerable importance that Dr. Wallenquist is at the present time preparing a catalogue in which he combines all the available data and appraises the accuracy of the various methods which have been used. The individual values in different series usually disagree by amounts in excess of what could be expected from the internal accuracy of the measures within a series.

With our present knowledge it would appear that the photoelectric method should be used in connection with instruments of long focal length to obtain the magnitudes and colors of pairs wider than $4''$, and the combined magnitudes for the closer pairs. The Kuiper and Muller methods should be used for obtaining the magnitudes of the individual components. The accuracy of the photoelectric method for this type of work is shown in a recent paper by Johnson in a three-color photometry of 40 wide doubles.³⁴ The initial program, if confined to the double stars for which such data are of special interest, could easily be completed by a single observer in each

of the two hemispheres in a period of two years. The new survey by Morgan and Harris of obtaining photoelectric magnitudes and colors as well as spectra in the revised Yerkes Atlas system (MK) for all stars brighter than 7.4 P.D. in the zone $+20^\circ$ to $+30^\circ$ is an important step in the right direction.

The main sources of information in regard to spectral classifications of double stars are still the Henry Draper Catalogue and the lists in the *Mt. Wilson Contributions*. These catalogues furnish in general the spectra for both components, if the pair is wider than $5''$ and each component is brighter than 8.0 mag. For the closer pairs with a combined magnitude brighter than 8.0 we usually have the spectrum of the brighter component, or the combined spectrum if the magnitude difference is less than 1.0 mag. We have in general spectra in the Morgan-Keenan-Kelmann system for the classical binaries within reach of the Yerkes telescopes. For certain binaries, which fall within groups of special interest, spectral classifications have usually been published if the objects are within reach of the observatories in the northern hemisphere, but in general the spectral classification of visual binaries is very incomplete, especially for the southern hemisphere.

It is, of course, impossible to obtain accurate spectral classification for all known binaries within a foreseeable future. Even in a program limited to binaries brighter than 6.51 H.S., as in Kuiper's survey, with a fairly limited number of objects to observe, the problem of obtaining the spectra of the faint components in close binaries poses a difficult problem.

The lack of spectroscopic observations is felt even more in connection with the need of radial velocities to determine the sign of inclination for the individual orbits as well as the relative orientation of orbits in multiple systems. In very few cases are radial velocities known with sufficient accuracy to give an independent check on the parallax of a double star with a well determined orbit.

A *General Catalogue of Stellar Radial Velocities* has recently been completed by R. E. Wilson.³⁵ Since it will contain the positions, magnitudes, spectral types, and definitive radial velocities of all stars whose velocities have been determined at any observatory, it will be of importance to the workers in the double star field. A total of 15,105 stars are included in the catalogue.

The only phase of double star astronomy

which leaves nothing to be desired is the orbit computation. Usually orbits appear for a double star long before the motion has been adequately measured, and revisions of earlier orbits have been published in great numbers for many of the classical binaries. For 70 Ophiuchi alone 42 orbits have appeared to date.

Catalogues of orbits are also published at regular intervals, the most recent ones being those by Finsen³⁶ and Baize.³⁷ It would be desirable to have future catalogues edited by a committee selected by Commission 26 of the I.A.U. in order to make them as correct and unbiased as possible in the selection of the orbits for the individual binaries.

In summarizing the present status of visual double star astronomy, we find that the current small number of observers will be able to provide the necessary observations for only a relatively small number of the approximately 40,000 pairs now known. The visual observer should accordingly be encouraged to observe only the close double stars which are too difficult for the photographic method. Since in general the pairs in this category most in need of observation are those which require the use of large telescopes, special efforts should be made to allow visual observers now using small and medium-sized telescopes the facilities of large refractors and reflectors. The photographic observer should select, among the vast number of faint double stars of moderate and large separations, those pairs for which the first traces of orbital motion may be measurable within a reasonable time. Since the orbital motion to be expected is proportional to the proper motion,³⁸ the pairs with known large proper motions should be given preference. For the very faint pairs in this category the photographic method should be extended to the use of large reflectors with observations made at the Cassegrain focus.

Further progress in the field also depends upon additional accurate data on parallaxes, mass ratios, magnitudes, spectral types and radial velocities.

REFERENCES

1. *New General Catalogue of Double Stars*, Carnegie Inst. of Washington, 1932.
2. *Pub. A. S. P.* **46**, 188, 1934.
3. *A. J.* **58**, 29, 1953.
4. *Ap. J.* **51**, 263, 1920.
5. *Union Obs. Circ.* **112**, 96, 1951.
6. *Ibid.* **93**, 1951.
7. *J. Observateurs* **34**, 85, 1951.
8. *Pub. Astroph. Obs. Potsdam* **24**, No. 75, 1920.
9. *B. A. N.* **1**, 193, 1923.
10. *Ann. Stw. Leiden* **18**, Part 2, 9, 1937.
11. Strand, *Ann. Stw. Leiden* **18**, Part 2, 70, 1937.
12. Strand, *A. J.* **57**, 97, 1952.
13. *Amer. Phil. Soc. Yearbook*, 1941, p. 95-96.
14. Boss, *General Catalogue of 33342 Stars* **1**, 49, 1937.
15. *Ibid.*, App. II, 151, 1937.
16. Lippincott, *A. J.* **55**, 236, 1951. Originally discovered by Reuyl, *A. J.* **45**, 133, 1936.
17. v. d. Kamp, *Proc. Amer. Phil. Soc.* **88**, 373, 1944.
18. *Ap. J.* **88**, 477, 1938.
19. Strand, *Ap. J.* **113**, 1, 1951.
20. Hall, *A. J.* **57**, 147, 1952.
21. Silbernegel, *A. N.* **233**, 168, 1928.
22. *A. J.* **55**, 215, 1951.
23. *Ann. Harv. Col. Obs.*, **11**, 105, 1879; **64**, 159, 1912; **69**, 180, 1913.
24. *Pub. Astr. Obs. Univ. Tartu* **25**, 6, 1924.
25. *Ann. Astroph.* **14**, 85, 1951.
26. *Pub. A. S. P.* **47**, 24, 1935.
27. *Ann. Obs. Strasbourg* **5**, Fasc. 1, 1948.
28. *A. N.* **273**, 253, 1943.
29. *Uppsala Astr. Obs. Ann.* **2**, No. 2, 1947.
30. *B. A. N.* **10**, 169, 1946.
31. *Ann. Obs. Strasbourg* **5**, Fasc. 3, 1951.
32. *Zs. Astroph.* **24**, 263, 1948.
33. *Veröff. Bad. Land. Stw. Heidelberg (Königstuhl)* **15**, No. 6, 1947.
34. *Ap. J.* **117**, 361, 1953.
35. *Carnegie Inst. Washington Year Book*, No. 50, 8, 1953.
36. *Union Obs. Circ.*, **100**, 466, 1938.
37. *J. Observateurs* **33**, 1, 1950.
38. Hertzprung, *B. A. N.* **6**, 57, 1930.

Dearborn Observatory,
Northwestern University,
Evanston, Ill.,
1953 September.