In general terms, the problem is this: The spread and excursion of a stellar image at the focus of a large telescope is determined almost solely by atmospheric seeing. During ordinary good seeing, star images are smeared out into blobs roughly one second of arc across, which corresponds to 0.1 mm at the 200-inch prime focus; this is at least 6 times larger than the resolving power of the photographic emulsion and 40 times larger than the optical resolving power of the telescope. The detection of a faint star image depends accordingly upon whether the picture receiver, which might be either a photographic emulsion or an electronic device, is capable of distinguishing the surface brightness within the small area occupied by the smeared image from the surface brightness of the sky background surrounding it. If we want to detect fainter images than we can now reach, we must improve the discrimination of faint images against the sky background, which in engineering terms means that we must improve the signal-to-noise ratio of the receiver. Merely increasing the sensitivity of the receiver, although potentially useful in high-dispersion spectroscopy, will not solve the image detection problem.

The present photographic limit of the 200-inch telescope can be very nicely understood on a signal-to-noise basis. After a suitable allowance is made for the statistical saturation due to the overlapping of photographic grains on a so-called full exposure, actual grain counts indicate that we should theoretically expect the threshold of a fully exposed Eastman 103a-O plate at the 200-inch prime focus to be in the vicinity of the 24th magnitude. This is in very satisfactory agreement with observation; the faintest photographically detected image thus far measured directly at the telescope with the photoelectric photon counter is found to be 23.8 magnitude.

Since the percentage statistical fluctuation in background grain counts is inversely proportional to the square root of the number of grains per background "picture element," the only way to extend the present threshold is to produce, or potentially be able to produce, more grains per element and at the same time avoid saturation due to the overlapping of grains either by making the grains smaller or by devising a scheme for subtracting a uniform excess portion of the background before the picture is photographically recorded. For practical reasons, both of these methods call for electronic techniques which take advantage of the high quantum efficiency of photoelectric cathodes. One electron is ejected

from a photoemissive cathode for every ten eligible photons incident upon it, as compared to one photographic grain being rendered developable for every thousand eligible photons incident upon an emulsion during unaided photography. If a photoemissive cathode is used in combination with photographic registration of the picture, each photoelectron can be caused to produce one or more photographic grains, making the combined process 100 times more efficient than unaided photography. The general feasibility of these television-like techniques has already been adequately demonstrated; it is now primarily a question of determining which particular method is best for the image detection problem. If the full advantage of photoelectric quantum efficiency could be ideally utilized, it would theoretically accomplish what unaided photography would need a 2000-inch telescope at $F/3.67$ to do.

The Carnegie Corporation recently put up funds for a concerted attack on this problem. Organizations at present cooperating in this project include the Carnegie Institution's Department of Terrestrial Magnetism, the National Bureau of Standards, the U. S. Naval Observatory, and the Mount Wilson and Palomar Observatories. Tentative plans are also being made to hold an image-tube symposium in connection with the 1955 I.A.U. meeting at Dublin.

*Mt. Wilson and Palomar Observatories,
Pasadena, Calif.*

Bidelman, W. P. The space distribution of supergiant stars of spectral types later than B.

The known stars of spectral types A5 and later and of luminosity classes Ia and Iab have been investigated to determine whether their space distribution is similar to that of the aggregates of early-type stars discussed by Morgan, Whitford, and Code.¹ While there is no doubt that these later-type supergiants are largely located in the spiral arms, they are not all intimately associated with the early-type aggregates. The η and χ Persei and the χ^2 Orionis groups include a number of M-type supergiants, and several other aggregates probably have middle- and late-type supergiant members, but a number of the aggregates do not seem to have any associated late-type stars of high luminosity. The general conclusion is that the supergiants of later type favor those early-type aggregates which also contain supergiant stars of early type, rather than those consisting predominantly of O-type main sequence stars.

Trumpler and Weaver have noted that the