scribed by M. Czerny and A. F. Turner,⁴ but the mirrors and grating are now placed in a vacuum tube 52 feet long and 4 feet internal diameter. Other tube specifications are: Rolled and welded plate, thickness $\frac{1}{4}$ inch, reinforced with rings every four feet. The end plates are $1\frac{1}{4}$ inches thick and are heavily reinforced with 6-inch deep ribs of 1-inch plate welded to the inside faces. Light from the entrance slit located behind a quartz field lens which also acts as a window to the vacuum tank is collimated by a 12-inch spherical mirror and returned to the grating located several feet inside the head of the tube. The dispersed beam is focused on the second slit by a second 12-inch spherical mirror and after passing through a quartz window is received by a photomultiplier. A third concave 12-inch mirror at the far end of the tube is arranged so that a short region of the spectrum can be returned to a photographic plate. All mirrors, and the grating table, are provided with complete motions in all degrees of freedom through the use of electrical motors. The final adjustments can be made by push buttons after the tank has been pumped down and assumed its slightly altered shape under the atmosphere pressure load of 700 tons. The head of the spectrograph is to be provided with air locks at every point. The adjustment and repair of the slits, or of the photocell, are thus facilitated without destroying the vacuum of the main tank. A special cassette and air lock will permit a quick plate change to be made.

The tank, constructed by Whitehead and Kales of Detroit, proved to be vacuum-tight on erection. The first pump-down with a Kinney vacuum pump, Model VSD-8811, of 45 cubic feet displacement per minute, required one hour to reach 1 cm pressure—1 mm after two hours. A pressure of 70 microns has been obtained and it is expected that still lower values will be achieved after the inside surfaces, painted with red lead, have occluded their more volatile constituents. Tests have shown that the rate of pressure rise on the tank is so small, 0.1 mm rise per 24 hours, that it will be possible to pump out the tank at night and then to shut off the pump during the operating day. This is a necessary requirement as the pump, though on an isolated pier with flexible connection to the tank, carries a perceptible vibration to the optics.

This spectrograph will allow the use of one of Babcock's superb gratings in the fifth order, and should allow the grating to develop its full re-

solving power of about 600,000. The linear dispersion in the fifth order will be 6.92 mm per angstrom at 5000 Å. This dispersion should make possible faint line profile work containing a relatively small instrumental contribution. The scattered light is to be evaluated by interferometric and other methods. Precision measurements of profiles of the stronger lines by photoelectric methods requires an exceptional sky. Our experience has been that sky transparency over long intervals of time is so rarely obtained that means of monitoring the background intensity is very desirable. For this purpose we are planning to use a modification of the system employed by Hiltner and Code.⁵ One photocell will monitor a 200 Å interval as passed by the predisperser through the first slit, and the other will record the spectrum through the second slit. Reflecting slits of stellite will allow one to guide accurately on details seen through a Lyot filter which will present a field bisected by the slit.

1. R. R. McMath and O. C. Mohler, *J. Opt. Soc. Amer.* **39**, 903, 1949.
2. *Ap. J.* in press.
3. *Ap. J.* **116**, 1, 1952.
4. *Zs. Phys.* **61**, 792, 1930.
5. *J. Opt. Soc. Amer.* **40**, 149, 1950.

*McMath-Hulbert Observatory,
University of Michigan,
Pontiac, Mich.*

Menzel, Donald H. and Fred L. Whipple. The case for H_2O clouds on Venus.

There has been a tendency in recent years for astronomers to subscribe to the theory that the clouds of Venus consist of dust rather than H_2O . The dust hypothesis, however, is inconsistent with Lyot's polarization measures, which, on the other hand, accord well with the behavior of light scattered by H_2O clouds.

Dunham has estimated that the total amount of water vapor above the clouds of Venus does not exceed 2 to 5 per cent of that above Mount Wilson. Recent measures of the planet's temperature by William Sinton indicate a nearly constant day-night value of -39°C . A saturated convective atmosphere having this temperature at the visible surface will contain appreciably less than 2 per cent the water vapor of a saturated atmosphere at 6°C above Mount Wilson. Therefore, failure to detect water vapor in the spectrum of Venus is at present no argument against the H_2O character of the clouds.

A thick atmosphere consisting largely of CO_2 could not exist on an earth-like planet with con-