Mumford, George S, III. The motions and distribution of red dwarf stars.

An analysis of the peculiar motions of 825 dwarf stars, spectral types dK8 to dM5, discovered spectrophotometrically at the Leander McCormick Observatory indicates no asymmetry in the distribution of peculiar velocities in the direction of the galactic center and anti-center; while the velocities in the direction of rotation and the opposite direction show a pronounced asymmetry traceable to velocities considerably less than 50 km/sec. The largest velocity in the direction of rotation is 54 km/sec, while that in the opposite direction is 171 km/sec.

The dispersions of the peculiar velocities along the axes were computed. The quantity $M\sigma^2$, where M is the mass, σ the total dispersion, is very nearly constant within the range of classes investigated here.

Projections of the positions of the stars on the galactic plane and perpendicular to it show a remarkably even distribution of these stars in space. From the absence of clustering we may conclude that the red dwarfs must be predominantly old stars.

*Leander McCormick Observatory,
Charlottesville, Vir.*

Münch, L. and Blanco, V. M. Red stars near η Carinae.

As part of a program of observations of the southern Milky Way with infrared sensitive spectroscopic plates, the region near η Carinae has been studied at Tonantzintla. In an area of 25 square degrees, about 800 M-type stars with magnitudes brighter than 10.4, infrared, were observed. A list of these stars with charts for their identification is under preparation. Some of the M stars appear to be supergiants, and these are found associated with the OB star groups studied by Münch and Morgan (1953). A group of 10 carbon stars was found centered approximately on η Car. Lack of accurate magnitudes, luminosities and absorption values do not permit us to tell whether these stars are associated with the η Car nebulosity.

Four S-type stars were found in the region, two of which are embedded in the densest regions of the nebulosity. Around each of these a small patch of bright nebulosity may be observed. Tentative luminosities for the S stars obtained by using the distance and absorption published by Bok and van Wijk (1952) are in agreement

with those of giant M-type stars, in accordance with Keenan's (1954) deduction and with the results obtained for π' Gruis (Feast 1953).

A group of 13 stars was also observed in which OI or CaII infrared emission lines near 8450Å are suspected. These stars were mostly found in the darkest absorption regions of the area. Some of the stars are probably A-type main sequence objects, while others appear to be M giants.

Bok, B. J. and van Wijk, U. 1952, *A. J.* **57**, 213.

Feast, M. W. 1953, *M. N.* **113**, 510.

Keenan, P. C. 1954, *Ap. J.* **120**, 484.

Münch, L. and Morgan, W. W. 1953, *Bol. Tonantzintla y Tacubaya* No. 8, 26.

*Tonantzintla Observatory,
Tacubaya, Mexico,
and Warner and Swasey Observatory,
East Cleveland 12, Ohio.*

Nicholson, Seth B. Short-period fluctuations in solar activity.

Superposed on the 11-year variation in the frequency of sunspots are fluctuations with amplitudes of 20 to 100% and with periods ranging from 8 to 12 months. The periods vary by about the same per cent as do those of the main sunspot cycle. The cause of these short period fluctuations is as unknown as is that of the 11-year cycle, but if one wishes to predict the number of sunspots in any month a year or less in advance, the predictions will be greatly improved, if the fluctuations are taken into account.

*Mount Wilson and Palomar Observatories,
Carnegie Institution of Washington,
and California Institute of Technology.*

Oke, J. B. The luminosity classification of late-type stars using microphotometer tracings.

Calibrated spectrograms having a dispersion of 33 Å/mm have been obtained for many late-type stars. From microphotometer tracings of these spectrograms ten pairs of lines were selected which showed marked changes with luminosity. For each pair of lines a quantity R , defined as the ratio of the logarithms of the depths of the two lines, was obtained. The weighted mean $\bar{R}$ was calibrated to absolute visual magnitude M_v using 25 G5 to K0 stars with the best trigonometric or cluster parallaxes. The results indicate that the standard deviation of the absolute magnitude as obtained from $\bar{R}$ and the calibration curve is approximately 0.2 mag. if two spectrograms are measured.

*David Dunlap Observatory,
Richmond Hill, Ont.*