provides a very large aperture area per unit cost. The design is especially well suited for background mapping, source location, and spectrum observations. Although essentially a meridian-transit instrument it can be arranged to track objects near the meridian for moderate intervals.

Downcoming celestial radio waves are deflected by the tiltable flat-sheet reflector into the paraboloid and thence to a horn or other antenna at the prime focus. The reflecting surface of both paraboloid and flat sheet consists of coarse mesh wire screen. The only moveable structure is the flat-sheet reflector. Since a flat surface can be very easily checked for any unevenness, it is not necessary that the tiltable reflector be a rigid structure. This results in a large saving in cost. The reflector is tilted by a number of independently operated truss members and the desired degree of flatness of the reflector obtained by adjusting them individually while monitoring or observing the reflector for flatness with a small optical telescope set up near one end.

It is much more difficult to maintain tolerances on a curved surface such as a paraboloid, especially if it is moveable. This problem is avoided and the cost correspondingly reduced by making the paraboloid fixed. In this way the ground itself provides the principal mechanical support for the paraboloid and the amount of structure required to maintain the paraboloid tolerances is a minimum. Movement is required of only the simple flat reflector and only 65 degrees of tilt of this reflector are necessary to cover nearly all of

the observable sky. By making the length of the paraboloid considerably greater than its height there is an additional large saving in cost since the expense of a structure is approximately proportional to the cube of its height. Furthermore, the lower overall height will make the telescope much less susceptible to interfering radio signals of terrestrial origin, since such signals increase greatly in strength with height above the ground.

To test the design a scale model has been constructed measuring 12 feet across and operating at a wave length of 1.25 centimeters. By scaling both the physical size and wave length in this manner, patterns can be measured that will duplicate those of a full-size system. Measured patterns and performance of the model have been so satisfactory that the model has been put into use as an actual radio telescope at 1.25 cm wave length for solar and other observations. Specifically the model duplicates the performance of a telescope with parabolic reflector 700 feet long by 70 feet high operating at a wave length of 70 cm. A grant of \$48,000 from the National Science Foundation has been received for the construction of the first half of a telescope of these dimensions.

REFERENCES

- Kraus, J. D. 1955, *Sci. Amer.* **192**, 36.
 Kraus, J. D. and Nash, R. T. 1955, *Ohio State Univ. Radio Obs. Rep.* No. 5 (Nov. 22).
 Mills, B. Y. and Little, A. G. 1953, *Australian J. Phys.* **6**, 272.
*Ohio State University,
 Columbus, Ohio.*

ABSTRACTS *

Aller, Lawrence H. Chemical composition of the planetary nebulae.

An analysis of the chemical composition of the planetary nebulae NGC 1535, 2022, 2165, 2440, 2392 and 7662, based on spectrophotometric observations by Minkowski and the writer at the Mt. Wilson Observatory, is reported. The method is very similar to that employed for NGC 7027. The effects of filamentary structure and the distribution of atoms among different stages of ionization introduce complications that appear particularly serious for elements such as nitrogen which appear only in the lowest stages of ionization. The adopted mean composition, including

NGC 7027, by relative numbers of atoms is as follows:

	log N		log N		log N
H	7.23	O	4.0	S	2.95
He	6.13	F	0.6	Cl	1.53
N	3.7:	Ne	3.18	A	2.11

No convincing evidence is found for variations in composition from one object to another. The scatter in the individual values is more likely due to variations in physical conditions.

*University of Michigan Observatory,
 Ann Arbor, Mich.*

* Of papers presented at the Ninety-fourth Meeting of the American Astronomical Society, Columbus, Ohio, March 21-24, 1956.