out to the surface. The writer feels that the humps occurring on the ascending branches of the radial-velocity curves are to be associated with this reflected pulse, the direct pulse having appeared near the phase of greatest velocity of expansion.

Without speculating further on the peculiarities of Cepheids with periods around 10 days, we may suggest that the hump phenomenon is restricted to the period interval 6–20 days for two reasons. The hotter Cepheids, with periods less than 6 days, possess ionization zones which are so shallow that the product PV is insufficient to produce appreciable hump energy. On the other hand, the cooler Cepheids, with periods greater than 20 days, possess ionization zones which are so deep that nonadiabatic effects are not important in the critical region.

Ledoux, P. 1953, *Ciel et Terre* **69**, 70.

Dumézil-Curien, Mme. P. 1954, *Ann. Astroph.* **17**, 197.

*Institut d'Astrophysique,
Cointe-Sclassin, Belgium.*

Wild, J. Paul and Zirin, Harold. On the association of solar radio emission and solar prominences.

We have examined prominence cinematograms made at Sacramento Peak and Climax in the years 1949–1955 and compared them with the solar radio records at 167 Mc/s. No close connection was found between limb events and radio events, but some eruptions were found to be associated with simultaneous radio bursts. Three such cases are discussed in detail. These appear to produce a series of storm bursts of spectral type 1. The radio emission appears to originate from points in the corona at the leading edge of the disturbance.

The limb passages of large sunspot groups in 1949–1955 were studied on both cinematograms and single exposure surveys with the view of finding a criterion for radio-active spots. It was found that spot groups showing looped prominences and downward streaming from the corona showed a marked tendency to produce radio storms. Of 14 spots producing enhanced radio levels, 12 showed this criterion; 9 out of 10 large spots ($>800 \times 10^{-6}$ of the disk) with loops and streamers produced enhanced radio emission. This result is ascribed to the fact that the seat of such storms must lie in the corona and in the presence of strong ordered magnetic fields; the existence of such conditions is indicated by the presence of looped prominences.

This work was supported by the Air Force Cambridge Research Center, Geophysics Research Directorate, through contract AF 19(604)-969 with the High Altitude Observatory.

*Commonwealth Scientific and Industrial
Research Organization,
Sydney, Australia,
and High Altitude Observatory,
University of Colorado,
Boulder, Colo.*

Wilson, Raymond H., Jr. Magnetic damping of rotation as a factor in stellar evolution.

A thin ring of conducting matter rotating in a magnetic field about an axis in its plane experiences a braking couple given by a well-known function of the area of the ring, its inductance, resistance, and angular velocity, and the effective strength of the magnetic field. Integrating this as an element of a conducting sphere leads to the approximate formula for its angular velocity ω as function of time t in seconds:

$$\log_e \frac{\omega_0}{\omega} = \frac{\sigma \mu^2 H^2}{\rho} (t - t_0).$$

Here σ (cgs, EMU) is the conductivity of the sphere, μ its magnetic permeability, and ρ its density, while H is the external magnetic field strength in gauss.

Such magnetic damping would tend to reduce rotation of all celestial objects in a magnetic field. As explained by Chandrasekhar and Fermi (1953) all present evidence indicates a magnetic field of 7×10^{-6} gauss in the sun's galactic region, due to motions of the ionized galactic plasma. The action of this field on a star like the sun depends partly on its conductivity. A reasonable estimate of the effective conductivity of the sun would be the logarithmic mean between center and photosphere: $\sigma = 10^{-6}$ cgs EMU (Spitzer 1954). Then by the above formula we may derive the sun's period of rotation at, say, $t_0 = 5 \times 10^9$ yr. = 1.6×10^{17} sec. ago. Its mean density in this interval could be estimated as perhaps half that at present = 0.7. Assuming the galactic field to have remained constant, the effective component would be about 4×10^{-6} gauss. The result is that $\log_e (\omega_0/\omega) = 3.7$ so that the rotation period at t_0 would have been 25/40 day = 15 hours. This is on the order of rotational velocities of young stars of early main-sequence type, from which, presumably, the sun and other similar stars have evolved (Struve 1950).

Since the conductivity of a star with increasing depth below the photosphere seems to increase more rapidly than its density, the rotation of the interior would be slowed down more rapidly than that of its surface. A result for a fluid or partly fluid sphere would be turbulence, and more rapid rotation at the equator, such as is observed on the sun and the planet Jupiter.

In planetary systems and double stars the action of any magnetic fields belonging to the separate members would add its effect to that of the general galactic field. For instance, in the case of close spectroscopic binary stars (Struve 1950), the total effective field might be on the order of 10^{-3} gauss. Assuming other data as above, except the conductivity like that of the solar photosphere, 10^{-8} cgs EMU, the time required for reduction of rotation by a factor of 100 turns out to be 10^7 years. This is the age estimated for such early-type stars (Struve 1950). Thus the theory of magnetic damping alone could possibly explain the abnormally slow rotation of such binary stars compared to that of single stars of the same age.

Struve, O. 1950, *Stellar Evolution* (Princeton Univ. Press), pp. 130, 83, 244.

Chandrasekhar, S. and Fermi, E. 1953, *Ap. J.* **118**, 113.

Spitzer, L. 1954, *Smithsonian Physical Tables*, p. 502.

*Naval Research Laboratory,
Washington, D. C.*

Wright, K. O. The chromospheric spectrum of 31 Cygni.

At the 1951 eclipse of the important system, 31 Cygni, a series of high-dispersion grating spectrograms in the region $\lambda\lambda 3700-4500$ was obtained during ingress. Sharp, chromospheric absorption lines of FeI could be detected at least twelve days before totality, and the intensities of these lines have been analysed. Turbulent velocities derived from curves of growth range from 8 to 15 km/sec. and may increase somewhat with height in the atmosphere. Excitation temperatures increase with height from 2500°K at -2^d to 4500°K at -10^d . The number of neutral iron atoms ob-

served in the column traversed by the light of the small secondary star decreases by a factor of 3000 in the same time. The analysis has been made by measuring total absorptions on intensity tracings and relating them to the apparent continuum of the B-type star at each wave length. The light ratio of the two stellar continua was obtained from a comparison of intensities in the composite spectrum taken well out of eclipse with those in the totally-eclipsed spectrum. Occultation effects could be detected at least two days before totality.

*Dominion Astrophysical Observatory,
Victoria, B. C.*

TITLES OF ADDITIONAL PAPERS PRESENTED AT THE MEETING IN DELAWARE, OHIO

Bell, G., Davis, M. H., King, R. B. and Routly, P. M. The absolute f -values of Copper I, Manganese I and Iron I.

Cameron, Donald and Nassau, J. J. Spectral classification of red variables along the galactic equator.

de Vaucouleurs, Gerard. Magnitudes and colors of the Magellanic clouds.

Fernie, J. D. The southern open cluster, NGC 2547.

Gallet, Roger M. and Bowles, Kenneth L. Some properties of the radio emissions of Jupiter.

Grant, Gordon. The eclipsing systems λ Tauri and RW Tauri.

Irwin, John B. A photoelectric study of southern Cepheids. McMath, R. R., Mohler, O. C., Pierce, A. K. and Goldberg, Leo. The contribution of the chromosphere to the Fraunhofer spectrum.

Mayall, Margaret W. Predicted and observed light curves of nine long period variables.

Odgers, G. J. Variations of BW Vulpeculae during 1955.

Petrie, R. M. Values of Oort's constant "A" derived from some B stars.

Pierce, Keith and McMath, R. R. Instrumental profile of the vacuum spectrograph.

Prendergast, Kevin H. Equilibrium of a fluid sphere with a magnetic field.

Rabe, Eugene K. On the origin of the Kirkwood gaps and the minor planet families.

Roman, Nancy G. and Haddock, Fred T. A model for non-thermal source spectra.

Schwarzschild, Martin, Howard, R. and Harm, R. A model for the sun with convective envelope.

Slettebak, Arne. Line broadening in the O- and early B-type stars.

Spitzer, Lyman, Jr. Properties of an interstellar galactic corona.

Urey, H. C. and Donn, B. Chemical heating for comets and meteorites.