order exponential integrals were carried out on the 1953 solar limb-darkening measures by Pierce (1954) throughout the visual and infrared regions. Each of the integro-exponential forms represents very well the limb darkening in the observed range, $\mu = 1.0$ to $\mu = 0.10$.

An indeterminacy formerly resulting from the inclusion of more than one exponential integral term in the expression for the source function is replaced by a continuum of solutions which are obtained by fractional superposition of the separate solutions described above.

Use of exponential integrals of non-integral order (Pecker 1952) enlarges the range of boundary conditions accessible with these analytic forms. The somewhat greater freedom in imposing the boundary conditions is shown by comparison with Michard's (1953) treatment of the limb-darkening data.

The distribution of temperature with optical depth for the 1953 measures is shown for the wave-length region $\lambda\lambda 6000-24000$.

It appears that the ambiguity of the $\tau - \lambda$ scale outside of the range of observation, i.e. at $\tau < 0.2$ and > 1.0 , may be removed by integration of the solar model at a number of wave lengths. The isobars of gas pressure would be expected, in all but possibly the uppermost layers, to parallel the isotherms from wave length to wave length. The suitability of the $\log \tau$ scale for the entire integration is examined in view of the greater weight it allows the line-forming regions.

A perturbation method is indicated for the solution of the equation of transfer to make it applicable to the case of the spherical sun.

Michard, R. 1953, *Ann. Astroph.* **16**, 217.

Pecker, J.-C. 1952, *Contr. Inst. Astroph.*, Ser. A, No. 132.

Pierce, A. K. 1954, *Ap. J.* **120**, 221.

*Ladd Observatory,
Brown University,
Providence, R. I.*

Newkirk, Gordon, Axtell, James and Wlérick, Gérard. Studies of the electron corona.

During the period September 11, 1956 to October 8, 1956 observations of the polarized radiation from the K corona were secured on a total of twelve days. Although the data are not yet on an absolute standard, polar diagrams of the relative polarization product for successive days show the electron corona rotating with the sun.

A region of enhanced brightness which ap-

peared on the west limb on September 17 and on the east limb on October 2 was remarkable for having its angle of maximum polarization displaced about 35° south of its normal position during the west limb passage. On September 17 a surge at the same position angle was inclined approximately the same amount to the solar limb. A tentative explanation for the departure of the polarization angle from normal has been found in the synchrotron radiation from relativistic electrons.

The present data on the K corona are not yet extensive enough to reach any definite decision concerning its connection with geomagnetic storms.

This research has been supported by the U. S. National Committee for the International Geophysical Year and by the HAO Institute for Solar-Terrestrial Research.

*High Altitude Observatory,
Boulder, Colo.*

Odgers, G. J. and Kushwaha, R. S. Shock waves in stellar atmospheres.

Owing to the decreasing density, an acoustic disturbance proceeding outwards from the photosphere of a star rapidly develops into a wave of large amplitude and then into a wave of discontinuity or shock wave. Observations of the β Canis Majoris star BW Vulpeculae show clearly that material is ejected from the surface at velocities corresponding to Mach numbers of nearly six. Clearly such motion corresponds to the dynamical effect of a shock wave.

Equations have been developed to describe the passage of a shock wave through a stellar atmosphere. The usual conservation laws at the shock front are used, but in addition the decay of the wave is accounted for by considering the rate of dissipation of energy at the front due to processes of an irreversible character which increase the entropy and enthalpy of an element of gas and so lose the energy available for further propagation of the wave. The equations are exact but in order to obtain solutions an approximation is made regarding a certain energy-time integral occurring therein which is equivalent to supposing that the pressure-time curve of the shock wave for a given element maintains the same shape as the wave decays.

The resultant equations can then be solved exactly in several cases of astrophysical interest; the most important case is shown to be that of