THE ASTRONOMICAL JOURNAL

FOUNDED BY B. A. GOULD

PUBLISHED BY THE AMERICAN ASTRONOMICAL SOCIETY

VOLUME 62

1957 July ~ No. 1249

NUMBER 5

ABSTRACTS *

Athay, R. G. and Zirin, Harold. On the relative intensities of chromospheric Balmer and Paschen lines.

The flash spectrum of the solar chromosphere includes hydrogen Balmer and Paschen lines originating in common upper levels. We denote the intensity of the Balmer line arising from the transition n to 2 by B_n and the intensity of the Paschen line arising from the transition n to 3 by P_n . If all terms of the same n are populated in proportion to their statistical weights and if self-absorption is small, $B_n/P_n \sim 3.3$ for $n \geq 9$. Self-absorption will tend to reduce the observed ratio of B_n/P_n .

From spectrograms obtained at the 1952 eclipse, we find $B_n/P_n \sim 8$ for $n \geq 14$. The discrepancy between observed and predicted ratios of B_n/P_n is too large to be accounted for by photometric errors. This may be illustrated by comparing continuum intensities in the ultraviolet and infrared and by comparing intensities of the H and K lines with the intensities of the infrared $CaII$ triplet. The relative continuum intensities for the corona, derived from the same spectrograms, follow the photospheric curve. Furthermore, the $CaII$ lines have very nearly the predicted relative intensities.

If we discount photometric errors, then the observed ratio of B_n/P_n implies that some of the multiplets normally present in the Paschen lines are suppressed. In fact, it seems necessary to assume that only the s , p and d terms are populated. In this case, the predicted ratio of $B_n/P_n \sim 6$, which is in satisfactory agreement with the observations. The above assumption can be justified if it can be shown that most of the electrons in the upper term arrive via excitations

from the $1s$, $2s$ and $2p$ terms. Quantitative estimates of rates of excitations from these low-lying terms and rates of three-body recombinations from the continuum using the Thompson collision cross-section indicate that three-body recombinations form the primary source of electrons for the upper levels. This would imply that all of the multiplets are populated and B_n/P_n should be ~ 3.3 . Thus, the Thompson collision cross-section leads to a contradiction between predicted and observed ratios of B_n/P_n . Observations of the Paschen-Balmer ratio in prominences are now in progress at Climax.

This work was carried out under the sponsorship of the Office of Naval Research and the Geophysics Research Directorate, Air Force Cambridge Research Center.

*High Altitude Observatory,
Boulder, Colo.*

Beals, C. S. Results of drilling operations at the Holleford crater.

The Holleford Crater, lat. $44^\circ 27'$, long. $76^\circ 38'$; diameter 1.46 miles, present depth 100 ft, reported as of possible meteorite origin by Beals *et al.* (1956), has been drilled with diamond coring bits in three locations designed as Holes (1), (2), (3), at distances of 1400 ft., 2500 ft. and 3780 ft. from the center, to depths of 1128 ft., 1486 ft. and 500 ft. respectively. The general area of Holleford is covered with paleozoic sediments which nearly fill the crater and cover the rim. In all three holes the contact between the sedimentary rocks and what may reasonably be assumed to be the floor or rim of the crater is

* Of papers presented at the Ninety-seventh Meeting of the American Astronomical Society, Cambridge, Mass., May 8-11, 1957.