prominent. Emission nebulosities are situated around NGC 2175, BD +20°1293, and HD 254577.

Analysis of the data reveals that there are probably two associations in this region, one having a distance modulus 10.3 mag. (1150 pc) and the other 11.5 mag. (2000 pc) rather than a single one at 1400 parsecs according to the earlier work. In obtaining these values, photoelectric measures of reddening for each of 24 stars having good spectral classifications were used to arrive at dependable apparent magnitudes corrected for absorption. Further confirmation of these values was secured from the data of Crawford *et al.* (1955).

- Crawford, D., Limber, D. N., Mendoza, E., Schulte, D., Steinman, H. and Swihart, T. 1955, *Ap. J.* **121**, 24.
 Keenan, P. C. and Morgan, W. W. 1951, *Astrophysics*, ed. J. A. Hynek (New York: McGraw-Hill Co.), p. 23.
 Merrill, P. W. and Burwell, C. G. 1933, *Ap. J.* **78**, 87.
 —. 1943, *ibid.* **98**, 153.
 —. 1949, *ibid.* **110**, 387.
 —. 1950, *ibid.* **112**, 72.
 Morgan, W. W., Code, A. D. and Whitford, A. E. 1955, *Ap. J. Suppl.* **2**, 41.
 Morgan, W. W., Whitford, A. E. and Code, A. D. 1953, *Ap. J.* **118**, 318.

Arthur J. Dyer Observatory,
 Vanderbilt University,
 Nashville, Tenn.

Hawkins, Gerald S. Variation in the occurrence rate of meteors.

Radio-echo observations are used to estimate the total influx of meteors on the earth's surface. The approximate distribution of sporadic meteor radiants is deduced by a statistical method, and from this distribution the diurnal variation in the occurrence rate of meteors is computed for geographical latitudes 0°, 30°N., 50°N., and 80°N. The variation is greatest at the equator and becomes progressively smaller at the higher latitudes.

Harvard College Observatory,
 Cambridge, Mass.

Heeschen, D. S. A 21-cm line study of extra-galactic nebulae.

An investigation of 21-cm line emission from extra-galactic objects has been initiated with the Agassiz Station equipment. Preliminary observations of the Coma Cluster of galaxies with the 24-foot radio telescope of Harvard Observatory show that this cluster contains about 10^{14} solar masses of neutral hydrogen, and that the hydrogen has a dispersion in radial velocity of 500 km/sec. The implications of these observations with

respect to the mass-luminosity ratio of the cluster, the effects of collisions of galaxies in the cluster, and the relative amount of hydrogen in Population II objects are discussed. Other objects on the observational program include the Virgo and Hercules clusters, the elliptical nebulae M 32, NGC 185 and NGC 205, and the spiral galaxies M 31 and M 33. Hydrogen line emission from most of these objects should be observable with a 60-foot radio telescope, and a meaningful upper limit may be obtained to the amount of hydrogen in those which cannot be detected. The elliptical nebula M 32, for example, should be observable if as much as 0.005 of its total mass is neutral hydrogen. One objective of the program is to determine what relationships, if any, exist between the total mass, hydrogen mass, mass-luminosity ratio and population characteristics of the clusters and field nebulae. The mass of neutral hydrogen and, in some cases, the total mass may be determined from observations of the 21-cm line. Preliminary results are described.

Associated Universities, Inc.,
 350 Fifth Ave.,
 New York, N. Y.

Hoffleit, Dorrit. Variable stars in Variable Star Field 193.

Magnitudes on 30 or more plates of the Harvard MF series, taken with the 10-inch Metcalf telescope, have been estimated for 450 unpublished variable stars in VSF 193 at 18^h 24^m, -23° 3' in Sagittarius. Periods have been determined for 65 of the long-period variables, or about half the total number of this type I discovered in the region. Their distances, based on the Wilson-Merrill period-luminosity relation, range from one to 13 kpc. Only those with periods shorter than 250 days have been found at distances beyond the galactic center. Selection effects are discussed which hinder the discovery of the stars of longer period either at great distances or in dense star-fields.

Cambridge, Mass.

Hosfeld, Roger and Hynek, J. A. A directional stellar scintillometer.

A small, compact and transportable instrument for the rapid determination of the direction and velocity of the stellar shadow-band pattern, responsible for stellar scintillation, consists of a rectangular portion of an off-axis parabolic mirror, of 2-foot focal length, rotatable around