On this model, prominence activity associated with spot groups should originate in the surrounding material rather than in the spots themselves. Careful comparison of the relative location of spots and prominences will provide an observational check. Also correlations should be sought between the detailed character of spot groups and the types of associated prominences.

The enhanced flux of mechanical energy involves increased dissipation and hence above-average heating of the chromosphere and corona near spots. These effects should appear, as in fact they do, before a spot has surfaced. Prominences visible as dark filaments on the disk may similarly be associated with submerged magnetic fields. Although the main energy source for prominence activity is probably mechanical, the actual motions will be influenced strongly by local magnetic fields and may, in turn, also influence those fields.

It is our view that the normal convection of the quiet sun is sufficient to produce both chromosphere and corona. The tenuous extremities of the latter may, as Chapman has suggested, reach beyond the earth. Shock waves, arising from the activity near spots and amplified by passage through a medium of progressively lower density, can produce many of the features of solar prominences. A knot ejected from an active region will exhibit a Doppler displacement even if the matter composing it changes constantly. The phase and amplitude cross-modulation, referred to in the accompanying paper, permits propagation velocities in excess of the sound velocity in the medium.

Stars having a general magnetic field sufficiently intense to inhibit atmospheric convection should possess extended envelopes to make up for the lower efficiency of energy transport. Although no detailed models have been worked out, we note that stellar magnetic fields and the mechanical disturbances associated with them may be responsible for certain varieties of stellar variability, especially those involving expulsion of matter from the surface and asymmetrically distended atmospheres.

The research in this paper has been supported in part by the Geophysics Research Directorate, Air Force Cambridge Research Center, Cambridge, Mass., under contract with Harvard University.

Krook, Max and Wild, Paul. 1957, *A. J.* 62, 22.

*Harvard College Observatory,
Cambridge, Mass.*

Krook, Max and Wild, Paul. On the theory of solar radio bursts.

If a sharp pulse of small amplitude is generated in an ionized gas, it propagates as two separate pulses—a slow one carried by electrons and ions together, and a fast one carried by the electrons alone. For pulses of large amplitude we have to consider the additional effects arising from cross-modulation in both amplitude and phase of the Fourier components. Various modes of behavior are then possible, including the generation and propagation of shock waves. Wild has classified solar radio bursts into three main types. The behavior of his types II and III is consistent with the interpretation that they are associated with shock waves in the electron-ion gas, and in the electron gas, respectively. This interpretation is supported by examples from Wild's "rogues gallery" of solar bursts, which includes combinations of events that are to be expected on the basis of cross-modulation.

The burst disturbances generally originate near sunspots. We suggest that they may arise from instabilities that are just enhanced forms of the instability responsible for the existence and maintenance of the chromosphere and corona. The fluctuating character of the outer boundary of the convective zone results in the transfer of mechanical energy through the overlying material and its eventual dissipation there. Biermann has suggested that the magnetic field of a spot inhibits convection. The energy prevented from flowing through a sunspot leaks out through the surrounding material. We note that this implies enhanced convective motions and enhanced mechanical energy transfer to the overlying material. In particular, the fluctuating top of the convective zone may approach the top of the photosphere and may even sometimes break through it.

The research in this paper has been supported in part by the Geophysics Research Directorate, Air Force Cambridge Research Center, Cambridge, Mass., under contract with Harvard University.

*Harvard College Observatory,
Cambridge, Mass.*

Kupperian, J. E., Jr. and Milligan, J. E. Sources and stellar fluxes in the far ultraviolet.

Identifications of the discrete sources observed by E. T. Byram *et al.* (1956) in the far ultraviolet have been made. The strongest are γ Velorum, ζ Puppis and Regulus. There also appear