The method of analysis is that of weighting functions, as developed by Minnaert and Claas for weak lines and by Pecker for lines of medium strength. In this method, the equivalent width of each line is related to the abundance by an integral over the model atmosphere, in which the integrand is the product of the number of absorbing atoms, the weighting function at each point in the atmosphere, and a so-called saturation function which reduces to unity for weak lines. Special numerical techniques for the evaluation of these integrals have been devised by Elste and adapted by him for computation with the Michigan high-speed computers, MIDAC and the IBM 650. An extensive set of these integrals has been tabulated by Elste for both weak lines and those of medium strength covering the wave-length region $\lambda\lambda 4000$ to 22,000 and a wide range of excitation potentials.

This tabulation has provided the basis for the present investigation of solar abundances. The model atmosphere employed was derived by Waddell and Pierce from the latter's recent observations of solar-limb darkening and spectral-energy distribution, but with the pressure distribution modified by Elste in accordance with the work of Weidemann.

The results are generally in good agreement with those of other workers, but there are notable exceptions, especially for those abundances that depend on the analysis of lines of medium strength. The reason appears to be that the shape of the curve of growth in the transition region differs markedly for lines originating at different depths in the atmosphere. It is anticipated that the results quoted here can be further improved when new photoelectric observations of weak lines are secured with the vacuum spectrograph of the McMath-Hulbert Observatory. However, for a large number of elements, uncertainties in the f -values present the major obstacle to definitive abundance determinations.

*University of Michigan,
Ann Arbor, Mich.*

Greenstein, Jesse L. Some high-velocity Population I stars.

A group of objects high above the galactic plane has been found in the investigation of faint blue stars. Some of these show such high radial velocities, that in spite of lack of knowledge of the transverse motions, we can be sure that the space motions are now large, and must have been

even larger, 200–300 km/sec, when the stars passed through the galactic plane.

In spite of the large motions, some of these objects clearly belong to groups associated with Population I, and it is proposed to call them "runaway" Population I. Included in the group are the following: (1) HD 172324, a supergiant of type A0 Iab, (2) A symbiotic Op and gM star, Mt. Wilson catalogue 603, with strong emission lines both permitted and forbidden, (3) Two distant B stars at intermediate galactic latitudes, (4) A spectroscopic binary of type B2, period 4 days and semi-amplitude 125 km/sec, located about 6000 pc above the galactic plane. (Collaborators in the various investigations listed above are (1) W. Bonsack (2) W. Tifft (3) D. MacRae and R. Fleischer.)

These objects show that some stars have been produced, very recently, from very high-velocity interstellar clouds, or in extraordinarily unstable associations.

*Mount Wilson and Palomar Observatories,
Carnegie Institution of Washington,
California Institute of Technology,
Pasadena, Calif.*

Grenchik, R. A model atmosphere for the white dwarf 40 Eridani.

A model atmosphere is calculated for the white dwarf 40 Eridani B using the parameters given by Schatzman (1949), effective temperature 13,800°K and surface gravity $\log g = 7.69$. A pure hydrogen atmosphere in radiative and hydrostatic equilibrium is assumed.

Various processes are considered as sources of continuous absorption including those due to negative hydrogen ion, neutral hydrogen and the overlapping of line wings. The contribution due to negative hydrogen ion can be neglected. The main contribution to the continuous absorption comes from the neutral hydrogen. The overlapping of line wings may be serious in the Lyman region. To set a range of applicability of the model-atmosphere calculations two models are constructed: Model 1, ignoring the absorption in the wings of the Lyman lines; and Model 2, including an approximation for this effect. The Lyman contribution adds about 0.2 to the $\log \kappa$. Chandrasekhar's (1950) mean absorption coefficients are used. The modified (2,1) approximation is used for the temperature distribution in the atmosphere and the hydrostatic equilibrium equation is solved numerically by the modified Euler method.