Gehrels, Thomas. *Polarization of the light from Moon, Mars, Venus, and NGC 7023.*

With the Wollaston photometer, described separately, the polarization of seven small lunar regions was observed at different phases and wave lengths. The results in visual light are similar to those obtained by Lyot (1929), but the wave length dependence is strong. Up to 23 per cent polarization was found in the ultraviolet for lunar maria at quarter phase.

Mars and Venus show the same wave-length dependence with strong polarization in the ultraviolet. In the ultraviolet, Mars at 33° phase has 6.7 per cent, while Venus at 59° phase has 2.4 per cent positive polarization. In visual light, the agreement with Lyot's results is nearly perfect, with Venus at 59° phase showing 1.7 per cent negative polarization.

The present program was initiated in 1956 to verify, or disprove, an explanation of lunar phase effects in terms of accreted interplanetary particles. The strong wave-length dependence of Moon, Mars, and Venus, and the peculiar polarization-reversal of Venus may indeed be explained by a partial coverage with very small accreted particles. The observed polarization of Mars and Venus is then a composite that is partly caused by suspended particles as well as by the planetary atmosphere.

Three regions of NGC 7023 were observed with wide filters at 3600, 5500, and 8300 Å; a typical region has 13, 19, and 22 per cent polarization with these filters, respectively. Particle sizes of the order of 0.5 microns, different from those for Moon, Mars, and Venus, are indicated for NGC 7023.

REFERENCE

Lyot, B. 1929, *Ann. Meudon* 8, 42.

*Yerkes and McDonald Observatories
University of Chicago
Chicago, Ill.*

Gehrels, Thomas and Teska, Thomas M. A *Wollaston photometer.*

A photometer of the McDonald Observatory was converted for interchangeable use in either *UBV* photometry, or in six-color polarization measures. The polarization is analyzed with a Wollaston prism and two multiplier phototubes by simultaneous integration. A complete series on a fairly bright object, at six position angles and in six colors, takes about 60 minutes; this requires 144 integrations as at each angle the

tubes are exposed twice, namely with and without a depolarizer in the incoming beam. Neither thin clouds nor poor seeing affect the polarization measures. The effective wave lengths range from 3250 to 9900 Å. Two types of tubes are used, namely RCA 7102, having S-1 response, and RCA 7265, with multi-alkali cathode. Reductions of the observations are made with the IBM 650 of Indiana University. The internal probable errors are $\pm 2^\circ$ in position angle, larger when the polarization is very small, and ± 0.1 in the percentage polarization.

Results on the main program of solar system studies are in a separate paper. As a check on this new equipment, eight strongly polarized stars were observed that occur in the lists of Hall and Mikesell and of W. A. Hiltner; in visual light the agreement is close. However, it was found that in the ultraviolet as well as in the infrared the interstellar polarization is less than in visual light. The sunlit blue sky shows approximately the same wavelength dependence. It was found that neither the 82-inch mirror nor the photometer cause depolarization in the ultraviolet; the observations and the depolarization test will be fully reported in another publication.

*Astronomy Department
Indiana University
Bloomington, Ind.*

Giordmaine, J. A., Alsop, L. E., Townes, C. H. and Mayer C. H. *Observations of Jupiter and Mars at 3-cm wave length.*

Radiation associated with the planet Jupiter in the wave length range 3.03 to 3.37 cm has been observed during the period April 16, 1958 to February 7, 1959. The observations were made with the U. S. Naval Research Laboratory 50-foot reflector in conjunction with a maser amplifier (Giordmaine *et al.* 1959). A total of 153 drift curves through the position of the planet was obtained.

The following are the average equivalent black body temperatures of the visible disc observed in the several wave length ranges: 3.36 ± 0.01 cm, April 16 to May 8, 1958 with the exception of April 30 to May 1, $189 \pm 20^\circ\text{K}$; 3.17 ± 0.02 cm, May 24 to July 29, 1958 and January 31 to February 7, 1959, $173 \pm 20^\circ\text{K}$; 3.03 cm, August 22 to September 4, 1958, $171 \pm 20^\circ\text{K}$. The probable errors quoted for the black body temperatures are mainly those arising from the uncertainty in the effective