2. Procurement of improved telescope mountings, with circles capable of giving position readings to an accuracy of at least $\frac{1}{10}$ degree of arc.

3. Servicing of stations on a wider and more frequent basis.

4. Development of techniques and instrumentation for producing greater accuracies in position and time measures.

5. Establishing a satellite photometric (light-curve) project.

Observations of a Large Redshift. WILLIAM A. BAUM AND R. MINKOWSKI, *Mount Wilson and Palomar Observatories, Carnegie Institution of Washington, California Institute of Technology, Pasadena, California.*—Investigation of radio sources has led Minkowski to the discovery of a very distant cluster of galaxies whose brightest member provides the identification for the radio source 3C295. The spectrum of this galaxy shows a strong emission line of $\lambda 5447.8$. By analogy with the spectrum of Cygnus A the line can only be identified with $[OII] \lambda 3726/29$. The redshift therefore is 0.4614 ± 0.0002 .

Redshifts above 0.2 can be observed spectroscopically only for objects showing strong emission lines, but for all galaxies of normal color they are readily accessible to the method of multicolor photoelectric photometry. Using this method Baum has observed a redshift of 0.44 ± 0.03 for two average galaxies in the central part of the cluster. The agreement with the spectroscopic value proves the validity of the photometric method for the measurement of redshifts of up to 0.5.

The present results establish the possibility of reaching redshifts at which cosmological effects are large enough to permit observational discrimination between different cosmological theories. The results, in line with two earlier photometric determinations of redshifts in the range from 0.3 to 0.4 by Baum, favor an exploding universe, but observations of more very distant clusters are needed to determine the relation between redshift and magnitude with adequate accuracy.

Major Flares and Geomagnetic Activity. BARBARA BELL, *Harvard College Observatory.*—The relation between geomagnetic storms and major solar flares was studied, with particular attention to the magnetic type and the location of the flaring sunspot. For the 23 years, 1937–1959, the IAU Quarterly Bulletins yielded 580 major (importance $\geq 2^+$) flares. Daily magnetic classifications were obtained for the associated sunspot groups from Mount Wilson records.

Each flare was assigned to one of four geomagnetic categories, according to the behavior of the Kp and Ap indices in the three days following the flare:

no storm, small storm, moderate storm, and great storm (an Ap ≥ 100 and or one 3-hr Kp $\geq 9^-$).

A flare occurring in a magnetically complex sunspot group (γ or $\beta\gamma$) is more likely to be followed by a major storm than is a flare in a unipolar (α) or bipolar (β) spot group. The percentage of major flares followed by a great storm for each type of spot is: (γ) 23%, ($\beta\gamma$) 23%, (β) 4.6%, (α) 11%, with an average of 12.6% for all 580 flares. The percents followed by a moderate or a great storm are: (γ) 42%, ($\beta\gamma$) 35%, (β) 16%, (α) 21%, with an average of 25% for all types.

The expected concentration of great-storm flares toward the central regions of the solar disk was found, but this effect was less striking than the concentration of great-storm flares in the northern solar hemisphere. In the 23 years studied, 62% of the major flares occurred in the solar north. Northern spot groups produced 86% of the major flares followed by great storms and 64% of those followed by moderate storms. The probability of a great storm after a northern flare is 17.6%, contrasted with only 4.5% after a southern flare. This surprising effect appears about equally in each of the three sunspot maxima studied, and tends to increase with the intensity of the storm.

The work here reported was supported by ARDC contract. The cooperation of Mount Wilson Observatory in making sunspot classification records available, and of the Meudon Observatory in providing a prepublication copy of the 1958–1959 flare lists, is gratefully acknowledged.

Absolute f -Value of the PbI Line, $\lambda 2833$. GRAYDON D. BELL, *Harvey Mudd College*, AND ROBERT B. KING, *California Institute of Technology.*—Improvements made in the atomic beam apparatus used for the measurement of absolute f -values of resonance lines of the metals and refinements in techniques are described. Recent measurements of the PbI line $\lambda 2833$ give an absolute f -value of 0.23 ± 0.02 for this line. This is in poor agreement with the f -value of 0.60 reported by H. D. Engler (1956, *Z. Physik* 144, 343). Our f -value, in combination with the measurements of the relative f -values of $\lambda 2833$, 3683, and 4057 by M. Z. Kokhlov (1959, *Contrib. Crimean Astrophys. Obs.* 21, 84), yields absolute f -values of 0.16 for $\lambda 3683$ and 0.35 for $\lambda 4057$.

Supported by the Office of Naval Research.

The Brighter Early-Type Stars in the Region of the North Galactic Pole. WILLIAM P. BIDELEMAN, *Lick Observatory*, AND BENGT WESTERLUND, *Uppsala Southern Station, Mount Stromlo, Australia.*—Spectroscopic and photometric observations, made at the McDonald and Lick Observatories, respectively, are reported for approximately 100 early-