the conclusion follows that only the modified gravity waves may be present.

Radiative and Conductive Opacities for Stellar Mixtures.

A. N. COX AND D. D. EILERS, *Los Alamos Scientific Laboratory*.—A program for the calculation of radiative and conductive opacities is in progress at the Los Alamos Scientific Laboratory. This work is appropriate because of new hydrogen-like bound-free and free-free gaunt factors of Karzas and Latter and because accurate opacities are needed for modern stellar models. Checks are being made of all the opacities given by Keller and Meyerott and by Zirin. Some of the opacities for this program are being calculated by considering the electronic occupation of individual ions instead of the usual methods which consider only the average atom of each element. Comparisons show that the previous radiative opacities are generally accurate to about 10% except in isolated cases. Conductive opacities are calculated by a method slightly modified from that described by Mestel. Within 10% the results for the Russell mixture agree with conductive opacities given by Lee. Errors in the values for radiative and the combined radiative and conductive opacities should be less than 10% if bound-bound absorptions are negligible as assumed.

Programs for Automatic Reduction of UBV and H_β Photoelectric Measures.

D. L. CRAWFORD, *Dyer Observatory, Vanderbilt University*.—Programs have been written for automatic reduction of photoelectric observations, utilizing an IBM 650 computer system. Three separate programs are available. The first is a general one for reduction of UBV observations. The second is a program especially written to determine extinction coefficients, and to keep close check on any variation of these coefficients. The third is a general reduction program for H_β narrow band photometry. An attempt has been made to keep all three programs as general and adaptable as possible, and all are optimized for efficient use of machine time. Some special features and techniques are pointed out. All three programs are available from the Vanderbilt University Computing Center at cost.

Color Classification of Galaxies.

G. DE VAUCOULEURS, *Harvard College Observatory*.—Colors of about 150 galaxies in the UBV system are used to define a standard "main sequence" in the $U-B$, $B-V$ diagram. Departures from the main sequence arise from ultraviolet excess in some peculiar galaxies (e.g., Mayall's nebula) and in the emission nuclei of some spirals (e.g., Seyfert's galaxies). Many of the other "blue" galaxies described by Haro are merely late-type irregulars which do not depart significantly from the main sequence. The color distribution in normal galaxies is analyzed as a function of tilt, galaxy type, and absolute magnitude. After correction for reddening caused by tilt effects the colors of spirals and irregulars are closely correlated with morphological type (Sa to Im) and with the ratio $A/D(O)$ of the field aperture to the reduced or "face on" diameter $D(O)$. This correlation can be used for an objective quantitative color classification of galaxies in good agreement with spectroscopic criteria. For ellipticals (E) and lenticulars (SO) there is a large variation of color with absolute magnitude, which can be used for distance determinations after correction for red-shift effects. Residual color excesses in low galactic latitudes permit a determination of interstellar absorption for individual objects of well-defined morphological type.

Flares of July 16, 1959.

H. W. DODSON AND E. R. HEDEMAN, *McMath-Hulbert Observatory*.—Two flares of importance 3 and 3⁺, respectively, occurred on July 16, 1959 in the north central part of the solar disk. They were separated in time by about five hours. The two flares differed greatly in their radio frequency emission. The earlier flare was accompanied primarily by a type II or slow-drift burst. The second was one of the relatively few flares known to be associated in time with great emission at centimeter wavelengths, with great con-

tinuum radiation at radio frequencies, and with enhancement of high energy particles in the neighborhood of the earth. For these two flares, the time intervals between start and maximum were similar, being 23 and 21 minutes, respectively. Although these values are longer than average for flares in general, they are not abnormally long for flares of importance 3. During the second of the two flares, spectroheliograms were made by systematic changes of wavelength throughout 6A of the spectrum centered on $H\alpha$. These spectroheliograms revealed numerous hydrogen absorption features, some with ascending and some with descending motions. In the late stages of the flare, spectroheliograms taken to the redward of $H\alpha$, showed an extensive system of curved dark features which may be loop-type prominences in projection against the disk. Comparison revealed a conspicuous difference in the *locations* of the two flares. Portions of the flare with the great centimeter and continuum radiation occurred *directly over* the umbrae of large spots. No portion of the earlier flare, which did not have this type of radio-frequency emission, appeared over the umbra of a large spot. Similar comparisons have been made for a small number of other flares. The association between very great centimeter wavelength radio emission and close proximity of $H\alpha$ flare and large spot umbra is sufficiently good to warrant further investigation.

Survey of Number of Flares Observed during the IGY.

H. W. DODSON AND E. R. HEDEMAN, *McMath-Hulbert Observatory*.—In the McMath-Hulbert Observatory working list of world-wide flares there are more than 6700 solar events during the IGY for which at least one observer estimated the importance to be ≥ 1 . According to reports to World Data Centers the flare patrol was gratifyingly complete and uniform. However, examination of the starting times of the reported flares reveals evidence for a serious lack of homogeneity in the IGY flare data. Approximately twice as many flares, ≥ 1 , were reported per hour from 05^h to 16^h UT as during the remainder of the Universal Day. This strong dependence on UT exists in the flare data for each month of the IGY. Examination of the patrol hours and flare reports for May and July 1958 for each of the 19 $H\alpha$ cinematographic patrol stations shows that the principal photographic patrol stations fail to confirm the very high rate of flare occurrence for 05^h to 16^h UT that appears in the total IGY data. Furthermore, for the hours 17 to 04 in the Universal Day the IGY flare reports are based primarily on the cinematographic patrol. Comparison of the number of flares and subflares indicates that in the IGY data for the hours 05 to 16 UT many flare events are listed as flares that probably would have been classified as subflares by the principal cinematographic observers.

Remarks on the Radio Emission from the Gamma Cygni Nebulosity.

F. D. DRAKE, *National Radio Astronomy Observatory*.—The existing radio data suggest that the radio emission from the Gamma Cygni nebulosity has the spectrum of an HII region. It is shown, using Stromgren's original relations, that the radio flux predicted for the Stromgren sphere of an F8 Ib star is adequate to explain the observed radio flux from Gamma Cygni. Thus the strong radio emission and bright nebulosity are not unusual, despite the late spectral type. It appears that the existence of a bright nebula close to Gamma Cygni is a result of the star's peculiar space motion, which has carried the star into a dense dark nebula. The resulting ionization has exposed portions of the dust cloud to quantitative study. Densities greater than 1000 atoms/cc probably are present in the cloud.

Detection of Discrete Solar Radio Emission Sources during the October 2, 1959 Partial Eclipse.

S. EDELSON, C. R. GRANT, AND H. CORBETT, *Radio Astronomy Branch, U. S. Naval Research Laboratory*.—On October 2, 1959, observations of the partial solar eclipse were made at the Naval Research Laboratory. Variations in the solar flux were recorded at 0.43-, 3.15-, and 9.4-cm wavelengths as the sun was gradually uncovered by the moon. H-alpha photographs of the