is determined by the shock strength and can exceed the piston speed by a factor of 2 or more. A change of sign of the derivative of velocity with depth after a quarter-cycle is due to the outward-propagating rarefaction wave.

Absorption contours were computed at $\mu=1$ for lines of two strengths— $\log \eta_0=0$ and $\log \eta_0=1.6$ —at different phases. These show a progressive line doubling as the wave front advances; intensity ratios for the two components of each line may be quite different, and even reversed, between the weak line and the saturated line.

Calculations with similar models in which the line contours were integrated over the disk have shown (1) that weak shocks may be present without causing line doubling and (2) that the velocity differences between components are not closely related to shock strength, but depend upon the sign and magnitude of $(d/d\tau)|v|$.

The results of these calculations also provide a foundation for an observational distinction between the view that solar granules are produced by compression waves generated in the convection zone and the view that the granules are themselves the convective cells, through analysis of high-dispersion spectra procured with the vacuum spectrograph at the McMath-Hulbert Observatory.

Lunar Temperature Measurements at 8.6-mm Wavelength. WARREN C. TYLER AND JACK COPELAND, *Research Laboratory, Army Rocket and Guided Missile Agency, Redstone Arsenal, Alabama*.—Radio observations of the moon have been made at 8.6-mm wavelength using a conventional Dicke-type superheterodyne radiometer. The antenna is a horn-fed paraboloidal reflector five feet in diameter. The beam width is about 20 minutes of arc to half-power points.

Effective lunar disk temperatures were measured during the eclipse of September 5, 1960. At this longitude, moonset occurred at the beginning of totality, by which time a net decrease of about 20°K was observed.

Observations have also been made in order to determine the variation in effective disk temperatures as a function of lunar phase. The results obtained from these measurements are in substantial agreement with previous measurements at this wavelength.

The Clustering of Galaxies. SIDNEY VAN DEN BERGH, *David Dunlap Observatory, Ontario, Canada*.—A cluster of galaxies is usually regarded as a distinct type of aggregate of matter which is intermediate between the individual galaxy and the universe as a whole. The concept "a cluster of galaxies" would

lose much of its usefulness if it could be shown that most clusters can be divided into subclusters or even smaller units. Alternately, one may inquire if clusters of galaxies are the largest aggregates of matter or whether they are themselves constituents of still larger organizations.

Examination of the Local Group and of the Virgo cluster shows that subclusters exist within each of these systems. An analysis of the angular separations of the galaxies in the Coma cluster indicates a weak subclustering tendency. In fact, subclustering appears to be a quite general phenomenon, although it may be less pronounced in dense clusters like the Coma cluster than in low-density clusters like the Local Group. The existence of binary galaxies (LMC, SMC) and multiple galaxies (M31, M32, NGC 205) within the local group is well established. An analysis of the radial velocities of galaxies in the Virgo cluster also indicates the existence of stable binary systems.

The Lick counts of galaxies give strong evidence for the existence of clusters of clusters of galaxies with characteristic dimensions of the order of 10 mpc. There is also fairly good evidence that the "Local Group" belongs to a vast supercluster containing the Virgo cluster. Do still larger aggregates of matter exist? An analysis of the distribution of the most distant ($V \approx 50\,000$ km/sec) very rich clusters observed by Abell indicates that inhomogeneities with characteristic dimensions in excess of 100 mpc may exist.

The available evidence strongly suggests a hierarchy of clustering ranging from binary galaxies to clouds of superclusters of galaxies. The universe may well be much less homogeneous than has been generally assumed.

The Rayleigh-Taylor Instability in Compressible Fluids. PETER O. VANDERVOORT, *National Radio Astronomy Observatory, Green Bank, West Virginia* (now at *Princeton University Observatory*).—The Rayleigh-Taylor instability of an isothermal atmosphere supported against gravity by a uniform incompressible fluid has been analyzed. A perturbation characterized by a periodic deformation of the boundary separating the two fluids is found to grow like $\exp \eta$, and the dependence of the growth rate η on the wavelength λ of the perturbation is determined. For very short wavelengths, η is proportional to $\lambda^{-1/2}$ (as is the case for all wavelengths when both fluids are incompressible); whereas for long wavelengths η is proportional to λ^{-1} . Thus, the effect of compressibility is to retard the growth rates for the longest wavelengths. The transition from the $\lambda^{-1/2}$ -dependence to the λ^{-1} -dependence of η occurs in a range of wavelengths of order λ_0 , where λ_0 is a determinate quantity in the theory.