Interferometric Stellar Spectrometry. LAWRENCE MERTZ, *Block Associates, Inc., Cambridge, Massachusetts* (Introduced by S. Goldstein).—A brief review of the properties of interferometric spectroscopy for astronomical applications is presented along with recent results.

(1) An interferometer and a conventional scanning monochromator give equal average signal to photon noise ratios, assuming the same monochromatic light transmission.

(2) An interferometer, however, can transmit much more light than a similar size instrument requiring slits, since the interferometer does not require collimated light. This advantage becomes even more important when one is obliged to compromise resolution with scintillation noise and energy loss both on the slit jaws.

(3) Dark current noise for an interferometer becomes distributed among all the measured spectral elements resulting in an effective reduction by the square root of the number of resolved elements. (Although the photon noise is also distributed this reduction is exactly compensated because, looking at more colors simultaneously, the interferometer suffers more photon noise.) This renders photomultiplier cooling unnecessary and thus permits better red sensitivity.

(4) Computation of a Fourier transform is required. This property has so far discouraged widespread use of interferometric spectrometry.

In order to demonstrate that interferometric spectrometry actually works a polarization type interferometer was constructed and used for stellar spectrometry. The instrument transmits an $f/15$ bundle through a 5-mm-diameter aperture, automatically subtracts sky background, resolves about 60 Å, and weighs only 7 lb.

Interferograms of a variety of sources (Orion Nebula, Mira, Gamma Geminorum) were made on two nights with the interferometer installed on the 60" telescope at Mt. Wilson and the spectra computed from these data were shown.

Dynamics of Spherical Stellar Systems: Properties of Theoretical Models, and Comparison with Clusters and Elliptical Galaxies. RICHARD W. MICHIE, *University of California, Berkeley*.—A wide range of spherical models are now available and result from a distribution function which for the inner regions is obtained from the Boltzmann equation with encounters. It is assumed that encounters and orbital mixing were important in the early stages of the formation and development of the star clusters and elliptical galaxies, and, since this time period, the structure of these systems has not significantly changed. The orbital mixing allows an extension of

the distribution function to the regions far from the center, where mathematically it satisfies the collisionless Boltzmann equation. The gravitational potential is obtained from Poisson's equation. The stellar orbits are not restricted to an isotropic velocity distribution everywhere—the models range from complete isotropy to highly radial orbits. While the distribution function is obtained for stars of different mass, the comparison with observation has assumed all stars to have the average stellar mass.

In a log density versus log distance plot, it is found that for certain degrees of velocity space anisotropy there is an inflection in the curve, and this is a result of the orbits becoming elongated as well as the energy cutoff becoming important. All the models have finite mass, are self-supporting and satisfy the virial theorem. They are all obtained from an analysis of the Boltzmann equation with particular emphasis placed at the high energies. In dimensionless variables the models are characterized by a single model parameter C . The model parameter involves three physical parameters which indicate the total mass, total energy, and degree of velocity space anisotropy.

A comparison with rich galactic clusters, globular clusters, dwarf galaxies and spherical galaxies has been made. The theoretical models appear to be in agreement with the observed systems. For example, the log density versus log distance inflection observed in systems such as NGC 3379 is exactly as predicted by the theory. There may be a small inflection for the globular cluster 47 Tuc. Both star clusters and spherical galaxies show a fairly wide range in the model parameter. The star clusters may favor a somewhat larger value of C , but the data are probably not sufficient to firmly establish this trend.

Luminosity of Galaxies in Clusters and in the Field. J. NEYMAN AND E. L. SCOTT, *University of California, Berkeley*.—Earlier study, based on 342 field and 104 small group galaxies in HMS catalogue (Humason, Mayall, Sandage, *Astron. J.* **61**, 97, 1956), provided estimates (see abstract by Marcus) of luminosity functions and selection probabilities for eight different morphological types. These are used for comparison with cluster galaxies.

System	No. of galaxies in HMS	Mean radial velocity	m_0	m^*	m
Near groups	32	629	11.4	10.7	10.7
Virgo	80	1197	12.9	11.6	11.4
Intermediate groups	72	2961	14.1	12.5	12.6
Coma	46	6866	16.7	13.9	14.8
"Far" clusters	48	11696	17.4	14.5	16.0