Santiago, Chile.—The sporadic radio noise emitted by Jupiter was monitored in 1961 at the University of Florida Radio observatory, near Gainesville, at the Fleurs Field Station of the Radiophysics Laboratory near Sydney, Australia, and at the Maipu Radioastronomical Observatory, near Santiago, Chile. The observations were made at 5, 10, 15, 18, 19.7, 27.6, and 85.5 Mc/sec. There was no observable activity at 85.5 Mc/sec. The observations at this frequency were carried out by O. B. Slee, using a section of the original Mills' cross antenna. The 5 Mc/sec observations, which were made in Chile, were sparse because of excessive interference. From 27.6 Mc/sec down to 10 Mc/sec, there was a monotonic increase in the percentage of nights on which activity occurred. Peak flux densities increased monotonically from about 2×10^{-21} W m⁻² cps⁻¹ at 27.6 Mc/sec to about 10^{-18} W m⁻² cps⁻¹ at 5 Mc/sec. An analysis of the distribution of pulse flux densities was made at each of 4 frequencies from 10 to 27.6 Mc/sec, from which the time average flux density was determined for each frequency. The average flux density increases monotonically with a decrease in frequency within this range, as does the flux density value which is exceeded any fixed percentage of the time. Further observations will be conducted to determine whether there is an asymptotic decrease or a sharp cutoff in activity between 27.6 and 85.5 Mc/sec.

The National Science Foundation, the U. S. Army Research Office (Durham), and the Office of Naval Research participated in the financial support of this project.

Planetary Distances According to General Relativity. G. M. CLEMENCE, *U. S. Naval Observatory*.—Radar echoes from Venus have been interpreted (*Astron. J.* 67, 181, 191, 1962) to mean that the distance from the earth to the sun is about 75 000 km greater than is indicated by the dynamical method (*Astron. J.* 59, 409, 1954). A search for relativistic effects previously neglected yields corrections to the radius vector of a planet amounting at most to about a kilometer. The cause for the discrepancy must be sought elsewhere.

Emission from a Rapidly Rotating Stellar Atmosphere. GEORGE W. COLLINS, II, *Washburn Observatory*.—The primary aim here is to describe the continuum emission from a rotating atmosphere. It

is assumed that the dynamical structure of the rotating star is that of a Roche model, in rigid rotation. The further assumption is made that the atmosphere at any point on the surface of such a star can be approximated by the atmosphere for a spherical star with the temperature and gravity of the point in question.

Calculations were carried out for the gray atmosphere approximation, and monochromatic fluxes and magnitudes were determined for various rotational velocities and angles of inclination for a specific model B3V star. Additional calculations were made to determine the variation of the Balmer jump and Lyman jump with angle of inclination for the model rotating with breakup velocity.

It was found that large variations in the observed total luminosity may result from rotation. Significant variation in the colors in the ultraviolet may also result. However, variation of the Balmer and Lyman jumps as well as the visual and blue colors remain essentially unchanged, regardless of the rotational velocity and angle of inclination.

Some Singular Orbits of an Earth-Moon Satellite with a High Area-Mass Ratio. G. COLOMBO AND D. A. LAUTMAN, *Smithsonian Astrophysical Observatory and Harvard College Observatory*.—No gravitational orbit will, in a few revolutions around the earth, lead from a perigee close to the earth to a capture by the moon. Within the framework of the restricted three-body problem, Egorov (*Russian Literature of Satellites*, New York, 1958) estimates that about 200 revolutions are needed before a particle may pass through the constriction in the Hill curve at point L_1 and become, at least temporarily, a satellite of the moon. It had therefore been thought that to effect a lunar capture would require a second active rocket phase close to the moon.

It is possible, however, considerably to speed up the capture process by utilization of the radiation pressure of the sun on a satellite with a high area-mass ratio. The basic mechanism consists of a secular increase of the angular momentum of an orbit initially highly eccentric with respect to the earth until the orbit becomes nearly circular and the particle passes through L_1 with a small velocity with respect to the moon.

For satellites with an area-mass ratio of 280 cm²/g, we have calculated several orbits that lead to captures by the moon 50 to 75 days after launch. These times correspond to 4.5 to 6.5 revolutions around the earth. The periods of the selenocentric orbits are of the order of 4.5 days.

Comparative Spectrophotometry of Selected Areas on the Lunar Surface. GEORGE V. COYNE, S.J., *Georgetown College Observatory*.—During the months September 1961 to January 1962, a Wadsworth spectrograph, giving a reciprocal dispersion of 4.9 Å/mm on the normal to the grating in the first-order spectrum, was used to obtain spectrograms of selected areas on the lunar surface. Identification of the observed areas was accomplished by photographing the slit-plane image of the moon at the time the spectrograms were taken. The spectrograms were measured on a recording microphotometer in a direction transverse to the dispersion at approximately every 100 Å in the spectral range 4380 to 6835 Å with a bandpass of 9.3 Å. These measurements provide a densitometric profile across the observed lunar regions in which various small areas can be identified. From these profiles comparative spectral energy distribution curves of one area with respect to another are constructed. The results are given in the form of a color-contrast function defined as follows:

$$CE_{a,b}(\lambda) = (m_{\lambda} - m_{5500})_a - (m_{\lambda} - m_{5500})_b,$$

where the subscripts a and b refer to different areas of the lunar surface and m_{λ} , m_{5500} are the magnitudes for the wavelengths λ and 5500 Å, respectively.

In general, the curves show a smooth but non-monotonic variation with wavelength. The maximum values of $CE(\lambda)$ lie between 0^m15 and 0^m20. The lunar maria are more homogeneous in blue light (below 5500 Å) than in red light (above 5500 Å). On several curves, and notably for lunar areas near the edges of maria, there are humps resembling luminescence bands observed by Dubois (*Publ. l'Obs. l'Univ. Bordeaux* Ser. A, No. 13, 1959) for various areas of the lunar surface. These bands are centered at approximately 5050, 5200, and 6100 Å. They are about 500 Å wide and have an amplitude varying from 0^m03 to 0^m18 above the smooth curve on which they are superimposed.

Variable Star Search in M53. JAMES CUFFEY, *Goethe Link Observatory*.—In order to provide a complete survey of variables in the globular cluster M53, including stars of small amplitude, all the resolved images on the 100-inch reflector plates have been measured with an iris photometer.

The expected increase in sensitivity over the blink-microscope technique has been realized, and in addition to rediscovery of the known variables, 4 new variables were found, plus 12 probable variables, and 30 suspects. Most of the suspects have small amplitudes approaching the limits of detectability set by the probable error, ± 0.03 mag., of measurement.

In the discovery survey, 620 images were measured on 30 plates; the stars showing the highest residuals, regarded as possible new variables, were then measured on 40 additional plates. Analysis of the residuals related to apparent magnitude reveals an interesting tendency for stars in the immediate vicinity of the absolute magnitude of the cluster variables to show less variability than stars a magnitude brighter and a magnitude fainter. Further measures for light curves are in progress.

Emission of Hydromagnetic Waves from Sunspot Umbras. ROBERT E. DANIELSON, *Princeton University Observatory*.—Linearized equations describing convection in the presence of a magnetic field predict overstable oscillations in sunspot umbras. No such oscillations have been observed, however. The equations which predict these oscillations have boundary conditions which allow only standing wave oscillations. It is suggested that these are not the correct boundary conditions to use. More physically realistic boundary conditions would allow the convective layer to emit hydromagnetic waves.

The critical equivalent Rayleigh number for the emission of hydromagnetic waves has been computed for various magnetic field strengths using these new boundary conditions. The equivalent Rayleigh number in sunspot umbras seems to exceed the critical equivalent Rayleigh number. Thus it appears possible that a significant nonthermal flux of hydromagnetic waves is emitted by sunspot umbras. Such a flux is consistent with the fact that higher velocities are observed in sunspots than in the normal photosphere. It might also account for the missing energy in sunspots.

This work is part of Project Stratoscope, which is sponsored by ONR, NSF, and NASA.

Statistical Analysis of Jupiter's Decameter Radiation 1950–1961. JAMES N. DOUGLAS, *Yale University Observatory*.—Observations of Jupiter's decameter noise storms furnished by all observers in the period 1950–1959, by Yale and High Altitude Observatory in 1960, and Yale in 1961 have been combined to form an observation time series for statistical analysis. No evidence for a change in the previously announced average period ($9^h55^m29^s37^{\pm}0^s13$) is found.

Slow variations in this period totaling 0^s4 could have been detected but were not. Autocorrelation studies provide a measure of the average intrinsic duration of a Jupiter noise storm, as well as demonstrating the nonindependence of the three significant regions of activity. Cross-correlation studies with various solar and geomagnetic indices, together with significance tests, are described.

This work was supported by the National Science Foundation.

Novel Interferometer for Studies of Discrete Radio Sources. JAMES N. DOUGLAS AND CLINTON C. BROOKS, *Yale University Observatory*.—A 250-Mc interferometer nearing completion at the Bethany Observing Station of the Yale University Observatory will measure the declination of 500 radio sources to an accuracy of 20". Obtaining its *N-S* directivity by virtue of a broad operating bandwidth, the instrument will utilize 10 simultaneous beams to compress observing time and to increase effective sensitivity and degree of redundancy. The potentialities of this technique as applied to much larger instruments are described.

This work has been supported by the National Science Foundation.

Performance of Multistage Image Amplifying Tubes; Astronomical Applications; and Proposed Use with Fabry-Perot Etalons for Stellar Spectroscopy. T. DUNHAM, JR., *Mount Stromlo Observatory*.—Three properties of multistage image tubes (Wilcox, W. L., Emberson, D. L., and Weekly, B., *Nature* **185**, 370, 1960) offer advantages for photometry at low flux levels: (1) high-speed recording; (2) background electrons are focused at the output, thereby reducing the number per resolution band to approximately 10^{-4} of the contribution encountered with photomultipliers; and (3) scintillations, corresponding to individual photoelectrons, are bright enough to permit recording a considerable fraction on moving film. Information storage is greater on moving film and scintillations in individual spectral resolution bands can be counted, so that the tubes can be used as multichannel pulse counters [Dunham, T., Jr., *J. Opt. Soc. Am.* **50**, 1129(A), 1960].

The factor of merit of an image tube for spectrophotometry may be expressed as G^2r^2/N , where G is the over-all light gain, r the linear resolution, and N the background.

For direct stellar photometry, an image tube offers the possibility of recording flux from a number of star images (and sky) simultaneously, using fiber light guides to transfer images to a linear array on the photocathode, so that scintillations can be recorded and counted on moving film.

Speed, resolution, and background have been investigated, using test spectra and coude spectra with Baum's microprojector (Baum, W. A., *Advances in Electronics and Electron. Phys.* **16**, 391, 1962). The percentage of incident photons that can be recorded on film has been determined by using a low-intensity

scattering source of approximately monochromatic light.

For high-resolution spectroscopy, the use of an image tube and moving film, with a scanning Fabry-Perot etalon (Jacquinot, P., and Dufour, C., *J. Rech., Cent. Nat. Rech. Sci.* **2**, 19, 1948) and rotating grating, appears to offer an efficient system that permits pulse counting of many fringes simultaneously. For space spectroscopy, direct electrical pickup at the output, on narrow conducting strips corresponding to the fringes, and connected to individual pulse-counting circuits, would have advantages.

The experimental part of this study was carried out at Imperial College, thanks to the kindness of Dr. J. D. McGee.

High-Luminosity, Fully Convective Contracting Sun. I. D. EZER AND A. G. W. CAMERON, *NASA Institute for Space Studies, New York*.—Last year Hayashi predicted that the sun would be fully convective in its early contracting phases and would have a high luminosity. This follows from the consideration that the photosphere must occur at about an optical depth of unity, which requires that the photospheric density and temperature must both be high in order to provide the necessary opacity. High temperature means high luminosity in the red-giant stage of contraction; high density and high luminosity both require deep outer convection zones. New models of the contracting sun have been constructed in which particular care has been taken to treat the outer convection zone and transition layer to the photosphere realistically. Opacities were calculated from the Los Alamos code and crude line corrections were added. For purposes of general exploration the models were not calculated in strict evolutionary sequences, but rather subject to the restriction that dr/dt is constant throughout the model. We find that Hayashi's predictions are fully correct. For all models with $R/R_\odot \geq 3$, convection is complete to the center. It is possible to construct models out to several hundred solar radii (but see accompanying Cameron-Ezer paper, p. 572); their surface temperatures lie in the range 2600°–4600°K. One can roughly say that these models descend nearly vertically in the color-magnitude diagram. The luminosity reaches a minimum near $R/R_\odot = 1.7$ and then increases slightly as further contraction proceeds towards the main sequence.

Atmosphere of Mercury. GEORGE B. FIELD, *Princeton University Observatory*.—Kuiper pointed out that radiogenic argon might be retained by Mercury. The critical temperature for escape in 4.5 billion years is 1800°K, close to the actual daytime temperature of the exosphere of the earth. The at-

mospheric polarization observed by Dollfus may be tentatively ascribed to argon since other abundant gases are much lighter and have therefore certainly escaped. Dollfus' measures imply a surface pressure of 1.2 mm and an atmospheric mass of 3.3×10^{18} g.

The pressure corresponds to a sublimation temperature of 56°K . Mercury's coldest spot must be hotter than this if the argon is not to freeze out. Walker calculated an antisolar temperature of 28°K , maintained by conduction of solar energy and by radioactive decays. Maintaining at least 56°K would require at least a thirty-fold increase in conductivity or radioactivity. Alternatively, atmospheric circulation at ~ 10 m/sec with temperature differences of $\sim 10^\circ\text{K}$ would also suffice. However, the observed near-agreement between insolation and infrared emission at the subsolar point suggests that the heat convected to the dark side is less than 3% of the total, and therefore, that the dark-side temperature is less than 250°K .

The suggested dark-side temperatures, between 56° and 250°K , may be detectable by radio methods. For example, 250°K would give a 340° disk temperature at 90° phase angle (including the known bright-side contribution), in agreement with the $370 \pm 125^\circ\text{K}$ observed at 3-cm wavelengths by Howard *et al.* Whether the dark side is really warm could be tested by observations near conjunction, predicted antenna temperatures (85-ft antenna) lying between $0^\circ 02$ and $0^\circ 09$ K for the conjunctions of October 1962 and January 1963.

The quoted mass of argon is 1.0×10^{-8} times the mass of Mercury; the same ratio for the earth is 1.1×10^{-8} . Is this merely a coincidence, or is it because the mass and constitution of the mantle, the effectiveness of degassing, and the lack of exospheric escape are all similar for the two planets?

Model-Independent Correlation of Stars and Neutral Hydrogen. GEORGE B. FIELD AND EMERY S. FLETCHER, *Princeton University Observatory*.—Radial velocities of O and B stars, Population I Cepheids, galactic clusters, and $H\text{ II}$ regions are compared with those of $H\text{ I}$ gas at the same galactic longitude. The method used is to obtain distances from the stellar velocities in accordance with the Schmidt model of galactic rotation and to superpose the resulting plot on the combined Leiden–Sydney map of the distribution of neutral hydrogen. A strong correlation is found, the stars falling preferentially at the same velocities as the gas arms. Where a random distribution would have caused 39% of the stars to be found in the gas, the observed percentages ranged from 73% for the Cepheids to 85% for the clusters; many, if not all, of the poorly correlated cases may be accounted for by real and observational

dispersion in the stellar velocities. This correlation is independent of the model of galactic rotation, a change in the model producing identical changes in the positions of stars and gas. The most straightforward interpretation is that stars and gas are associated, and that the poor correlation between distances of stars and gas found by several investigators is due to errors in the distance assigned to the neutral hydrogen.

In principle, this method may be used to order the ages of objects less than a few times 10^8 yr old by noting the relative tendency to lose correlation with the gas. It may also be used to assign distances to gas features when stellar distances are available.

Final Model of β Lyrae. SERGEI GAPOSCHKIN, *Harvard College Observatory*.—Among puzzling binaries, β Lyrae has occupied a foremost place. More than half a century of spectroscopic and photometric observation has brought a bewildering array of phenomena, many of which have defied explanation. Two main mysteries surrounded the system, one with reference to the invisibility of the eclipsing star, the other to the mass of the eclipsed one (and consequently to the mass of the whole system, often given as about 100 solar masses).

In 1936, I suggested a solution to the first mystery, finding it necessary to assume that the eclipsing star should be the smaller; so the constant visibility of the spectrum of the brighter one was explained. Twenty years later I proposed another solution which removed the major puzzle by concluding that the eclipsing star, although smaller and apparently fainter, must be heavier (not lighter), must have a smaller range in radial velocity, and must have a thick moving envelope changing or camouflaging its spectrum so that according to the theory of moving envelopes a set of emission rather than absorption lines may occur.

Support for this conclusion was found in the old observations made in 1896 exhibiting the predicted range of radial velocity of the heavier component. It appears now that the recent astrometric determination of the absolute magnitude of the visible star, on the basis of the assumption that β Lyrae is astrometrically a multiple system, provides a striking confirmation of the model proposed six years ago. The astrometric value of the magnitude is $-4^m 1$ (*Astrophys. J.* 135, 429, 1962), mine was $-4^m 01$ (*Z. Astrophys.* 39, 133, 1956). Thus, the current model of β Lyrae, no longer enigmatic, may be said to consist of two moderately heavy stars, surrounded by a large atmosphere in which all sorts of orthodox streams and currents occur. In addition, there is a moving envelope around the heavier star, making it fainter and camouflaging its spectrum.

Magnetosphere of the Moon. THOMAS GOLD, *Cornell University*.—The flow of solar plasma in the vicinity of the moon is considered and the magnetic configuration is deduced which will be set up by the interaction of the flow with the moon. It is shown that this configuration will tend to hold off solar plasma and prevent it striking the moon. A small magnetization of the moon will result. Variations in this magnetization will be measurable and will allow deductions to be made concerning the conductivity of the moon's interior.

An Attempt to Observe 21-cm Line Emission from Intergalactic Hydrogen. S. J. GOLDSTEIN, JR., *Harvard College Observatory*.—In an expanding universe containing a uniform distribution of optically thin neutral hydrogen atoms, the brightness temperature of the sky has a step in its spectrum at 1420.4 Mc. The amplitude of this step is proportional to the density of neutral hydrogen; for a Hubble parameter 100 km/sec/Mpc, a density of 10^{-4} atoms/cm³ gives a step of 1.7°K, if the gas is in thermal equilibrium. Field (*Astrophys. J.* **135**, 684, 1962) has attempted to observe 21-cm absorption in the spectrum of Cyg A. Observing emission has the advantage that the measurement is independent of the spectra of sources and that the observed brightness temperature can be expressed as a density without knowledge of the excitation temperature of the gas.

A horn antenna with half-power beamwidths of 24×27 deg, directed at the celestial north pole, fed one input of a switched maser radiometer. A coaxial resistor immersed in liquid nitrogen fed the other input. The difference in available power of these sources was measured at 2.0-Mc intervals with 1.0-Mc bandwidth over a 16-Mc range centered at 1420.4 Mc. The system noise temperature was 190°K including 110°K from the antenna and feed line. Test spectra of the temperature difference between two coaxial resistors clamped together in an aluminum block at room temperature show a maximum variation of 0.4°K at the switch terminals (0.6°K in brightness temperature). Observations of the north pole show an increase in brightness temperature of 2°K at the center frequency with respect to the adjoining frequencies. I conclude that a step as large as 1.1°K is not present, if the random motions of the intergalactic gas have a standard deviation of 400 km/sec or less. Hence, subject to the same condition, the average density of optically thin neutral hydrogen atoms in the intergalactic space observed is less than 7×10^{-5} atoms/cm³.

This research was supported by the National Science Foundation.

Evolution of Massive Stars after Helium Burning. CHUSHIRO HAYASHI AND ROBERT C. CAMERON, *Kyoto University and Goddard Space Flight Center*.—Our model of a star of 15.6 solar masses, with age-zero composition, $X=0.90$, $Y=0.08$, and $Z=0.02$, has been modified to include an outer convection zone for the carbon-burning phase, whose onset was treated in our Paper III (*Astrophys. J.* **136**, 166, 1962).

The following general expectations of Paper III have been confirmed: (1) the stellar radius is now considerably reduced, (2) the parameters describing the interior were already essentially correct, and (3) the red-supergiant branch in clusters such as h and χ Persei is accounted for by carbon burning and later evolutionary phases.

For the early stages of carbon burning, we find $E \approx 15.5$, with the bottom of the outer convection zone so deep ($q \approx 0.27$) as to cause mixing of previous hydrogen-burning products. Detailed computations have also been made for the contraction phase (lifetime: 0.5×10^5 yr) which precedes carbon burning.

The carbon-burning lifetime proper has an upper limit of 2.3×10^5 yr. The total lifetime from central helium exhaustion to iron decomposition (supernova explosion) has been carefully estimated to be nearly 11×10^5 yr, agreeing well with the preliminary estimate of Paper III and with star-count ratios in the appropriate regions of the H-R diagrams of h and χ Persei and NGC 330.

The inclusion of neutrino loss, due to the $(\bar{\nu}e)(\bar{\nu}e)$ interaction, as proposed by Feynman and Gell-Mann, and computed by Chiu *et al.*, shortens drastically the above-mentioned lifetimes and destroys the observational agreement mentioned. Assuming, and later verifying, nondegeneracy of the core, we find that, when neutrino effects are considered, the lifetime as a red supergiant becomes essentially that of carbon burning (now reduced to only 0.8×10^5 yr, at most), since, for example, the neon- and oxygen-burning lifetimes are reduced by factors of 10^3 and 10^4 , respectively.

The observational data of h and χ Persei therefore strongly suggest that the $(\bar{\nu}e)(\bar{\nu}e)$ interaction is forbidden.

UBV Photometry of DH Cephei. ARNOLD M. HEISER, *Dyer Observatory*.—DH Cephei is a spectroscopic binary located in the galactic cluster NGC 7380 and consists of two early O stars. The spectroscopic observations (Pearce, J. A., *Astron. J.* **54**, 135, 1949) indicated an inclination of 62 deg and prediction of an eclipse amounting to 0.1 mag. at primary minimum. *UBV* observations of this suspected eclipsing variable have been obtained and

are incorporated with the *UBV* data of NGC 7380 (Hoag, A. A., *et al.*, *Publs. U. S. Naval Obs.* 17, Part 7, 1961) to ascertain further physical quantities of the binary components.

Tentative Model of Beta Lyrae on the Assumption of a Less Massive Primary. SU-SHU HUANG, *Goddard Space Flight Center.*—That the primary component of Beta Lyrae should be regarded more massive than the secondary is inferred from the observed difference in luminosity of the two components. Struve, Sahade, and S. Gaposchkin have individually pointed out the difficulties and even suggested the contrary view of a primary less massive than the secondary. A serious consideration of this contrary view becomes necessary after the recent work by Abt, Jeffers, Gibson, and Sandage on the distance of this binary system, since the absolute luminosity thus determined by them can only lead to a less massive primary. How could we then account for the observed difference in luminosity of the two components? It is proposed here that the secondary has received a large amount of mass from the primary but has not yet adjusted the luminosity to its new mass. Also, an opaque disk rotating around the secondary, formed from the ejected matter from the primary, obscures partially the light coming from the secondary. The presence of this disk would also make the system look photometrically like a contact binary while it actually could be a semidetached system, i.e., only the primary has filled the equipotential lobe. The emission lines of a large positive γ velocity, discussed by Sahade, perhaps come from the gas falling into the secondary star from the space on both sides of the opaque plane. Their profile shows a peak to the red of the center of the broad feature according to Sahade and Houziaux, and is therefore in agreement with the proposed model because we observe most emitting atoms only on one side of the plane if the inclination of the system is not exactly 90° .

Distribution of Binary Stars among the Bright Stars. J. ALLEN HYNEK, *Dearborn Observatory.*—Since present theories of stellar evolution indicate that the normal giants were once members of the upper main sequence, it might be expected that the frequency of binaries among the giants should be essentially the same as that among the present upper-main-sequence population.

Using the well-observed naked-eye stars alone, nearly one-half of which are G, K, and M giants, as a test population, we find that the present upper main sequence has more than twice as many spectroscopic and visual binary systems as the present giants do.

Further, the median period of 248 BAF spectroscopic binaries is 8 days; that of 76 GKM spectroscopic pairs, 463 days.

The implications of these paradoxical relationships appear to offer material for discussion. It is tentatively suggested that those members of an earlier epoch main sequence that possessed short-period spectroscopic companions, either singly or as members of a visual system, could not have become the primaries of binary systems now on the giant branch of the H-R diagram. Mass transfer and a possible cataclysmic stage as the main-sequence primary expanded to engulf the short-period secondary would remove this system from the conventional evolutionary path. This would account for a deficiency both of spectroscopic and visual binaries on the giant sequence, the "missing" visual systems being those whose primary was once itself a short-period spectroscopic system.

Computer Program for a Steady-State Gas. HOLLIS R. JOHNSON, *Yale University Observatory.*—In radiating gases of interest in astrophysics the existence of strict (or even "local") thermodynamic equilibrium is questionable. The absence of such equilibrium leads to departures from the equilibrium distribution functions, but quantitative estimates of the amount of these departures have not been established for many interesting cases. In general, the answer requires the solution of the set of equations expressing steady state in the atomic energy levels and the set of equations of radiative transfer through the gas.

A computer program which will permit comprehensive calculations of steady-state equilibrium in a self-consistent fashion is described. The complete program performs these operations: (1) Determines the monochromatic optical depth $\tau_\nu(\tau_0)$, where τ_0 is some standard optical depth, using as input the absorption coefficients κ_ν and κ_0 throughout the atmosphere. (2) Finds $J(\nu, \tau_0)$, the mean intensity, at all wavelengths of interest and at many points throughout the atmosphere from the integral equation. Input for this program are $S(\nu, \tau_0)$ and $\tau_\nu(\tau_0)$, where S is the source function. (3) Calculates the integral of the mean intensity and the absorption coefficient over frequency for all transitions of interest at many values of τ_0 . Input are $J(\nu, \tau_0)$ and the absorption coefficients as functions of frequency. (4) Solves the steady-state equations to obtain at a number of points in τ_0 the population of the energy levels, κ_ν , and S_ν . The input data are the radiation fields (step 3), atomic parameters, collisional and radiative cross sections, and the temperature and density throughout the gas.

An application of the program is made to obtain a self-consistent solution for neutral helium in the shell surrounding a star. A five-level model atom is employed in an atmosphere of constant $T_e=10\,000^\circ$, $N(\text{He})=10^{11}\text{ cm}^{-3}$, and $N_e=10^{12}\text{ cm}^{-3}$ which is excited by a B star.

Evidence for Nitrogen Dioxide in the Martian Atmosphere. C. C. KIESS, C. H. CORLISS, AND H. K. KIESS, *Georgetown College Observatory and National Bureau of Standards*.—During the oppositions of Mars in 1956 and 1960–61 we obtained several spectrograms of the planet and the moon, with our concave-grating spectrograph giving a dispersion of 5 Å/mm . Spectra of the moon were recorded in juxtaposition with those of Mars when the moon, on the same date, was as near as possible in altitude to that of Mars.

Microphotometer tracings of these spectra show not only the decline of the Martian spectrum toward shorter wavelengths relative to that of the moon, but also numerous dips and depressions in the spectrum of Mars that do not occur in the lunar spectrum. These features suggested absorptions by an unknown atmospheric constituent of the planet. Comparisons with published laboratory data showed an almost one-to-one correspondence between the Martian features and those of nitrogen peroxide. This induced us to observe the absorption spectrum of this gas from the near infrared to the vicinity of 3600 Å . Nearly all the so-called bands revealed in this investigation from $H\alpha$ to the H and K lines agree in wavelength with similar absorptions in Mars. Of particular interest is the spectrogram of 12 October 1956 taken at a time when other observers noted a yellow veil over the planet. About 100 Martian features between $H\alpha$ and 6150 Å agree in wavelength with absorptions measured on a laboratory spectrum of NO_2 .

Secular Perturbations of Asteroids with High Inclination and Eccentricity. YOSHIHIDE KOZAI, *Smithsonian Astrophysical Observatory*.—Secular perturbations of asteroids with high inclination and eccentricity moving under the attraction of the sun and Jupiter are studied on the assumption that Jupiter's orbit is circular. After short-periodic terms in the Hamiltonian are eliminated, the degree of freedom for the canonical equations of motion can be reduced to 1.

Since there is an energy integral, the equations can be solved by quadrature. When the ratio of the semimajor axes of the asteroid and Jupiter takes a very small value, the solutions are expressed by elliptic functions.

When the z component of the angular momentum (that is, Delaunay's H) of the asteroid is smaller than a certain limiting value, there are both a stationary solution and solutions corresponding to libration cases. The limiting value of H increases as the ratio of the semimajor axes increases, i.e., the corresponding limiting inclination drops from 39.2 to 1.8 as the ratio of the axes increases from 0.0 to 0.95 .

Study of Degeneracy in Very Light Stars. SHIV S. KUMAR (National Academy of Sciences—National Research Council, Postdoctoral Research Associate with NASA), *Institute for Space Studies*.—Convective models have been constructed for stars of mass $M < 0.1M_\odot$. For a given mass, the internal structure of the star is computed at several radii, taking into account nonrelativistic degeneracy of the stellar material. It is found that for these stars the central temperature has a maximum value at one particular radius and this value decreases when the radius is greater or smaller. For example, for a Population I star of mass $0.05M_\odot$ the maximum temperature of two million degrees occurs at a radius of $0.12R_\odot$, and the temperature decreases if we increase or decrease the radius. For Population I stars with mass less than $0.05M_\odot$ the central temperatures and densities are too low for energy production by the proton-proton cycle and even for a star of mass $0.07M_\odot$ this process may not produce sufficient energy. Application of these models is made to contracting stars and it is found that Population I stars of mass less than $0.05M_\odot$ become degenerate objects and that they do not reach the main-sequence stage. The limiting mass for such an evolution may be as high as $0.07M_\odot$. For a star of mass $0.05M_\odot$, the time scale for contraction to the stage of maximum central temperature is estimated to be approximately one billion years, which is small as compared with the age of the galaxy. After this stage the star begins to cool slowly and evolves towards a completely degenerate configuration. Thus it becomes a "black" dwarf without ever going through normal stellar evolution.

Steady-State Atomic Populations in Stellar Atmospheres. MYRON LECAR, *Yale University Observatory and Institute for Space Studies*.—A collection of computer programs has been developed to compute steady-state atomic populations in stellar atmospheres. First, the temperature distribution satisfying radiative equilibrium is calculated, assuming LTE. Then, the resulting radiation fields are used to compute the non-LTE steady-state populations. The calculation is applied to the atmosphere of an A0 star, assuming a composition of pure hydrogen.

Steady-state populations for pure hydrogen have been computed for dilute Planck radiation fields, for hydrogen densities ranging from 10^{14} to 10^{16} cm^{-3} and for election temperatures ranging from 7000 to 30 000°K. By successively adding bound states it has determined that six bound states are sufficient to calculate the populations of the first five bound states to an accuracy of 0.1%.

1417 Mc/sec Observations of M31. JOHN W. LEIBACHER, *Harvard College Observatory* (Introduced by A. E. Lilley).—M31 has been observed with the maser-equipped Harvard radio telescope at 1417 Mc/sec. The low-noise temperature of the radiometer permitted the detection of the halo of M31 over a region $3.5 \times 8^\circ$. The integrated value of the flux density was found to be $(68 \pm 17) \times 10^{-26}$ W m^{-2} cps^{-1} . Comparing this with the 408 Mc/sec flux density of Large, Mathewson, and Haslam (*Nature* **183**, 1250, 1959) gives a spectral index of -0.58 ± 0.13 . This value should be compared with the spectral index obtained from the flux determinations at 408 and 158 Mc/sec (Hanbury, Brown, and Hazard, *Monthly Notices Roy. Astron. Soc.* **119**, 297, 1959) of -0.38 ± 0.11 . Brightness temperature profiles along the major and minor axes at 408 and 1417 Mc/sec have been compared; and the 1417 Mc/sec data have been smoothed and compared with the 38 and 178 Mc/sec data of Baldwin and Costain. Both of these comparisons of the distribution of emission show no significant differences in the distribution of intensity across M31, and are thus in agreement with the results of Costain and of Adgie and Smith for the Galaxy.

The Harvard Radio Astronomy Project is supported by the National Science Foundation.

On the Nature of the Wolf-Rayet Phenomenon. D. NELSON LIMBER, *Yerkes Observatory*.—The basic observational data pertaining to the Wolf-Rayet phenomenon have been reviewed with particular reference to the Wolf-Rayet component of V 444 Cygni. The validity of the conventional interpretation in terms of the radial ejection of matter at high speeds has been re-examined. The conventional interpretation appears to be inadequate on a number of independent counts.

An alternate interpretation of the phenomenon in terms of a forced rotational instability consequent to contraction in a post-main-sequence evolutionary stage is put forward. This latter interpretation appears to account for many of the available observational data in a much more satisfactory way than does the conventional one and does not appear to be inconsistent with any of the data studied thus far.

Spectrum of Coherent Electron Waves in the Solar Corona. J. M. MALVILLE, *University of Michigan, Radio Astronomy Observatory*.—The interaction between density fluctuations of an electron stream and longitudinal waves carried by the plasma through which the stream is passing can lead to an effective conversion of kinetic energy into energy of electron oscillations. Using simple energy considerations the steady-state amplitudes of coherently excited electron waves are calculated. To take account of the bandwidth of the oscillations a Gaussian profile is used for the rate of transitions from the free to trapped electron energy states. Assuming Rayleigh scattering for transforming the longitudinal waves into transverse waves, the radio-frequency spectrum is calculated for an electron stream with a velocity corresponding to type-III bursts. A turnover in the spectrum is predicted near 50 Mc/sec. Between 70 and 140 Mc/sec the slope of the spectrum agrees reasonably well with Wild's (Wild, J. P., *Australian J. Phys.* **3**, 541, 1950) statistically determined spectrum of type-III bursts. The frequency of the turnover and the slope of the spectrum are critically dependent upon the temperature gradients in the corona. It is suggested that type-III bursts first become visible at frequencies corresponding to the steep temperature gradient in the chromosphere-corona transition region. This work was supported by National Aeronautics and Space Administration.

Precision of the Estimates Relating to the Selection Factor of Galaxies. ALLAN H. MARCUS, *University of California, Berkeley*.—A catalogue of galaxies for which the criteria of selection depend essentially on only magnitude and type is characterized by a "selection factor" $\Phi(m, t) = \text{probability that a galaxy of type } t \text{ and apparent magnitude } m \text{ will be included in the catalogue}$. Neyman and Scott suggested (*Handbuch der Physik* **53**, 416, 1959) the approximation $\Phi(m, t) = \text{Gaussian probability integral from } (m - \alpha_t)/\beta_t \text{ to infinity}$, where α_t and β_t are parameters. Subsequently (Neyman and Scott, *Astron. J.* **66**, 148, 1961) maximum likelihood estimates of α and β were published for eight types of field galaxies, together with estimates of mean absolute magnitude M_0 and dispersion σ_M . However,

Type	M_0	σ_M	α	σ_α	β	σ_β	ρ	n
E0–E3	–18.0	1.02	12.0	0.4	1.13	0.12	–0.94	83
E4–E7	–17.0	1.33	10.9	0.9	1.30	0.22	–0.95	28
SB0	–18.5	0.58	11.4	0.8	1.10	0.23	–0.94	21
SBb	–18.7	0.86	11.2	0.7	1.10	0.21	–0.94	26
S0, S0p	–17.0	1.25	11.7	0.5	1.10	0.13	–0.94	66
Sa, Sap, Sab	–18.4	0.82	11.9	0.4	0.90	0.13	–0.92	51
Sb, Sbc	–17.9	1.05	10.8	0.4	1.10	0.12	–0.94	77
Sc, Scp, SBc	–16.7	1.28	11.6	0.2	0.75	0.08	–0.90	94