the convection zone is of crucial significance, the deeper the zone the smaller the importance of diffusion.

The Hydrogen-Deficient Star HD 96446. L. H. ALLER, W. BUSCOMBE, AND T. DUNHAM, JR., *Mount Stromlo Observatory*.—HD 96446 (galactic coordinates $l^{\text{II}}=290^\circ$, $b^{\text{II}}=0^\circ$) is an early B-type star with abnormally strong helium lines (Jaschek 1959). In view of the pronounced Stark broadening of the hydrogen and helium lines, we suggest that the star must be on, or below, the main sequence. Photoelectric spectrum scanner measurements show the star to be rather blue, apparently because Balmer line and continuous absorption are unusually weak.

The sharpness of absorption lines of heavier elements, noted previously at lower dispersions, is confirmed by coude spectrograms (dispersion 6.7 Å/mm). The radial velocity and proper motion are representative of stars more distant than those a few degrees further north of the galactic plane which share in the stream motion of the Scorpio-Centaurus association. The H and K lines show two components, of which at least one set is probably of interstellar origin.

About the 400 lines for which wavelengths have been measured in the spectrum interval $\lambda\lambda 3550\text{--}4850$ are those due to H, He, CII, CIII, NII, NIII, OII, OIII, NeII, SiIII, SiIV, PII, SII, ClII, ArII, KII, CaII, FeIII, FeIV, and NiIII, and probably KrII. Several lines of the latter ion appear to be present although perhaps the strongest, $\lambda 4355$, has not been found. Most of the unidentified lines are very weak. Analysis of the spectrum by curve of growth and model atmosphere methods is being undertaken.

Contribution of Absorption Lines to the Opacity of Matter in Stellar Interiors. ALBERT ARKING AND JACKSON HERRING, *Institute for Space Studies, New York*.—Using line profiles calculated from the theory of Griem, Kolb, and Shen, the Rosseland mean opacities for several stellar mixtures in the range of 10^6 to 2×10^7 °K have been computed. The corrections to the continuous opacity are less than 15% except at temperatures $T \lesssim 3\times 10^6$ and values of the ionizing potential $\psi \lesssim 8$ kT.

The Griem-Kolb-Shen theory (Griem, H. R., *et al.*, *Phys. Rev.* **116**, 4, 1959) which includes the effects of both ions and electrons, was modified to take into account the electrostatic repulsion between the perturbing ions and the radiating ions by a method similar to that of Baranger and Mozer (*Phys. Rev.* **115**, 521, 1959). This procedure is necessary for the heavier elements.

The continuous opacities and the occupation numbers were taken from a program developed at Los

Alamos (opacity code supplied to us by A. Cox), although our results, expressed as the line contribution divided by the continuous opacity, are essentially unchanged when simpler methods for the continuous background and occupation numbers—e.g., the method of Strömgren (Strömgren, B., *Publ. Copenhagen Obs.* **83**, 118, 1932)—are used.

Ionization and Excitation Equilibrium of CaII in the Solar Atmosphere. R. GRANT ATHAY, *High Altitude Observatory*, AND J. B. ZIRKER, *Sacramento Peak Observatory*.—The ionization and excitation equilibrium of CaII is computed for conditions appropriate to the outer solar atmosphere. Populations of energy levels are found to depart sharply from the LTE populations at corresponding values of T_e . The dominant ionization process of CaII is shown to be via collisions from the 3D level and to exhibit fundamental differences from H, He, and MgI. For electron densities and temperatures such as are expected in most of the chromosphere and corona, the ratio CaII/CaIII is shown to be inversely proportional to electron density, rather than directly proportional as predicted by the Saha equation. The 50% ionization stage of CaII is reached for $T_e = 10\,000\text{--}14\,000^\circ$, depending upon the degree of radiative detailed balance attained in the H and K lines. Computations of opacities and occupation numbers indicate that the CaII H and K lines in the Fraunhofer spectrum are formed near the top of the hydrogen temperature plateau where $T_e \approx 13\,000^\circ$ and that $T_e \approx 15\,000^\circ$ in the regions within spicules and prominences where the CaII emission arises.

A Spectrum of the Sun from 168 to 700 Å. W. E. AUSTIN, J. D. PURCELL, AND R. TOUSEY, *U. S. Naval Research Laboratory*.—A solar spectrum covering the range 168 to 700 Å was obtained on June 21, 1961, from an Aerobee-High rocket. The instrument employed a 40-cm radius, 600-line/mm grating at 85° incidence. Stray light background was completely eliminated by introducing a self-supported film of aluminum, 1000 Å thick, in front of the slit. This was opaque for $\lambda > 850$ Å, but transmitted well from 700 Å to about 170 Å, the $L_{2,3}$ edge of Al. An exposure of 110 sec made from 186 km to peak at 197 km and down to 178 km, showed approximately 50 emission lines between 167.9 and 310 Å, in one or more orders. Twenty-five lines were photographed in three orders, and a line at 174.5 Å in four. From 310 to 700 Å only a few faint lines were recorded, because of strong attenuation at these wavelengths by N_2 remaining overhead.

Wavelengths have been established to 0.1 or 0.2 Å. Surprisingly few lines agree with lines that are