Only the very smallest cometary meteoroids can spiral into the sun by the Poynting–Robertson effect if the assumptions of this paper are true. Strong evidence for a high concentration of dust in the earth's immediate vicinity suggests that erosion rates may be greater near the earth than in deep space from Venus to Mars and even beyond. The surface of space vehicles should be carefully chosen or protected to minimize etching by interplanetary dust, if long exposure times are anticipated. The hazard may be greatest at the moon's surface.

Theoretical Aspects of the W Virginis Phenomena. CHARLES A. WHITNEY, *Smithsonian Astrophysical Observatory, Harvard College Observatory*.—The physical basis of an earlier shock-wave model for the atmospheric motions of W Virginis is discussed. Implications of this model for spectrum line profiles and curve-of-growth turbulence are shown to be either (a) consistent with observations or (b) undetectable from present data. The expected duration of the doubling of absorption lines is computed and found to agree with observed values. Results of recent calculations on the structure of the shock front are presented and are used for a preliminary interpretation of the observed hydrogen-emission lines.

Age of Messier 3. N. J. WOOLF, *Princeton University Observatory*.—Sandage (*Astrophys. J.* **126**, 326, 1957) studied the energy expenditure of stars in M3 using semiempirical evolutionary tracks. He found that if the absolute magnitude of the RR Lyrae stars is 0.0, for a cluster of 5.1×10^9 years, stars emitted energy equivalent to conversion of their entire mass from hydrogen to helium. Woolf (*Astrophys. J.* **135**, 644, 1962) pointed out the discrepancy between this result and estimated large ages using stellar models by Hoyle (*Monthly Notices Roy. Astron. Soc.* **119**, 124, 1959).

The study by semiempirical tracks has been repeated using Hoyle's stellar models as a basis. There are two modifying effects on the previous conclusion. The semiempirical study did not allow for evolved stars being slightly more massive than those at the turnoff point. The age for complete fuel consumption becomes 6.8×10^9 years. Sandage's study used theoretical early evolution tracks giving a 0.94-mag. rise from main sequence to turnoff point, while Hoyle's tracks give a rise of 1.45 mag. This affects the result through the derivation of the initial luminosity function. The maximum permissible age rises from 6.8×10^9 to 10.5×10^9 years. For comparison, Hoyle's models give the age as 12.0×10^9 years using M_v RR Lyrae stars=0, $m_v=15.67$ and for the turnoff point $m_v=+18.79$.

This discrepancy is still serious because the age from semiempirical tracks is a maximum one, not

allowing for incomplete combustion, rapid phases at high luminosity, or a possible 20% mass loss (Oort, J. H., and van Herk, G. *Bull. Astron. Inst. Neth.* **14**, 299, 1959; Woolf, N. J. *Publs. Astron. Soc. Pacific* **73**, 339, 1961). However, Hoyle's models still have uncertainties in the nuclear reactions and may give ages 30% too large (Hoyle, F. Report to IAU Commission 35). The age then, 8.4×10^9 years, would be consistent with the semiempirical tracks. Uncertainties in the absolute magnitude of the RR Lyrae stars are large. The latest value is +0.25 (Arp, H. C. *Astrophys. J.* to be published), and there are also uncertainties of helium abundance that may reduce these ages. There is no good evidence that M3 is more than 10^{10} years old.

Temperature Effects in Photomultipliers. ANDREW T. YOUNG, *Harvard College Observatory*.—The effect of temperature changes on the spectral response curves of RCA 1P21, EMI 6256 and 9558, and Farnsworth FW-118 photomultipliers has been measured for two tubes of each type. The cathodes are respectively S-4 (Sb-Cs), S-13 (Sb-Cs), S-20 (Sb-Na-K-Cs), and S-1 (Ag-O-Cs). All but the FW-118 have Sb-Cs dynode surfaces.

Cooling from room temperature to dry-ice temperature causes changes in relative spectral response of the order of 10% or less for the Sb-Cs cathodes for $\lambda \leq 5000$ Å, for the S-20 cathode for $\lambda \leq 6250$, and for the S-1 cathode over at least the range from 6250 to 10 000 Å. At $\lambda=7300$, the S-20 cathode loses about 30% of its normal sensitivity when it is refrigerated. The data are consistent with a temperature coefficient of $-0.04\%/^{\circ}\text{C}$ per stage for the Sb-Cs dynodes.

The Sb-Cs cathodes lose a large fraction of their sensitivity for $\lambda > 5000$ Å when they are refrigerated. This loss is more than enough to account for the discrepancy between observed $B-V$ colors of stars and colors computed from stellar atmosphere theory. The coefficient of the temperature effect between 5000 and 6000 Å is 1.12×10^{-5} mag./ $^{\circ}\text{C}/\text{Å}$. This typically requires a temperature stabilization of one or two degrees if a 1P21 is required to carry a color zero point in this part of the spectrum with an accuracy of 1%.

Preliminary Report on a New Computation of the Orbit of Halley's Comet. PEDRO E. ZADUNAI-SKY, *Smithsonian Astrophysical Observatory and Faculty of Sciences, Buenos Aires, Argentina*.—An improvement of the orbit computed by Cowell and Crommelin has been made by J. Bobone, from Cordoba. He used 1146 observations covering the period from August 1909 to March 1910. A final computation is being made based on Bobone's results and using 2862 observations made between August 1909 and May 1911.