made relative to 3C71, and their energy calibration thus depends on its spectrum, and should be revised slightly to agree with its spectral index as newly determined by Kellerman. This correction to the calibration is also indicated by the fact that with the previous calibration NGC 7027, which is optically very thick, has a steeper spectral slope than a black-body. Comparison of the observed fluxes at the two frequencies enables the optical depth of each observed planetary to be determined, relatively independently of the density, temperature, and distance. The derived optical depths at 1400 Mc/sec are as follows:

NGC 6720	0.1	NGC 6543	0.9
NGC 6853	0.2	IC 418	>0.9
NGC 7662	0.2	NGC 7027	~20.
NGC 7009	0.5		

These optical depths are mostly in approximate agreement with expected values derived from optical-frequency measurements of the planetary nebulae. However, the radio-frequency optical depth of NGC 7027 is larger than expected, and sets it apart from the other planetaries. There is also other evidence from optical-frequency observations that this nebula is a different type of object from the typical planetary nebulae.

Evolution of Extended Radio Sources. A. G. PACHOLCZYK, *Joint Institute for Laboratory Astrophysics, Boulder*.—The dynamics and radiation of an expanding plasma containing relativistic particles and a magnetic field are examined. The evolution of an extended radio source located outside the parent galaxy, approximated to by the expanding plasma, is considered. Physical properties of extended radio sources are calculated for the evolutionary stage in which production of relativistic particles was terminated. In this stage the dimension of a typical source is 1 kpc (0.1–10 kpc), the total energy is 10^{60} erg (10^{59} – 10^{62} erg) and the magnetic field is 10^{-1} G (10^{-2} – 10 G). It is shown that the assumption that production of relativistic particles stopped at some earlier stage of expansion is compatible with the observational data concerning extended radio sources. The energy requirement can be lowered if the radiation of extended sources is not supposed to be isotropic. Correlations between separation of components of double sources and other properties of these sources (size, luminosity) suggest some degree of anisotropy in their radiation.

A New Class of Long-Period Trojan Librations. EUGENE RABE, *University of Cincinnati Ob-*

servatory.—Numerico-analytical results show the existence of a new class of periodic librations, of nonsimple shape, about the two equilateral points of the restricted problem of three bodies. Using the Fourier series representations of the “simple-shaped” periodic librations (Rabe, *Astron. J.* **66**, 500; **67**, 382) as reference solutions, any superimposed oscillations of short periods are found to produce a normally nonperiodic combined solution, which represents the most general form of librational motion in the sun–Jupiter plane. For any given basic or reference orbit, the characteristic exponent c associated with the superimposed oscillations is a certain constant, independent of oscillation amplitude, as long as only the first and second powers of the displacements are considered in the differential equations of motion. c becomes a function of oscillation amplitude, however, when the third-order terms are included, and this leads to integral values of c and thus restores periodicity of the total librational motion for a discrete set of specific amplitudes about the basic periodic solution. This finding of the existence of long-period librations of more complicated forms is in agreement with observations of orbital trajectories actually obtained by numerical integration for a variety of starting conditions.

A Photoelectric Area Scanner. KARL D. RAKOS, *University Observatory, Graz, Austria, and Lowell Observatory*.—This new astronomical instrument is designed primarily to measure magnitude differences of very close double stars, or of a planet and its satellite.

The fundamental principle is based on a linear scanning technique, the use of a narrow slit and a data presentation system which includes a linear digital output in addition to photographic recording with an oscilloscope.

As the instrument was recently tested on the Navy's new 61-in. telescope, the slit was 0'6 wide and 6'5 long. The area exposed to the sky was therefore about 4 sq sec of arc in area. In ordinary photoelectric work it is rarely possible (even under ideal conditions) for an observer to use less than a 10" aperture or about 80 sq sec of arc. The large “off” time during each scan is more than compensated by the fact that accurate photoelectric photometry can be performed on very closely separated objects such as double stars which have heretofore been entirely beyond the reach of photoelectric photometers.

Preliminary tests under poorer-than-average seeing conditions show stars 3'5 apart to be clearly separated. An eclipse of Phobos when only 5" from Mars has been observed when the brightness ratio was more than 10^5 .