1963 there occurred changes in the proper motions of the larger sunspot umbrae, a change in the position and configuration of the magnetic neutral line of the longitudinal magnetic field, and a rapid growth of penumbra in the group. An importance 1 flare on 19 September 1963 occurred near two strong spots of opposite polarity and at the time when their relative proper motions brought them closest together. Faint "lanes" or arcs in the photospheric granulation appeared between spots of opposite polarity at the time of maximum brightness of an importance 1 flare on 24 September 1963. The arcs lay directly under the flare.

These observations suggest that the larger and brighter flares may be accompanied by observable changes at the photospheric level. A study of more cases is needed to establish a more definite relationship between changes in sunspots and solar flares.

Positions of Certain Recurrent Flares, 12-16 July 1959, Relative to Sunspots and Features of the Calcium Network. SUSAN M. P. MCKENNA (introduced by Helen Dodson Prince), *The University of Michigan*.—A study is in progress of the flares and subflares that occurred in the great center of activity at $N 15^\circ$ that traversed the solar disk between 8 and 23 July 1959. A number of these flares consisted, in whole or in part, of brightenings of apparently the same portions of the calcium plage. One such pattern of repeatedly flaring regions dominated many of the flares observed 12-16 July inclusive.

During this interval, as has been shown already by Bruzek (*Z. Astrophys.* **50**, 110, 1960), the associated spot structure was undergoing continual change. At the photospheric level the relative positions, numbers, and field strengths of the sunspot umbrae changed radically with time. At the chromospheric level, the most notable feature of the flares was their marked similarity in position relative to one another and to features in the calcium plage. Seven individual areas with hard, well-defined outlines can be identified day after day as the site of flares and these impart to the separate events a common pattern. The components of this pattern retained their positions relative to one another in the chromospheric network in spite of the changing pattern exhibited by the underlying sunspots.

The leading spot which was the main umbra of south polarity drifted westward faster than the motion imparted to the chromospheric layers by solar rotation, and the flaring regions moved westward faster than did the following spots of northern polarity. As a consequence, several of the components of the flare pattern appeared variously with time over umbral, penumbral, and "normal" photospheric regions.

The study is based on concomitant records of flares and sunspots obtained with the spectroheliograph of the McMath-Hulbert Observatory, using a telescope of 40-ft focal length, and on photographs from the worldwide flare patrol organized by H. Smith.

Some Consequences of Large Redshifts. G. C. MCVITTIE, *University of Illinois Observatory*.—Seven zero-pressure models of the universe, whose characteristics are as different as possible, are employed to calculate the luminosity distances and the linear diameters corresponding to an angular diameter of $1''$, of sources of radiation with redshifts up to 3. The intrinsic emissions at 10^{15} cps of the nine quasi-stellar radio sources, whose redshifts are now known, are calculated in each of the seven models. These emissions turn out to be very similar in any one model, except perhaps in the model of the steady-state theory where an increase of intrinsic emission with redshift is suggested. If the emission at 10^{15} cps is indeed constant for quasi-stellar sources, a new determination of the value of the Hubble constant and of the acceleration parameter should be possible.

Beryllium in G- and G-Type Dwarfs. ANN E. MERCHANT, *Lick Observatory*.—The abundance of beryllium has been determined in 10 F- and G-type stars, for comparison with the lithium abundances in these same stars. One hundred twenty-inch coude spectrograms with a dispersion of 8 Å/mm were used. Since the resonance lines of Be II used for the analysis lie in the ultraviolet ($\lambda\lambda$ 3130, 3131) where the spectra are very crowded with metallic absorption lines, special care was taken in establishing the position of the continuum with consistency.

Two of the stars, α CMi and θ Cyg, were found to show no detectable Be; upper limits on the Be content of 25 times less than the solar value were assigned. Neither of these stars has a detectable amount of Li. The evolutionary effects of the companions of these two stars are considered as a possible explanation. The other eight stars and the sun all have approximately the same amount of atmospheric Be.

Four of these latter eight stars were found by Herbig (*Astrophys. J.* **140**, 702, 1964) to have appreciable Li^6 . The presence of Li^6 is taken to mean that there has been no destruction of Li or Be by convective mixing to high enough temperatures for (p, α) reactions to take place. The mean abundance ratio for these four stars is +0.6 with a variation about this mean of 50%; it is expected that a constant ratio of Li/Be should result if these light elements were produced by spallation reactions.