Observatory with a velocity range of 10.1 km/sec. A total of 19 published radial velocities exist covering the interval 1897–1924. In addition, 29 spectrograms have been secured at Allegheny Observatory covering the interval 1960–1963. Analysis of the radial velocities from the Allegheny observations revealed a 2.158 day period. The old observations also fit this period. These old observations, however, show a different velocity range, a marked change in the systematic velocity, and a remarkable change in the value of ω . The evidence points to a three-body system with strong apsidal motion for the close pair. Preliminary elements are:

1908	1963
$P=2.158$ days	$P=2.158$ days
$e=0.35$	$e=0.35$
$\omega=315^\circ$	$\omega=185^\circ$
$\gamma=+3.0$ km/sec	$\gamma=-6.0$ km/sec
Velocity range=18 km/sec.	Velocity range=13 km/sec.

The long period is still unknown.

Parallax determinations exist from Allegheny, McCormick, and Yale Southern Observatories. Unfortunately each of these series covers only a two-year interval, which is insufficient to show astrometric orbital variation. However, the values of μ_α determined by each observatory are significantly less than the GC value: a possible indication by itself of an astrometric orbit. The star is undoubtedly an astrometric binary. Both astrometric and spectroscopic observations are being continued at the Allegheny Observatory.

Variability among Red Giants. V. M. BLANCO AND L. PLAUT, *Warner and Swasey Observatory and Kapteyn Astronomical Laboratory*.—The observations published by Stebbins and Huffer (Stebbins, J., and Huffer, C. M., *Publ. Washburn Obs.* 15, 137, 1930) are analyzed statistically to determine the frequency of variability of different spectral types in the giant sequence. The variability of M-type giants in three selected regions is also studied. The results show that among the giants, variability is rare up to class K4 and that with advancing class the frequency of variables then increases to practically 100% for stars of class M4 or later. Up to class M7, however, the range of variability is generally less than 0.5 photographic magnitudes. Past class M7, the giants generally show well-defined large-amplitude variations.

Circularly Polarized Components of the Decimeter Radiation from Jupiter. T. D. CARR, S.

GULKIS, H. BOLLHAGEN, D. J. KENNEDY, AND A. G. SMITH, *University of Florida*.—Right and left circularly polarized components of the radiation from Jupiter were measured at 10, 15.8, and 22.2 Mc/sec. Measurements were made during the 1962, 1963, and 1964 apparitions at the University of Florida Radio Observatory and at the Maipú Radioastronomical Observatory in Chile. The quantity α was determined for each noise pulse, α being the difference between the square roots of the flux densities of the two circular components divided by the sum (negative for right circular). Algebraic averages of α for 20° intervals of System III longitude were computed and were plotted as functions of longitude. The function was almost sinusoidal at 10 Mc/sec and became progressively more asymmetrical with increase in frequency. At 15.8 Mc/sec, the width of the zone of positive α was reduced to 120° , while at 22.2 Mc/sec it was only about 60° . These results are consistent with the idea of emission at frequencies near the gyrofrequency from localized regions in the field of an inclined and eccentric dipole, the directions of emission being approximately tangent to the field lines. The results indicate that the field-source and field-sink poles were apparently at longitudes of about 220° and 40° , respectively, in 1963, and that the dipole is displaced from the center of the planet toward the field-source pole. Corresponding features of the curves of α versus longitude occurred 10° to 20° later in 1963 than in 1962, substantiating the need for a slight correction in the System III rotation period.

Random-Hole Mask Image Spoiler for Schmidt Camera. ROBERT J. CHAMBERS, *Frank P. Brackett Observatory*.—The results are described of use of an experimental image spoiler mask placed in front of the corrector plate of the 48-in. Schmidt at Mount Palomar.

The mask incorporated 400 holes $\frac{1}{2}$ -in. in diameter and 100 holes 1 in. in diameter. Their positions were computed by a random number generation program. The number of holes that can be placed randomly on a mask is limited by overlapping. Photographic speed of the system is reduced because of increased image size for the light gathered. The resulting images had intensity distributions nearly that given by the superposition of the theoretical intensity patterns of 1-in. and $\frac{1}{2}$ -in. holes. The improved accuracy of photometric interpolation of the larger images is paid for in decreased speed and limiting magnitude. In future applications, enlargement of images may be controlled by selection of mask apertures, but should be done with regard to crowding of star images as well as desired image growth.