$2GM/c^2$. Configurations with $M > 10^6 M_\odot$ and $\alpha \gtrsim 10^{-3}$ are dynamically unstable as a consequence only of general relativity.

Helium in the Interstellar Medium. J. W. TRURAN, C. J. HANSEN, AND A. G. W. CAMERON, *Yale University and Institute for Space Studies, Goddard Space Flight Center, NASA*.—The usual assumption of modern theories of nucleosynthesis is that the initial composition of the galaxy was pure hydrogen. The large content of helium in the sun and recently formed stars then constitutes a problem; it is usually argued that an exceptionally large amount of stellar processing of nuclear fuels is required in the early history of the galaxy to form the observed abundance of helium. But this would require that the helium be produced in massive stars, which is not possible since in the late stages of evolution of massive stars only a small amount of helium is present between the hydrogen-and-helium-burning shells. Modern studies of stellar evolution suggest that more efficient ejection of helium into the interstellar medium comes from stars of roughly solar mass. Numerical integrations have been carried out to follow changes in composition in the interstellar medium as a result of star formation, nucleosynthesis, and mass ejection, for a variety of assumptions about the rate of star formation. These studies show that the observed helium content of the sun and of O and B stars can be obtained only if the galaxy is quite old ($\sim 2 \times 10^{10}$ yr). The ratios of abundance of uranium and thorium isotopes are consistent with those observed in the earth and meteorites for a wide range of helium contents of the interstellar medium, including the solar value. Hence these radioactivities give little information about cosmochronology.

Zeta Tauri—a Star in Eruption. ANNE B. UNDERHILL, *Sonnenborgh Observatory, University of Utrecht*.—About 1960 it was noticed that the shell spectrum of Zeta Tauri had increased in complexity. Observations obtained in France by R. Herman and in Italy by M. Hack confirmed that material was streaming from the star. At this time the absorption line profiles appear to have been symmetric with shortward displacements. There were also some longward displaced components. This paper draws attention to further changes in the shell spectrum seen on a spectrogram obtained 19 October 1964. This spectrogram covers the red–yellow region at 15 Å/mm.

The Na D lines and the Si II lines at 6347 Å and 6371 Å are strong in absorption, the profiles being sharp on the longward side but winged on the shortward side. This asymmetry indicates that a consider-

able range of velocities exists in the absorbing shell. This is the first time that such asymmetrical lines have been reported for Zeta Tauri. Strong, shortward-displaced symmetrical lines of He I are visible at 5876 Å and at 6678 Å. The H α emission is about twice as strong as in 1953 (Underhill, A. B., *Monthly Notices Roy. Astron. Soc.* 113, 477, 1953), the longward side of the emission being the stronger component. The absorption core is asymmetrical and displaced shortward about 40 km/sec from the position expected according to the well-established binary orbit for Zeta Tauri. A similar displacement is found for the He I lines and for some Fe II lines.

The simultaneous great strength of the Na D lines and of the Si II lines, which comes from a level of 8.12 V excitation, pose an interesting problem in interpretation. The shortward displacement of the D lines is about 13 km/sec; the Si II lines are displaced shortward by about 30 km/sec from the velocity expected owing to binary motion.

Proper Motions of Stars in the Galactic Cluster M 37. A. R. UPGREN, *U. S. Naval Observatory*.—The proper motions of about 600 stars in the region surrounding the galactic cluster M 37 have been determined from photographic plates taken with the 40-in. refractor of the Yerkes Observatory. Eleven plates covering the interval 1903–1964 were measured.

No systematic differences are present between the proper motions of 47 member stars inhabiting the cluster main sequence with intrinsic colors indicating spectral types of A0 to A5 and those of 27 members of the cluster's giant branch. The motions of the stars belonging to both of these groups indicate a contraction of the cluster along the direction of galactic rotation. This contraction appears to be consistent with the proper motions given by Meurers and Schwarz (*Bonn. Ver.* 53, 1960) and by Jefferys (*Astron. J.* 67, 532, 1962).

Recent Planetary Observations at Wavelengths Near 1 cm. W. J. WELCH AND D. D. THORNTON, *Radio Astronomy Laboratory and Space Sciences Laboratory*.—Following the inferior conjunction of Venus in June 1964, the microwave emission spectrum of the planet was measured at seven wavelengths in the range 8–16 mm, the measurements made with the 10 ft reflector of the Space Sciences Laboratory in Berkeley. The following brightness temperatures were obtained: $560^\circ \pm 51^\circ \text{K}$ at 16.53 cm, $595^\circ \pm 50^\circ \text{K}$ at 14.56 mm, $530^\circ \pm 45^\circ \text{K}$ at 13.50 mm, $451^\circ \pm 53^\circ \text{K}$ at 12.45 mm, $495^\circ \pm 50^\circ$ at 11.58 mm, $412^\circ \pm 55^\circ \text{K}$ at 9.71 mm, $390^\circ \pm 45^\circ \text{K}$ at 8.35 mm. No significant absorption features are evident. In