axy with 50 pc diameter and $M = -18$ brightness, one obtains that in the 21 cases of quasars with known redshift, the deflector galaxy is with at least 2 mag fainter than the "quasar." As an example we give the mass and the apparent magnitude of the deflector galaxy for 3C9 ($m = 18.21$; $z = 2.012$) in three model universes.

	$\log(\mathfrak{M}/\mathfrak{M}_\odot)$	m_D	M	$\Delta n/\Delta z$	α
Bondi-Gold	13.55	22.62	-28.59	1.7×10^{-3}	59.0
Evolut. ($q_0 = +1$)	12.11	23.42	-26.20	0.05	14.2
Fib	10.57	27.13	-24.43	13	1.2

The last three columns list the absolute magnitude M of 3C9; the probability to observe a "quasar" with redshift between $z = 1.5$ and $z = 2.5$ having equal or higher intrinsic luminosity than 3C9; in one-quarter region of the sky; and the apparent diameter α of the image in seconds of an arc. We can see that the observed number (four so far) as well as the starlike appearance of quasars favor the Fib model. Should our universe correspond to this latter model, the search for objects with large cosmological redshifts could be facilitated by the detecting devices described in *Z. Physik* 176, 435, 1963.

Cosmic X-Ray Sources—Galactic and Extragalactic. E. T. BYRAM, T. A. CHUBB, AND H. FRIEDMAN, *The E. O. Hulburt Center for Space Research*.—An Aerobee rocket survey of the night sky for x-ray sources in April 1965 revealed fluxes from discrete sources in the directions of Cygnus A, M87, and Cassiopeia A with positional uncertainties of 1.5 deg. The intensity of Cyg XR-1 was about one-fourth as great as had been observed in June 1964.

Properties of Vibrating Neutron Stars. A. G. W. CAMERON, *Goddard Institute for Space Studies*.—The range of uncertainty in the upper limit to the stable mass of a neutron star overlaps the uncertainty in the mass of a remnant left in current hydrodynamical studies of supernova explosions. Thus neutron stars remain legitimate objects for theoretical astrophysical investigation. The supernova implosion may store upwards of 10^{50} ergs as thermal, vibrational, or rotational energy. In the age range 10^3 – 10^5 yr, surface temperature consistent with photon and neutrino removal of thermal energy are in the range 1 – 2×10^6 °K, too small for identification with x-ray sources like the Crab Nebula. The principal internal damping mechanism for the vibrational energy is the URCA neutrino process. After 10^3 yr the amount of vibrational energy remaining is model-dependent but probably in the range 10^{47} – 10^{49} ergs.

This is a sufficient energy source for the Crab Nebula if suitable mechanisms exist to transfer vibrational energy to the surrounding environment. Such mechanisms include shock heating of a neutron star corona and the subsequent formation of a stellar wind which will draw the external magnetic field into a radial configuration. The vibrationally induced hydromagnetic waves in this field can accelerate electrons and ions of photospheric composition. The hot coronal plasma and the energetic electrons can each form an extended x-ray emitting region. The ion acceleration forms an appealing mechanism to account for the large excess of medium and heavy nuclei in the cosmic rays.

Operation Moon-Blink. WINIFRED SAWTELL CAMERON, *Laboratory for Theoretical Studies, Goddard Space Flight Center*.—Operation Moon-Blink is a project to develop, fabricate, and test James Edson's concept of using a blink technique to detect lunar color phenomena with a rotating two-color filter wheel. The filtered transmission is passed onto an image-converter tube whose face may be viewed or photographed directly. The whole unit is attached to the telescope at its focus. A color phenomenon is seen as a black, blinking area. In progress is the development of a multipurpose detector, composed of a camera, polarimeter, spectrograph, and infrared detector, each to be successively rotated into the focal plane during an event. When completed, it will give the first multifaceted assault on the nature of these phenomena.

Ten Moon-Blinks have been distributed across the nation to amateur groups, headed by a professional, who maintain a lunar surveillance. The device can detect a red or blue color event whose brightness is as little as 2% above the lunar background.

In conjunction with the Moon-Blink Network, a Confirmation Network of 12 professional and 4 amateur observatories has been established along the east coast. They have agreed to turn their telescopes to the moon, when feasible, if alerted by a conference call that an event is in progress. Standard forms, giving observing details are then sent to W. S. Cameron at Goddard Space Flight Center.

I have received many other reports of transient phenomena, details of which are presented in tabular form. The last Moon-Blink observation was made on 15 November 1965. It marked the first completely successful Confirmation Network alert; two visual partial confirmations were received. Also, photographs and radio data were obtained. These observations are discussed in some detail. Other groups on the Moon-Blink Network have also been successful in photographing transitory phenomena in the Arcturus region.