Velocities of the Solar Wind Plasma from the Orientations of Ionic Comet Tails. JOHN C. BRANDT, MICHAEL J. S. BELTON, AND MARVIN W. STEPHENS, *Kitt Peak National Observatory*.—This paper presents the preliminary results concerning the velocity of the solar wind plasma as deduced from the observations of Type I (ion) tails contained in the comet catalogue of Belton and Brandt. The analysis is in the spirit of the early work by Biermann where the orientations are interpreted on the basis of dynamical aberration. The major conclusions are as follows:

(i) The solar wind has a fairly sharp velocity minimum at 175 ± 25 km/sec.

(ii) The solar wind velocity is correlated with geomagnetic activity in a manner similar to that found by the Mariner II deep space probe and by studies of travel times for geomagnetic storms.

(iii) Evidence is found for a mean tangential component of the solar wind near the earth of 10 km/sec directed in the sense of the solar rotation; the value of the tangential velocity is apparently lower for quiet conditions and higher for disturbed conditions in the interplanetary medium.

(iv) The sample of Type I comet tails (both direct and retrograde) is not uniformly distributed in space and time, and care must be exercised in drawing conclusions from the sample presently available; such bias may (at least in part) account for the curious dependence of the solar wind velocity on heliographic latitude found by Pflug where the *minimum* velocity is found in the sunspot zones.

An Astrometric Reflector for Canada. ELMAR BROSTERHUS, *Dominion Observatory*.—While trigonometric methods are inoperative for the remote stars of high luminosity, an almost unlimited amount of astrometric data could be accumulated by extending photographic astrometry to very faint limiting magnitudes and concentrating on the great number of stars which populate the faint end of the luminosity function in the sun's vicinity.

The discovery of increasingly faint white and red dwarfs in the solar neighborhood indicates that, in spite of the small distance range within which trigonometric methods produce reliable results, future astrometric telescopes will have to be as powerful as possible and yet maintain the highest level of optical performance and mechanical stability.

The Dominion Observatory plans to build an astrometric reflector at the site of the 150-in. QE II on Mount Kobau, B. C. It is hoped that financial support will be such that the order for the main mirror blank can be placed five years from now. In the meantime comprehensive design studies will be undertaken. The experience gained by K. Aa. Strand and his group in the construction and operation of

the 61-in. astrometric reflector of the U. S. Naval Observatory will be most valuable.

No decisions concerning the optical and mechanical specifications of a Canadian astrometric reflector have yet been made except that the aperture shall be not less than 60 in.

The Formation of the Iron Equilibrium Peak.

A. G. W. CAMERON AND J. W. TRURAN, *Institute for Space Studies*.—At temperatures in the vicinity of 10^9 °K and above, it is expected that energy losses from stellar interiors by neutrino–antineutrino pair emission will become enormous and dominate the last presupernova contraction stages of a star. The last effective energy generation stage is oxygen-burning; following oxygen destruction in a stellar core it has been shown by Chiu that the core evolves to high density at relatively low temperature. Hence the iron equilibrium peak is not made before the supernova implosion. We have examined the types of thermonuclear reactions that can occur in the supernova shock wave which ejects the stellar envelope, in which temperatures $\sim 5 \times 10^9$ °K may occur for $\sim 10^{-1}$ sec. The calculations are performed using a network of nuclei connected to their neighbors by absorption or emission of neutrons, protons, α particles, and photons. Cross sections for these reactions with probable errors of a factor 2 have been calculated based on newly developed expressions for general nuclear parameters. The results show that iron peak elements will be produced by the supernova shock wave, but the iron peak composition observed in nature is not produced unless electron captures have occurred in the presupernova material. We believe that the most important such captures are those which convert S^{32} (a principal product of oxygen-burning) to Si^{32} in the inner parts of the presupernova at densities near 10^9 g/cm³.

Rotation Curves of the Extragalactic Nebulae NGC 3310, NGC 4258, and NGC 4736.

GUIDO CHINCARINI AND MERLE F. WALKER, *Osservatorio Astrofisico, Asiago, Italy, and Lick Observatory*.—During the past several years, spectroscopic observations of a number of galaxies have been obtained by one of us (MFW) using the 120-in. reflector and the prime-focus spectrograph for the outer parts of the nebulae, and the Lallemand electronic camera and coude spectrograph for observations of the nuclear regions. Attention has been directed primarily to those objects in which motions in the nucleus and/or noncircular motions might be detected. From this material, three galaxies are discussed. The particular features of interest shown by these objects are as follows:

NGC 3310: Type Sb. A well-defined, symmetric