common by SAL and Boggess at essentially identical wavelengths agreed to within 0^m25 , which probably represents a realistic value for the best absolute observational accuracy obtainable by these means.

(2) Strong absorption lines can seriously affect the comparison of observed fluxes with model stellar atmospheres in which lines are not taken into account, especially shortward of 1500 Å. For example, at $\lambda 1314$ and $\lambda 1115$ line blanketing reduces the observed fluxes from B-type stars by about 0^m6 – 0^m7 .

(3) A value for the interstellar reddening at $\lambda 1115$ was derived and found to fit, within the large errors, on a straight line through points previously determined by Boggess and Borgman at $\lambda 2600$ and $\lambda 2200$ and by Smith at $\lambda 1376$.

(4) The uv observations were compared with model atmospheres chosen to agree with the ($B - V$) color of the star in question, corrected for reddening when necessary. With the exception of the few measurements at $\lambda 1427$ and $\lambda 1115$, all the observations agree with the model predictions to within 0^m5 ; generally the agreement is better than this. Small changes (a few hundred degrees) in the stellar effective temperature scale used (that of Morton and Adams) would improve the agreement, but they were not felt justified.

Bless presented this paper at the IAU Colloquium on Space Spectroscopy held at Northwestern University, 3–5 May 1967.

X-15 Program. Doherty obtained ultraviolet stellar photographs with the X-15 aircraft. Two morning flights from Edwards Air Force Base in August 1966, to an altitude of 75 km, show images of η Aur, α Aur, ρ Aur and α Gem. Interference filters give these photographs an effective wavelength of 2500 Å and a bandwidth at half-maximum response of 300 Å. The intensity of the dayglow was determined from the films and from a photoelectric scanning spectrometer carried on one of the flights. For the same instrumental response and an image size of one square minute of arc, he finds that a B3 star with visual magnitude 8.0 should equal the brightness of the dayglow at this altitude. The fluxes of η Aur (B3V) and ρ Aur (B5V) agree with the predictions of model atmospheres within the error of measurement. α Aur (G8III) is 2.4 mag fainter than η Aur.

Instrumentation and Laboratory Work. Schroeder considered the design of spectrographs capable of a spectral purity of 1–2 Å per millimeter of entrance slit. Study of various types of dispersing systems showed that a combination echelle spectrometer-spectrograph of moderate size was feasible. A laboratory version of this instrument was set up and performed as expected. The echelle used in this design has 30 lines/mm and is used in the 120th order at $\lambda 5000$ Å. Construction will soon begin on an instrument of this type to be used at the Casse-

grain focus of the 36-in. telescope at Pine Bluff. Some characteristics of the system are (1) beam diameter at the echelle of 74 mm, (2) dispersion of 2.5 Å/mm at 5000 Å with a camera focal length of 0.5 m, (3) spectral purity of 1.25 Å per millimeter of entrance slit, and (4) recording of spectra with a photomultiplier, photographic film, or electronic camera. The details of this design are in a paper which has been submitted for publication in *Applied Optics*.

Fairchild investigated the possibility of using radiation from a storage ring fed by the University of Wisconsin 280-MeV electron synchrotron as an absolute calibration standard in the 1000–2000 Å region. The technique seems to be feasible and attempts at calibration will soon begin.

A small telescope has been placed under computer control and has been operated since late last year. It utilizes an 8-in. filter photometer module developed under the OAO project. This module has remotely controllable filter selection and amplifier gain. The digital computer used is a PDP-8. Stepping motors and shaft position encoders together with an interface provide attitude control and feedback. This system was designed to determine extinction coefficients in four wavelength bands by automatically observing a preselected group of stars throughout the night. There are sensors to detect the presence of rain, morning and evening twilight, and clouds. The telescope housing may be opened or closed and electronics may be turned on or off as circumstances require, all automatically.

DONALD E. OSTERBROCK, *Director*

University of Washington, Seattle, Washington

I. FACULTY AND GRADUATE STUDENTS

The Department of Astronomy now consists of Professors Bardeen, Böhm, Böhm-Vitense, Hodge, Jacobsen, and Wallerstein. Dr. R. Michie was Visiting Associate Professor for one month in May. Six graduate students are enrolled, two of whom have started thesis research. L. Tomley is studying the atmospheres and chemical composition of three hot subdwarfs from spectrograms secured at the Kitt Peak 84-in. coude. T. Greene is studying the CN, C₂, and O I features in several K giants from 2 Å/mm spectrograms taken by Dr. Wallerstein at Lick. With the help of model atmospheres he is deriving the abundances of C, N, and O.

A site survey in eastern Washington has been completed. Observations with automatic sky cameras, visual observations by fire lookouts, and seeing observations with the 16-in. telescope and a portable 6-in. telescope have been combined and evaluated. It appears that Manashtash Ridge near Ellensburg has the best observing conditions in the

state and is also closest to the University of any potential site east of the Cascades.

II. RATTLESNAKE RIDGE OBSERVATORY

The observatory on Rattlesnake Ridge near Hanford, Washington, currently has a Boller and Chivens 16-in. reflector equipped with a photoelectric photometer. Photocells with the S-20 and S-1 surfaces are used for photometry from 3000 to 10 000 Å. Observations on a *UBVr* system (Sandage and Smith 1963) are being used to establish the correlation between the color difference $(B - V) - (V - r)$ and metal abundance in G8-K2 giants and dwarfs. Observations of the variable stars P Cyg, CH Cyg, Nova Del 1967, and SS Cyg are being carried out. Sco XR-1 was observed by E. Mannery in a cooperative effort with Dr. H. Mark of Berkeley to search simultaneously for optical and x-ray variations. Unfortunately the object was quiescent on the night of observation. The 16-in. telescope has also been used by Dr. G. Morris of Boeing Scientific Research Laboratory to search for high-frequency plasma oscillations in stellar atmospheres.

III. STAFF RESEARCH

Solar System—Meteoritic Dust Research. Dr. Hodge and D. Brownlee, predoctoral research associate, flew guest experiments on the S-12 dust collection devices flown on Gemini 10 and Gemini 12 satellites and prepared collection surfaces for the Venus Fly Trap rocket probe flown into the Leonid meteor shower. Because of loss of the Gemini 10 experiment by the astronaut and failure of the Venus Fly Trap rocket to open, only the Gemini 12 experiments have been successful so far.

The Gemini 12 samples have been investigated to study the various types of contamination present, using two control slides, and to detect and study craters. Contaminating particles were counted and identified to evaluate the problems involved in future collections of dust, though the present experiments were designed to be usable only in detecting cratering particles. The total number density of contamination particles is surprisingly large. The analysis indicates that many of these reached the surfaces during flight. No craters larger than $2\ \mu$ in diameter were found, and the resulting implied upper limit to the space density of particles is several orders of magnitude lower than densities predicted from early satellite and rocket results, but is consistent with recent satellite penetration data.

For nondestructive collection of particles, Hodge and Brownlee have designed and built a balloon experiment based on a new principle that is expected to have orders of magnitude better contamination control than previous methods. The first flight, to 120 000 ft, is planned for December 1967.

They have built up a high-specification ultra-clean-room facility, in our Meteoritic Dust Laboratory, for particle analysis, payload cleaning, and collection surface preparation that will enable them to maintain the highest possible level of laboratory contamination control.

Stellar Astronomy. Dr. Wallerstein has completed a search for lithium in stars of types F and G of luminosity class Ib and in classical Cepheids. Of 48 stars observed, not a single one shows a lithium line. The data are being combined with a similar search for beryllium by Dr. P. Conti of the Lick Observatory. The absence of lithium in these stars indicates either that all of them have diluted their surface lithium by mixing with lithium-poor material or that massive stars never produce a sufficient amount of lithium on their surfaces to be detectable spectroscopically.

Wallerstein has completed an analysis of 16 Å/mm Lick spectrograms of RU Cam taken in January 1965 when the star was nearly constant in light. A curve of growth analysis of the atomic lines using ϵ Vir as a comparison star yields a normal ratio of iron to hydrogen and an electron pressure that is lower than that in ϵ Vir by a factor of 10. A discussion of the abundances leads to the conclusion that about one-half of the material presently in the atmosphere of RU Cam has been processed through partial helium burning and subsequent carbon cycling before being mixed to the surface. A discussion of the mass and absolute magnitude leads to limits of 0.5 and $2.0M_{\odot}$ for the mass and -2.0 to -3.5 for the absolute visual magnitude.

Tomley and Wallerstein have obtained 15 spectra of HD 34452 for an analysis of the atmosphere and composition of this silicon-type spectrum variable. Similar spectra of 108 Aqr will be taken in the autumn of 1967.

X-Ray Astronomy. Dr. Wallerstein has observed the interstellar *H* and *K* lines in the star associated with Sco XR-1 at 18 Å/mm dispersion. They are stronger than the interstellar lines in any nearby star and set a lower limit to the distance to the x-ray star of 270 pc.

Extragalactic Research. Using three photoelectric sequences obtained at the Mt. Stromlo Observatory, Dr. Hodge, with F. W. Wright of the Smithsonian, has studied the variable stars of a section of the Large Magellanic Cloud centered at $\alpha = 5^h 12^m$, $\delta = -67^{\circ}5$. Measures in two colors, *B* and *V*, have been obtained on a series of 60 ADH Schmidt plates and measures in the blue were made on approximately 400 Harvard Bruce plates for each variable star. The period-luminosity relation for classical Cepheids has the form

$$\langle B \rangle = 17.94 - 2.38 \log P.$$

The derived slope lies about midway between the value published for the SMC by Arp and that published for the LMC by Woolley *et al.* It is believed that sampling problems for the longest-period Cepheids are a serious problem. Also studied are an R Corona Borealis variable, two W Virginis variables of changing period, two long-period variables, and two eclipsing binaries.

In order to understand better the pattern of evolution of galaxies, Hodge and Wills have attempted to estimate ages for most of the clusters of the LMC. So far, they have completed measures of the brightness of the brightest main sequence stars of 1300 LMC clusters, using photographic transfer techniques. They plan to use these data to estimate cluster ages and to explore the spatial pattern of cluster formation at various times in the LMC past.

Dr. Hodge found that the irregular galaxy G.R.8 is an extremely small galaxy, possibly close enough to be considered a local group member. Its mean dimension is on the order of only 100 pc. Photoelectric measurements show that it is considerably bluer than any other measured (non-quasi-stellar) galaxy, and that its color-magnitude diagram is like that of a very young, not very populous open cluster. Its luminosity function contains only 100 stars in the top five magnitudes, and no H II regions have been detected. Its structure, as revealed by isophotes constructed from photoelectric and photographic photometry, is like that of other irregular galaxies; an approximately exponential decrease of brightness characterizes the outer, unresolved portions.

Hodge and Hitchcock, of San Francisco State College, studied the irregular galaxy NGC 4449, using photographs in V , B , and $H\alpha$, with photoelectric surface photometry, for the purpose of studying its structure and content. The distance to NGC 4449, determined from the diameters of the largest of the H II regions detected, is 3.3 ± 0.2 Mpc. Its mean dimension is 4.5 kpc and its absolute magnitude is $M_V = -18.0$. Isophotes in two colors show the dominance of a large number of resolved features. Of these, at least 81 are H II regions and the remainder are globular clusters, associations and bright stars ($M_V \approx -6$). Resolved features are virtually absent from one quadrant. In an attempt to determine the underlying structure of the galaxy, they plotted a set of isophotes from tracings from which the resolved features that stand alone have been removed. The result shows that the older star population is arranged roughly rectangularly, with one corner anomalously rounded.

Hodge began a large-scale survey of galaxy structure using isophotometric diagrams made from large-reflector and Schmidt plates. So far isophotes are complete, but not yet completely calibrated, for 43 galaxies.

Hodge has begun study of the spatial distribution of H II regions in 25 galaxies. He finds that the H II region surface density varies in a characteristic way with radius for each galaxy type. The most surprising result is that the H II distributions are *very* different from the H I distributions found by M. S. Roberts for the galaxies that they have studied in common. Also, he has begun an investigation, using several nearby galaxies, of the across-arm distribution of luminosity, color, and H II regions, in an attempt to test various theoretical models of spiral arms.

Hodge published a special study of H II regions in Irregular II (I0) galaxies, showing that most of these galaxies are not similar to M82, contrary to what might have been expected. However, several of them were found to be extremely peculiar in many respects and further photometric investigations have been begun. Wallerstein and Welch began a spectrographic study of the most remarkable of these, NGC 5253.

General Relativistic Astrophysics. Dr. Bardeen is continuing work on the properties of spherically symmetric gravitational collapse inside the gravitational radius and on the formation of singularities in the space-time manifold. Certain solutions of the Einstein-Maxwell equations for the collapse of charged dust have the property that even though collapse proceeds inside a trapped surface in the sense of Penrose the density of the dust remains finite for an infinite proper time. The only singularity is in the exterior Reissner-Nordstrom metric at a finite spatial distance from the surface of the dust cloud and in a sphere of zero area at which the electric field and some components of the curvature tensor are infinite. The dust cloud collapses from an asymptotically flat universe, for a while is inside a "bubble" of finite volume, and then expands into an asymptotically flat region of space-time in the absolute future of the first such region. Although these solutions are too artificial to have direct application in astrophysics, they do provide a specific model for the anticollapse which has been proposed as an energy source for quasars. To an observer in the universe into which the dust cloud is expanding the light from the surface of the dust cloud first becomes visible with a redshift decreasing exponentially with time, for a dust cloud with mass M with a time constant the order of GM/c^3 . Eventually the redshift will change to the blueshift ordinarily expected from an expanding object.

These astrophysical applications should not be taken too seriously because quantum-mechanical effects, particularly pair formation in the infinitely strong electric field at the singularity in the exterior metric, may drastically alter the above classical picture based on an analytic extension of the exterior metric.