

GAMMA-RAY OBSERVATIONS OF THE GALACTIC CENTER AND SOME POSSIBLE POINT SOURCES

C. E. FICHEL, R. C. HARTMAN, D. A. KNIFFEN, AND M. SOMMER*

NASA/Goddard Space Flight Center, Greenbelt, Maryland 20771

Received 1971 June 1; revised 1971 August 19

ABSTRACT

A 0.5×0.5 m digitized spark-chamber γ -ray telescope was flown on three balloon flights to look at the galactic center region, Virgo, and the Crab Nebula. An excess flux above atmospheric background of over 4 standard deviations was found for γ -rays exceeding 100 MeV coming from the galactic center region, corresponding to a line intensity of $(2.0 \pm 0.6) \times 10^{-4} \gamma (\text{cm}^2 \text{ rad s})^{-1}$, but there was no statistically significant excess in the 50–100-MeV interval, with a 95 percent confidence limit for the ratio $[J(50\text{--}100 \text{ MeV})/J(>100 \text{ MeV})]$ being 0.50. As a result, there is only a 6 percent chance that Compton or synchrotron radiation from electrons with a power-law spectrum having an exponent of 2.6 could make as much as a 50 percent contribution to the γ -radiation in this energy range. No positive evidence was found for radiation from any of the sources M87, 3C 273, or the Crab Nebula, and 95 percent confidence upper limits for the γ -ray flux from these objects were set at 1.0×10^{-5} , 1.0×10^{-6} , and $6 \times 10^{-5} (\text{cm}^2 \text{ s})^{-1}$, respectively. The M87 limit eliminates the possibility that the power-law spectrum observed at X-ray energies extends unchanged to the high-energy γ -ray region.

I. INTRODUCTION

In spite of the considerable effort that has been extended during the last decade to detect high-energy (>20 MeV) celestial γ -rays, relatively few positive results have been obtained, although significant upper limits on the flux from several possible discrete sources have been set by the use of data from balloon experiments. The most positive evidence for an extraterrestrial source of high-energy γ -rays has come from the counter-telescope experiment of Clark, Garmire, and Kraushaar (1968) flown on OSO-3. These authors have presented convincing evidence for the detection of γ -radiation from the galactic plane, with the greatest intensity coming from the region of the galactic center.

Both in an attempt to measure the intensity and energy spectra of galactic γ -rays above 25 MeV from balloon altitudes and with the goal of developing a satellite-qualifiable γ -ray telescope, a large digitized spark-chamber γ -ray detector was built and flown on high-altitude balloons. This detector, whose active spark-chamber area is about $2.5 \times 10^3 \text{ cm}^2$, is about 10 times the size of the original Goddard spark chamber first flown in 1966. Preliminary results from one of the flights with the larger telescope, which confirmed the existence of radiation from the galactic center, were presented earlier (Kniffen and Fichtel 1970). Here, the experiment will be described in more detail together with the final results from the balloon flights made thus far.

II. DESCRIPTION OF THE GAMMA-RAY TELESCOPE

A schematic diagram of the spark-chamber γ -ray telescope is shown in Figure 1. The active parts of the detector are the 0.5×0.5 m digitized spark-chambers and the triggering counter system. The spark-chamber stack is assembled from a total of 32 modules, 11 of them being placed above the central scintillator level. Each module consists of two orthogonal sets of 400 parallel wires forming two wire grids with 4 mm vertical spacing. Stainless steel plates with a thickness of 0.03 radiation lengths between

* NAS-NRC Postdoctoral Resident Research Associate. Present address: Max-Planck Institut für Extraterrestrische Physik, München, Germany.




FIG. 1.—Schematic diagram of the 0.5×0.5 m digitized spark-chamber γ -ray telescope

the upper wire modules provide conversion of γ -rays into pairs. Plates of 0.02 radiation lengths between the lower modules are used to obtain information on the energy of the individual positron and negatron from Coulomb scattering. Sparks will occur along the paths of the electrons when a high-voltage pulse is applied to the wire grid shortly after detection of the electron pair by the counter telescope. The resultant spark current propagates along the orthogonal wires, setting ferrite cores at the end of the wires. In this manner, x - and y -coordinates are obtained for each electron track at every module or z -level. The spark coordinates are described by a number of 16-bit words which are transmitted by telemetry to the ground station within a total time of about 0.3 s for an average event.

The spark-chamber triggering logic is chosen to accept γ -rays and to reject other events to a high degree. A coincidence signal is required from one of the nine charged-particle telescopes. Each of these is formed by a $16.5 \times 16.5 \times 0.48$ cm plastic scintillator and a $16.5 \times 16.5 \times 5$ cm Plexiglas Cerenkov counter. The spark chamber is triggered when a neutral particle is converted into one or more downward-moving relativistic charged particles actuating one or more of the nine charged-particle telescopes. Primary charged particles are rejected by anticoincidence signals produced in the surrounding scintillator dome. The dome is a highly polished single-piece casting of plastic scintillator measuring 1.2 m inside diameter, 0.8 m in height, and 1.6 cm in thickness. It is viewed by 18 photomultiplier tubes spaced evenly around the lower edge.

A more detailed description of the 0.5×0.5 m digitized spark-chamber γ -ray telescope is given by Ross *et al.* (1969).

III. DATA REDUCTION, ANALYSIS, AND DETECTOR CALIBRATION

Data including a variety of housekeeping parameters and information on which of the magnetic cores in the spark chamber have been set in each event are telemetered from the balloon gondola and recorded on magnetic tape together with time signals accurate to 2 ms. The housekeeping information includes three-axis magnetometer readings which, together with time and balloon position, provide the basis for determining the experiment aspect, individual counter and coincidence rates, experiment dead time, and engineering parameters such as temperatures, pressures, voltages, and currents.

The spark-chamber data reduction includes identification of γ -ray events, estimation of the γ -ray energy from the scattering of the electron pair in the steel plates, and determination of the celestial arrival direction, utilizing the magnetometer data to obtain the instrument orientation. The general approach and details for the calculation of the energy and arrival direction of individual γ -rays have been described previously, together with the method of determining flux values (Fichtel *et al.* 1969); therefore, they will not be repeated here. In the interim, however, the computer programs have been developed to the extent that γ -ray identification and electron-track recognition are largely automatic, and the human interface time has been greatly reduced. A minimum interface for the purpose of verifying that the program is functioning properly is maintained.

The procedures for automatic recognition will now be summarized. The automatic analysis program rejects any event which (1) has any set cores in the top deck (representing generally either a γ -ray which has converted in the shell of the pressure vessel under the anticoincidence dome or a charged particle missed by the anticoincidence system), (2) has a set core within two wires of the edge of any of the top 10 modules (and therefore would usually correspond to a γ -ray which interacted in the spark-chamber walls), or (3) has less than 15 cores set in either orthogonal view (indicating no event, usually a false trigger).

The program then begins the search for γ -ray events by averaging groups of two or more adjacent set cores to obtain points which will represent the basis for the "picture" of the events in each of the two orthogonal views (x - z and y - z). The program then scans downward from the top of the spark chamber until it finds the occurrence of three decks containing points within a five-deck span. Within those three decks, it forms all possible triads with one point in each deck. For each triad it forms the second difference $\beta_i = X_i - 2X_{i+1} + X_{i+2}$. Triads which lead to a β_i -value greater than a fixed limit are rejected under the assumption that they cannot represent a segment of an electron track of interest. After all possible triads have been examined, the unmatched points are saved and the next deck with points is found. New triads are found, and the procedure iterates. Track segments containing the most points are examined further to ascertain their interrelationship in an attempt to form full tracks from the segments.

For a γ -ray, two basic forms occur depending on how fast the electrons separate, the "inverted Y" and the "inverted V." In the case of the Y configuration, the stem is considered to be two unresolved electrons and is associated with each of the two separated tracks. Some events appear to contain only a single-track segment. These are classified as either low-energy electrons or an unresolved pair (high energy, straight track), depending on the amount of Coulomb scattering. Experience has shown that the single tracks do indeed fall in these two groups with essentially no tracks having intermediate Coulomb scattering values.

There remain some spark-chamber events which the computer is unable to handle satisfactorily. The following are examples of situations which cause the program to abandon an event: (1) less than ~ 45 percent of the total X or Y points being contained in the two longest segments; (2) less than 13 decks containing points; (3) no acceptable triads; (4) no acceptable tracks (≥ 5 points), (5) too many segments (≥ 15 , which cannot be formed into longer tracks, (6) Ambiguous relationship between the two main tracks, (7) two main tracks not joined.

Events which are abandoned by the computer are then subjected to visual analysis in the following manner. The digitized "picture" of each of these events is displayed on an automatic graphics display unit where the necessary editing is accomplished by light-pen interaction and the remainder of the analysis is completed by machine, if it is an acceptable event. If it is evaluated to be a false event, it is rejected.

Although most of the results presented here are from the computer-analyzed data, extensive checks have been made by reanalyzing samples of the data visually. In its present stage of development, the computer analysis produces results of comparable or even slightly superior quality, when it is applied to high-quality data. When the efficiency of the spark chamber is less than optimal, the computer analysis is less dependable, particularly for γ -ray energies below about 70 MeV, where the electrons undergo considerable scattering. For that reason, all of the results presented here for γ -ray energies below 100 MeV are based on either visual analysis of the events or computer analysis checked by visual examination of the pictures.

Two separate approaches have been used in the calibration of the γ -ray telescope being discussed here. The first is a direct one using the γ -ray facility associated with the synchrotron at the National Bureau of Standards. Here it is possible to study the detection efficiency of the instrument as a function of energy, angle, and position of arrival in the detector. The other approach to calibration is a comparison with the intensity and energy spectrum of the atmospheric secondary γ -radiation measured at the same location by the smaller GSFC γ -ray telescope flown previously (Ehrmann *et al.* 1967; Fichtel, Kniffen, and Ögelman 1969). This smaller telescope has also been calibrated at an accelerator. The intensity and energy spectra of atmospheric γ -rays measured by these two instruments agree within the quoted uncertainties.

With regard to the calibration at the National Bureau of Standards, a bremsstrahlung beam from the 170-MeV electron synchrotron is directed into a tantalum target where

it is converted into electron pairs. The electrons are then doubly momentum analyzed to obtain a clean monoenergetic beam of electrons with selectable energy in the range 10–150 MeV. The monoenergetic electrons are then passed through a counter telescope consisting of two plastic scintillators, S_1 and S_2 , a lead target, and finally a thick plastic scintillator, $\bar{S}_3$. In this manner, γ -rays are obtained with the signature $S_1 \times S_2 \times \bar{S}_3$. Furthermore, the photons will have only a very narrow energy range, since electrons with any significant residual energy following a bremsstrahlung interaction in the lead target will activate the S_3 counter and inhibit the tag. The photons closely reflect the narrow collimation of the original electron beam, as confirmed by the γ -ray spark-chamber results.

The analysis of individual events from the γ -ray telescope is the same for the machine calibration as it is for the balloon flights, both in terms of the computer programs and in the visual analysis of a small sample for comparison purposes.

IV. THEORETICAL BACKGROUND AND RESULTS

a) *The Galactic Center Region*

It is well known that mesons are produced in the interaction of cosmic rays with interstellar matter, and by far the most common mesons are π -mesons, both charged and neutral. The π^0 decays into two γ -rays, each with about 70 MeV in the rest frame. Many of the other mesons and hyperons also decay into neutral π -mesons. As a result, the great majority of the high-energy γ -rays resulting from nuclear interaction are from π^0 mesons, most of which are formed in the initial interaction.

The energy spectrum of the γ -rays resulting from π^0 decay is quite different from that resulting from most of the other astrophysical processes, such as bremsstrahlung, synchrotron radiation, and the inverse Compton effect. The basis for the relatively unique spectrum is the neutral pion's isotropic decay into two γ -rays of equal energy (~ 70 MeV) in the rest frame. As a result, for a given velocity of the parent π^0 the probability of observing a given energy for the secondary γ -ray is a constant from a minimum energy of $W_0[(1 - \beta)/(1 + \beta)]^{1/2}$ to a maximum energy of $W_0[(1 + \beta)/(1 - \beta)]^{1/2}$, where W_0 is the energy of the γ -ray in the rest frame. Thus, if the observed γ -rays come from neutral pions formed by the interaction between cosmic rays and interstellar matter, a spectrum will be formed which peaks at W_0 since this energy is always included, and extends further on the high-energy side. The exact shape of the spectrum depends on the energy distribution of the cosmic rays and the details of the interaction process; the related calculations have been completed in detail in the literature (e.g., Stecker 1971). The actual intensity of the γ -rays was also calculated, with the most uncertain parameters being the density of hydrogen and the cosmic-ray intensity.

The OSO-3 satellite experiment of Clark *et al.* (1968) and Garmire (1970) has detected a celestial γ -ray flux concentrated in the galactic plane. Their angular resolution ($\pm 15^\circ$) is substantially larger than the structure of the plane, so the detailed structure cannot be seen. It is possible to quote the result as a line intensity and compare it with the result deduced from the 21-cm data integrated over the appropriate solid angle. The value of the galactic line intensity averaged over the observing angle of the OSO-3 experiment is in the range $(2-4) \times 10^{-5} \gamma(\text{cm}^2 \text{ rad s}^{-1})$, in reasonable agreement with expectation except near the galactic center. The intensity from the direction of the galactic center is not immediately explained. Even after normalizing to the rest of the galactic plane, the intensity averaged over a 60° wide region including the galactic center is about 3 times the expected flux. Kniffen and Fichtel (1970) in a preliminary Letter have confirmed the existence and intensity of the flux from the galactic center with a balloon flight.

The explanation of the enhanced γ -radiation might be postulated to be due to an increase in the intensity of the cosmic radiation and interstellar matter near the center of the Galaxy, or it may be due in part to interactions of electrons with matter, magnetic

fields, or photons. The unique energy spectrum of the π^0 decay makes it possible to separate this mechanism from bremsstrahlung, synchrotron radiation, and Compton radiation—all of which reflect the presumed power-law spectrum of the parent electrons. Specifically, if the parent electrons have a spectrum of the form $j_e = CW^{-\alpha}$, the γ -ray spectrum will have the form $j_\gamma \propto W^{-\alpha}$ for bremsstrahlung radiation, and $j_\gamma \sim W^{-(\alpha+1)/2}$ for Compton and synchrotron radiation.

The radiation observed from the galactic center and reported previously (Kniffen and Fichtel 1970) has now been analyzed fully, including information on the energy spectrum. Figure 2 shows the ratio of the observed intensity to the intensity expected for atmospheric background as a function of galactic latitude for intervals of galactic longitudes from -25° to $+20^\circ$. The atmospheric-background normalization was determined from regions away from the galactic plane in the same flight and was found to agree to within better than 5 percent with that determined previously with a similar but smaller γ -ray telescope (Fichtel *et al.* 1969). The excess of γ -rays above atmospheric background was determined for the interval $-6^\circ \leq b^{\text{II}} \leq +6^\circ$ and $-35^\circ \leq b^{\text{II}} \leq +25^\circ$ for the energy intervals 50–100 MeV and greater than 100 MeV. In this solid angle, there were 274 γ -rays whose measured energies were above 100 MeV compared with 210 expected from atmospheric background. Thus, the excess for γ -rays above 100 MeV corresponds to more than 4 standard deviations and leads to a line intensity of $(2.0 \pm 0.6) \times 10^{-4} \gamma (\text{cm}^2 \text{ rad s})^{-1}$. The value is in good agreement with the corrected line intensity of $(1.2 \pm 0.3) \times 10^{-4}$ photons $(\text{cm}^2 \text{ rad sec})^{-1}$ for energies above 100 MeV observed by Clark, Garmire, and Kraushaar (Garmire 1970). Both of these results are inconsistent with the upper limits (95 percent confidence) set by Frye *et al.* (1969) for




FIG. 2.—Ratio of observed line intensity of >100 MeV γ -rays to expected background intensity for $-25^\circ \leq l^{\text{II}} \leq +20^\circ$, plotted as a function of galactic latitude, b^{II} . Lower curve shows data plotted in 2° bin widths, the approximate angular resolution of the detector. Upper curve includes the same data in $\pm 6^\circ$ bins to increase statistics and more clearly show the latitudinal dependence.

energies above 50 MeV of $3 \times 10^{-5} \text{ (cm}^2 \text{ rad sec)}^{-1}$ for a region bounded by $-2^\circ 0' \leq b^{\text{II}} \leq 2^\circ 3'$ and $7 \times 10^{-5} \text{ (cm}^2 \text{ rad sec)}^{-1}$ for region bounded by $-15^\circ \leq b^{\text{II}} \leq 15^\circ$. The l^{II} range is similar to that of this experiment.

There is no statistically significant excess in the 50–100-MeV interval, and the following 95 percent confidence upper limit was obtained:

$$R = [J(50\text{--}100 \text{ MeV})/J(> 100 \text{ MeV})] < 0.50 .$$

For π^0 decay alone, this ratio is expected to be 0.12 (Stecker 1971), consistent with the experimental result. Current estimates of the spectra of cosmic-ray electrons, in the high-energy region which would produce these γ -rays indicate that the spectral index for the electrons is between 2.5 and 2.8 (Anand, Daniel, and Stephens 1970; Agrinier *et al.* 1970; Bleeker *et al.* 1970; Earl, Neely, and Rygg 1970; Fanelow *et al.* 1969; Marar, Freier, and Waddington 1970; Nishimura *et al.* 1970; Rockstroh and Webber 1969). For an electron spectrum with $\alpha = 2.6$, the ratio R would be 2.03 for bremsstrahlung radiation and 0.74 for Compton or synchrotron radiation. The experimental result, therefore, suggests that the radiation from the galactic center region is probably predominantly from π^0 decay. More quantitatively, there is only a 6 percent chance that Compton or synchrotron radiation from electrons with a spectrum having a 2.6 exponent could make as much as a 50 percent contribution to the γ -radiation in this energy range, and the data are in agreement with the radiation being entirely of π^0 -decay origin.

b) Virgo

One of the most surprising results of X-ray astronomy has been the discovery (Byram, Chubb, and Friedman 1966; Friedman and Byram 1967; Bradt *et al.* 1967) of a surprisingly intense extragalactic discrete source of 1–10-keV X-rays in the direction of the Virgo cluster. Recent high-resolution satellite studies (Kellogg *et al.* 1971) have provided convincing evidence that the source coincides with the radio galaxy M87 (Virgo A) and that M84 and the QSS 3C 273 are other likely discrete-source emitters in the direction of Virgo. Measurements of high-energy (40–100 keV) X-rays from Virgo A have also been reported (Haymes *et al.* 1968; Fishman *et al.* 1970), but these measurements lack independent verification and, in fact, conflicting results have been reported (Peterson 1970; McClintock *et al.* 1969; Webber and Reinert 1970). Felten (1970) has summarized the existing observations over the entire electromagnetic spectrum of the “jet” of Virgo and finds that the data over 13 decades of energy are reasonably consistent with a single power-law spectrum given by

$$N(E) = 8.2 \times 10^{-4} E^{-1.75} \text{ photons (cm}^2 \text{ s MeV)}^{-1} .$$

Extending this spectrum 2–3 decades above the highest observed energies gives an expected flux above 30 MeV of about $9 \times 10^{-5} \text{ (cm}^2 \text{ s)}^{-1}$.

It is most probable that the power law in the visible and radio portions of the spectrum is generated by the synchrotron emission of energetic electrons traversing the magnetic fields in the source galaxy. However, until the X-ray spectrum is better known, it is not clear whether this emission represents an extension of the power-law synchrotron emission at lower frequencies, which would create a very puzzling dilemma because of the very short lifetimes of the electrons at these energies, or whether it indicates the presence of additional processes such as the thermal bremsstrahlung suggested by Sartori and Morrison (1967). The resolution of this question depends upon further experimental results, and for this reason we undertook a balloon flight to examine the Virgo region for the emission of $> 30 \text{ MeV}$ γ -radiation.

The detector was flown 1970 October 14 from Palestine, Texas, on a balloon with a volume capacity of 27 million cubic feet. It reached a depth in the atmosphere of 2.5

g cm^{-2} at 8:55 A.M. CDT and remained at float until terminated by command at 5:00 P.M. CDT. The detector was tilted 20° and oriented to point toward the south in order to more efficiently examine M87 which lies at a declination which passes the meridian plane about 20° south of the Palestine latitude. M87 passed overhead at about 12:15 P.M. CDT. At about 2:15 P.M. the detector was commanded level and allowed to examine the sky centered at a declination of about $+31.5^\circ$ corresponding to the latitude of the balloon.

The data from this flight were analyzed by the automatic analysis program, with no human interaction. Since a significant portion of the events below about 70 MeV require human interpretation, the analysis reported here includes only γ -rays with measured energies in excess of 100 MeV.

The background was determined by analyzing all portions of the sky which were examined except for the areas surrounding possible sources. Table 1 summarizes the results obtained from M87 and 3C 273. The 95 percent confidence limits were obtained by the method outlined by Fichtel *et al.* (1969). The M87 limit clearly rules out the possibility that the power-law spectrum extends unaltered to γ -ray energies, as seen in Figure 3. However, the limit places no constraint on the secondary-production model

TABLE 1
DATA SUMMARY OF POSSIBLE DISCRETE SOURCES IN
THE DIRECTION OF M87 AND 3C 273

Variable	M87	3C 273
Number of gammas expected ($6^\circ \times 6^\circ$)..	7.4	5.0
Number of gammas observed..	8.0	3.0
95 percent confidence number limit ..	7.8	4.8
95 percent confidence flux limit ($10^{-5}/\text{cm s}$)	1.0	1.0

NOTE.— γ -ray limits $E_\gamma \geq 100$ MeV.




FIG. 3.—The upper limit to >100 -MeV γ -rays from M87 obtained in this experiment, plotted together with the empirical curve for the jet in M87 obtained by Felten (1970). Dashed curve, extrapolation of the curve above the highest observed energies.

of Felten (1968) which attempts to explain the presence of electrons in the jet as arising from the decay of pions produced in collisions of energetic protons from the core with matter in the jet. The decay of accompanying neutral pions would be expected to produce a γ -ray flux orders of magnitude below the observed limit. (The upper limit to the M87 γ -ray flux reported here is consistent with that reported by Frye and Wang 1969.)

c) Crab Nebula

On 1969 July 15 a balloon flight of the 50×50 cm spark chamber was conducted at Palestine, Texas, to examine the Crab Nebula for possible discrete-source γ -ray emission, either steady or pulsed. Unfortunately, a random electronic component failure prevented a full exposure to this region. No evidence was obtained for discrete-source γ -ray emission in the Crab Nebula, and 95 percent confidence limits of 6×10^{-5} (cm² s)⁻¹ and 8×10^{-5} (cm² s)⁻¹ were obtained for integral fluxes above 50 and 10 MeV, respectively. The arrival times of the γ -rays coming from the direction of the Crab Nebula were tested against the observed optical period for this date (Fazio and Helmken 1971), and no evidence for pulsation was seen. The limited statistics prevent the upper limit for the pulsed mode from being substantially lower than those for the steady mode given above.

We wish to express our appreciation to S. Derdeyn, C. Ehrmann, A. Mascaró, R. Ross, and R. Smith for their outstanding engineering support and to R. Marsh for her long hours of careful work in support of the data analysis. Also, we acknowledge the stimulating discussions that we had with H. Ögelman during the early phases of the experiment. For the three successful flights, we are indebted to the balloon-launching crews of the Australian Department of Supply at Mildura, Victoria, and the NCAR at Palestine, Texas, and to the personnel of the office of Naval Research.

REFERENCES

- Anand, K. C., Daniel, R. R., and Stephens, S. A. 1970, *Acta Phys. Suppl.*, **1**, 229.
 Agrinier, B., Koechlin, Y., Parlier, B., Paul, J., Vasseur, T., Boella, G., Sironi, G., Russo, A., and Scarsi, L. 1970, *Acta Phys. Suppl.*, **1**, 203.
 Bleeker, J. A. M., Burger, J. J., Deerenberg, A. J. M., Hulst, H. C. van de, Scheepmaker, A., Swanenburg, B. N., and Tanaka, Y. 1970, *Acta Phys. Suppl.*, **1**, 217.
 Bradt, H., Mayer, W., Narayan, S., Rappaport, S., and Spada, G. 1967, *Ap. J. (Letters)*, **150**, L199.
 Byram, E. T., Chubb, T. A., and Friedman, H. 1966, *Science*, **152**, 66.
 Clark, G. W., Garmire, G. P., and Kraushaar, W. L., 1968, *Ap. J. (Letters)* **153**, L203.
 Earl, J. A., Neely, D. E., and Rygg, T. A. 1970, Univ. Maryland Tech. Rept., No. 70-076.
 Ehrmann, C. H., Fichtel, C. E., Kniffen, D. A., and Ross, R. W. 1967, *Nucl. Instr. and Methods*, **56**, 109.
 Fanselow, J. L., Hartman, R. C., Hildebrand, R. H., and Meyer, P. 1969, *Ap. J.*, **158**, 771.
 Fazio, G., and Helmken, H. 1971, Smithsonian Institution Astrophysical Observatory Computer Programs, private communication.
 Felten, J. E. 1968, *Ap. J.*, **151**, 861.
 ———. 1970, *I.A.U. Symposium No. 37, Non-Solar X- and Gamma-Ray Astronomy*, ed. L. Gratton (Dordrecht: D. Reidel Publishing Co.), p. 216.
 Fichtel, C. E., Kniffen, D. A., and Ögelman, H. B. 1969, *Ap. J.*, **158**, 193.
 Fishman, G. J., Harnden, F. R., Jr., and Haymes, R. C. 1970, *I.A.U. Symposium No. 37, Non-Solar X- and Gamma-Ray Astronomy*, ed. L. Gratton (Dordrecht: D. Reidel Publishing Co), p. 116.
 Friedman, H., and Byram, E. T. 1967, *Science*, **158**, 257.
 Frye, G. M., Staib, J. A., Zych, A. D., Hopper, V. D., Rawlinson, W. R., and Thomas, J. A. 1969, *Nature*, **223**, 1320.
 Frye, G. M., and Wang, C. P. 1969, *Ap. J.*, **158**, 925.
 Garmire, G. P. 1970, *Bull. A.P.S.*, **15**, 564.
 Haymes, R. C., Ellis, D. V., Fishman, G. J., Kurfess, J. D., and Tucker, W. H. 1968, *Ap. J. (Letters)*, **151**, L131.
 Kellogg, E., Gursky, H., Leong, C., Schreier, E., Tananbaum, H., and Giacconi, R. 1971, *Ap. J. (Letters)*, **165**, L49.
 Kniffen, D. A., and Fichtel, C. E. 1970, *Ap. J. (Letters)*, **161**, L157.

40 C. FICHTEL, R. HARTMAN, D. KNIFFEN, AND M. SOMMER

- Marar, T. M. K., Freier, P. S., and Waddington, C. J. 1970, *Acta Phys. Suppl.*, **1**, 137.
- McClintock, J. E., Lewin, W. H. G., Sullivan, R. J., and Clark, G. W. 1969, *Nature*, **223**, 162.
- Nishimura, J., Mikumo, E., Mito, I., Niu, K., Ohta, I., and Taira, T. 1970, *Acta Phys. Suppl.*, **1**, 239.
- Peterson, L. E. 1970, *I.A.U. Symposium No. 37, Non-Solar X- and Gamma-Ray Astronomy*, ed. L. Gratton (Dordrecht: D. Reidel Publishing Co.), p. 59.
- Ross, R., Ehrmann, C., Fichtel, C., Kniffen, D., and Ögelman, H. 1969, *IEEE Trans. Nucl. Sci.*, **NS-16**, 127.
- Rockstroh, J., and Webber, W. R. 1969, *J. Geophys. Res.*, **74**, 5041.
- Sartori, L., and Morrison, P. 1967, *A p. J.*, **150**, 385.
- Stecker, F. W. 1971, *Cosmic Gamma Rays*, NASA SP-249 (Washington: Government Printing Office).
- Webber, W. R., and Reinert, C. P. 1970, *A p. J.*, **162**, 833.